NMR Imaging of Drought-induced Vessel Blockage and Restoration of Hydraulic Conductivity in Citrus and Blueberry Plants

C.R. Brodersen (U.F., Citrus Research & Education Center / Horticultural Science)

Introduction

The mechanics of water movement in plants is unusual because water is transported under a state of hydrostatic tension through vessels known as xylem. Under drought conditions this tension can become severe and lead to cavitation, resulting in trapped gas bubbles that block flow. The resulting loss of hydraulic conductivity deprives leaves of water, leading to wilting and ultimately death. Recently, NMR imaging has become a powerful tool for studying plant hydraulics *in vivo* because of its non-destructive nature. The goal of this study was to determine whether the functional status of citrus xylem vessels could be determined (i.e. water-filled and functional vs. air-filled and non-functional) using nuclear magnetic resonance imaging at the UF AMRIS facility, and whether citrus fruit development could be non-destructively studied.

Experimental

Stem and fruit samples from Hamlin Orange trees were collected at the UF-IFAS Citrus Research & Education Center in Lake Alfred, FL and sent overnight to the Gainesville campus. Samples were loaded into the 4.7T and 11T NMR imaging systems to establish which instrument and imaging parameters (e.g. gradient echo, spin echo, 2D vs. 3D, etc.) would provide sufficient images for this study.



Figure 1: Two dimensional NMR imaging of Hamlin Orange fruit in the xy, xz, and yz planes (A-C, respectively). Fruit is approximately 4cm in diameter. Three dimensional NMR image of an orange rendered with Avizo software (D). Transverse cross sections through citrus stems approximately 8mm in diameter.

Results and Discussion

The preliminary data generated using the 11T instrument provided the best resolution when comparing the two instruments (11T vs. 4.7T). The spin-echo parameters provided excellent detail of the Hamlin Orange, including the individual segments, juice sacks, oil glands in the peel, and seeds. The 2D images (**Figure 1A-C**) proved to be of sufficient quality that we will begin a season-long developmental study of citrus with and without “Citrus Greening” disease that causes fruit to become misshapen. Using Avizo 7.0 software we were able to visualize the Hamlin Orange in 3D (Figure 1D). The 11T instrument also provided the best images of stem cross sections (**Figure 1E-F**). Bright white spots in the images are the vessels filled with water. We are currently working with the AMRIS staff to optimize the scanning protocol, but the initial images are promising.

Conclusions

While this project was started late in the year (November, early-December) we have already made excellent progress in selecting the instruments and parameters necessary for completing the objectives of the study.

Acknowledgements

Dr. Brodersen would like to thank the NHMFL and AMRIS for the funding opportunity to support this pilot project (ML-BRODERSEN-001) and Huadong Zeng for his assistance with the initial imaging.