



## 1.1 Billion Years Old Porphyrins Establish a Marine Ecosystem Dominated by Bacterial Primary Producers

Gueneli, N. (The Australia National U.); McKenna, A.M. (NHMFL, Chemistry); Ohkouchi, N. (Japan Agency for Marine-Earth Science and Technology); Boreham, C.J. (Geoscience Australia); Beghin, J., Javaux, E.J. (U. of Liege) and Brocks, J.J. (The Australia National U.)


### Results and Discussion

Porphyrins are the molecular fossils of chlorophylls. Our recent discovery of porphyrins from 1.1 billion-year-old marine black shales of the Taoudeni Basin in Mauritania pushes back the geological record for photosynthesis by 600 million years. These porphyrins are nitrogen-containing molecules preserved from bacteriochlorophylls, and quantitation of their nitrogen isotopic composition provides quantitative information about the type of organisms in past ecosystems that utilized sunlight energy to synthesize organic compounds for nutrition.

Fourier Transform Ion Cyclotron Resonance (FT-ICR) mass spectrometry resolved and identified porphyrins from the tens of thousands of other organic compounds in each shale. Indeed, FT-ICR MS was able to identify the elemental composition and structure for nickel and vanadyl porphyrins in this highly-preserved shale.

The porphyrin nitrogen isotope ratio is different in this shale than in younger sediments. The isotopic offset between sediment nitrogen and the porphyrin nitrogen measured in this experiment indicates cyanobacteria were the dominant photosynthetic organisms 1.1 billion years ago in the Taoudeni Basin. The data indicate that purple sulphur bacteria and green sulfur bacteria also thrived, while larger planktonic algae were scarce.

The populations of bacteria through the mid-Proterozoic (1.8 to 0.8 billion years ago) and the perceived later transition to a more eukaryotic world - filled with large, active organisms - is one of the most intriguing yet unresolved phenomena of Precambrian biology.



**Figure 1: Top:** Identification of two classes of Ni- and VO-porphyrins by FT-ICR MS, plotting relative abundance versus double-bond equivalents (DBEs) and carbon number. **Left:** The structure of  $\text{C}_{30}$  to  $\text{C}_{32}$  DPEP, one of the many porphyrin structures identified supports phototrophs as dominant photosynthesizing organisms on Earth 1.1 billion years ago. Identified porphyrins likely derived from oxygenic phototrophs and anoxygenic phototrophic bacteria.

### Acknowledgements

The National High Magnetic Field Laboratory is supported by the National Science Foundation through NSF/DMR-1157490/1644779 and the State of Florida.

This work was supported by NSF Division of Materials Research through DMR-11-57490, the State of Florida, National Institutes of Health grants AI 033993 (DFH) and GM 037537 (DFH).

### References

[1] Gueneli, N., *et al.*, Proc. Natl. Acad. Sci. U.S.A., **115**, 1-9 (2018) DOI: 10.1073/pnas.1803866115