

LETTER TO THE EDITOR

Reply: Thalamotomy for tremor normalizes aberrant pre-therapeutic visual cortex functional connectivity

Jesse C. DeSimone,¹ Derek B. Archer,¹ David E. Vaillancourt^{1,2,3} and Aparna Wagle Shukla^{3,4}

1 Department of Applied Physiology and Kinesiology, University of Florida, Gainesville, FL, USA

2 Department of Biomedical Engineering, University of Florida, Gainesville, FL, USA

3 Department of Neurology, College of Medicine, University of Florida, Gainesville, FL, USA

4 Norman Fixel Institute for Neurological Diseases, College of Medicine, University of Florida, Gainesville, FL, USA

Correspondence to: Aparna Wagle Shukla

Department of Neurology, College of Medicine, University of Florida, Gainesville, FL, USA

E-mail: aparna.shukla@neurology.ufl.edu

Sir,

We thank Tuleasca and colleagues for their insightful comments on our study that compared the distinct neural network signatures identified with functional imaging analysis for dystonic tremor (DT) and essential tremor (ET). We chose to compare these tremor disorders as these entities are frequently confused in the clinical setting. In our experimental paradigm, we examined the effect of visual feedback on grip-force tremor and how that relates to blood oxygen level-dependent (BOLD) activity and functional connectivity in the two tremor disorders. We found increased visual feedback led to similar force tremor exacerbation; however, the functional activation abnormalities (termed $BOLD_{\Lambda}$) in the cortical and subcortical regions were different between dystonic and essential tremor. More importantly, the seed-based functional connectivity (termed FC_{Δ}) from the sensorimotor cortex/inferior parietal lobule, globus pallidus internus, ventral intermediate nucleus, and dentate nucleus in dystonic tremor was strikingly far more impaired within the cortical, subcortical and cerebellar regions in dystonic tremor compared to essential tremor (DeSimone et al., 2019). Our study highlights the importance of the visual network in essential tremor generation, which is consistent with previous studies by our group and other investigators (Tuleasca et al., 2017, 2018a; Archer et al., 2018; Benito-Leon et al., 2019). Additionally, our imaging analysis extends the involvement of a widespread visually sensitive functional network to dystonic tremor, which is a novel insight for

understanding the pathogenesis. While we are just beginning to understand the neural network signatures for dystonic tremor, future multimodal studies at a molecular, structural and functional level similar to those performed by Tuleasca *et al.* and other groups in essential tremor will further parse out the critical differences between these two disorders. Furthermore, the effects of therapeutic intervention such as deep brain stimulation (DBS) or thalamotomy procedures on tremor suppression and the corresponding change in the network behaviour similar to studies performed in essential tremor (Tuleasca *et al.*, 2017, 2019) will add clarification to their exact role in pathogenesis. Ultimately, these advances will contribute to the development of a robust diagnostic framework and focused treatments for the two disorders.

Dystonic tremor versus essential tremor

Functional perspective

A few years ago, the Movement Disorders Society officially designated tremor as a part and parcel symptom of dystonia (Albanese *et al.*, 2013). Dystonic tremor is defined as a tremor manifesting in a body part affected with dystonic features. Despite these efforts, owing to clinical resemblance dystonic tremor is frequently misdiagnosed as essential tremor. While behavioural studies have provided some clues for tremor distinction (DeSimone *et al.*, 2019), there is

© The Author(s) (2019). Published by Oxford University Press on behalf of the Guarantors of Brain. All rights reserved.

For permissions, please email: journals.permissions@oup.com

Advance Access publication October 11, 2019

limited knowledge on disorder-specific brain deficits that can reliably differentiate these two disorders. In our previous studies, we learned visuomotor processes influenced the amplitude of tremor and force error in essential tremor; the current study evidenced similar overlapping findings in the dystonic tremor cohort. We agree with Tuleasca and colleagues who draw attention in their letter to converging pieces of evidence in support of a widespread dysfunction of a visually sensitive functional network underlying tremor generation in essential tremor. What remains less understood is the detailed role of the visuomotor network in the pathogenesis of dystonic tremor.

We found the BOLD activity change in response to visual feedback was reduced in essential tremor compared to the dystonic tremor cohort when examined across many brain regions including the sensorimotor cortex, inferior parietal lobule, cingulate cortex, medial premotor cortex, and lingual gyrus (DeSimone et al., 2019). Dystonic tremor versus healthy control comparisons also revealed different activation patterns in many cortical regions of the brain, including middle and medial frontal gyri, cingulate gyrus, and cerebellar lobule I-IV. As pointed out in the letter, a recent resting state functional MRI study in a cohort of spasmodic dysphonia with and without tremor found that compared with healthy controls, there was abnormal hyperexcitability of the premotor-parietal-putaminal circuitry likely related to deficient connectivity and thus the aberrant flow of information between these regions (Battistella and Simonyan, 2019).

Visual network FC_{Δ} was impaired in dystonic tremor relative to controls for all seed locations, whereas essential tremor demonstrated reduced visual network FC_{Δ} for seed locations in the sensorimotor cortex/inferior parietal lobule and dentate nucleus. Furthermore, visual cortex FC_{Δ} was significantly reduced for the dystonic tremor group compared to the essential tremor group for seed locations in the sensorimotor cortex/inferior parietal lobule, ventral intermediate nucleus, and dentate nucleus. In line with our task-based functional MRI analysis, Tuleasca et al. (2019), report resting state functional MRI findings in essential tremor cohort analysed before and after patients receive thalamotomy surgery. The authors used a co-activation pattern analysis to understand how a specific seed region in the brain interacts in a time-varying manner with the rest of the brain. They found the co-activation patterns involving the visual network, particularly the extrastriate visual areas normalize when the tremor severity is controlled with thalamotomy surgery. The current treatment interventions for dystonic tremor are noted to significantly overlap with essential tremor, e.g. DBS and focused ultrasound thalamotomy targeted to the ventral intermediate nucleus (Fasano et al., 2017), but what remains to be seen is the short-term and long-term relationship between tremor control and the pathogenic networks.

Structural perspective

While our study focused on neural signatures from a functional perspective, Tuleasca *et al.*, in their previous work involving essential tremor, found structural changes in the visual regions of the brain. They implemented voxel-based morphometry analysis to track the longitudinal effects of ventral intermediate nucleus radiosurgery on the grey matter volume (Tuleasca *et al.*, 2017). They found higher baseline grey matter volumes in Brodmann area 19 (visual cortex region) were associated with higher improvements in tremor scores. In another voxel-based morphometry study, they found that a decrease in grey matter density at 1 year in the occipital cortex after ventral intermediate nucleus radiosurgery was related to improvements that are more significant in the hand tremor score (Tuleasca *et al.*, 2018*b*).

For a long time dystonia was regarded as a disorder purely affecting brain function. With the advent of voxelbased morphometry, many studies in dystonia uncovered distinct structural abnormalities in the grey matter of the cortical and subcortical regions (Lehericy *et al.*, 2013). Furthermore, diffusion tensor imaging has shown changes in the white matter fibre tracts in many forms of dystonia (Colosimo *et al.*, 2005). We agree our study does not address the structural changes in dystonic tremor; however, a well designed imaging study combining structural and functional aspects of the pathogenic networks with parallel monitoring of treatment response is the next promising step to pursue.

Data availability

Data sharing is not applicable to this article as no new data were created or analysed in this study.

Acknowledgements

A portion of this work was performed in the McKnight Brain Institute at the National High Magnetic Field Laboratory's AMRIS Facility.

Funding

This work was supported by grants from the National Institutes of Health (R01 NS058487, K23 NS092957) and Tyler's Hope Foundation. The National High Magnetic Field Laboratory's AMRIS Facility, which is supported by National Science Foundation Cooperative Agreement No. DMR-1644779 and the State of Florida.

Competing interests

D.A. received grant support from the Parkinson's Foundation. D.V. received grant support from NIH, NSF, and Tyler's Hope Foundation. He is co-founder and manager of Neuroimaging Solutions, LLC. A.W.S. received grant support from Benign Essential Blepharospasm Research

Foundation, Dystonia Coalition, Dystonia Medical Research Foundation, and National Organization for Rare Disorders and grant support from NIH (KL2 and K23 NS092957-01A1).

References

- Albanese A, Bhatia K, Bressman SB, Delong MR, Fahn S, Fung VS, et al. Phenomenology and classification of dystonia: a consensus update. Mov Disord 2013; 28: 863–73.
- Archer DB, Coombes SA, Chu WT, Chung JW, Burciu RG, Okun MS, et al. A widespread visually-sensitive functional network relates to symptoms in essential tremor. Brain 2018; 141: 472–85.
- Battistella G, Simonyan K. Top-down alteration of functional connectivity within the sensorimotor network in focal dystonia. Neurology 2019; 92: e1843–e51.
- Benito-Leon J, Serrano JI, Louis ED, Holobar A, Romero JP, Povalej-Brzan P, et al. Essential tremor severity and anatomical changes in brain areas controlling movement sequencing. Ann Clin Transl Neurol 2019; 6: 83–97.
- Colosimo C, Pantano P, Calistri V, Totaro P, Fabbrini G, Berardelli A. Diffusion tensor imaging in primary cervical dystonia. J Neurol Neurosurg Psychiatry 2005; 76: 1591–3.

- DeSimone JC, Archer DB, Vaillancourt DE, Wagle Shukla A. Network-level connectivity is a critical feature distinguishing dystonic tremor and essential tremor. Brain 2019; 142: 1644–59.
- Fasano A, Llinas M, Munhoz RP, Hlasny E, Kucharczyk W, Lozano AM. MRI-guided focused ultrasound thalamotomy in non-ET tremor syndromes. Neurology 2017; 89: 771–5.
- Lehericy S, Tijssen MA, Vidailhet M, Kaji R, Meunier S. The anatomical basis of dystonia: current view using neuroimaging. Mov Disord 2013; 28: 944–57.
- Tuleasca C, Bolton TAW, Regis J, Najdenovska E, Witjas T, Girard N, et al. Normalization of aberrant pretherapeutic dynamic functional connectivity of extrastriate visual system in patients who underwent thalamotomy with stereotactic radiosurgery for essential tremor: a resting-state functional MRI study. J Neurosurg 2019; 10: 1–10 doi: 10.3171/2019.2.JNS183454.
- Tuleasca C, Regis J, Najdenovska E, Witjas T, Girard N, Thiran JP, et al. Visually-sensitive networks in essential tremor: evidence from structural and functional imaging. Brain 2018a; 141: e47.
- Tuleasca C, Witjas T, Najdenovska E, Verger A, Girard N, Champoudry J, et al. Assessing the clinical outcome of Vim radiosurgery with voxel-based morphometry: visual areas are linked with tremor arrest! Acta Neurochir 2017; 159: 2139–44.
- Tuleasca C, Witjas T, Van de Ville D, Najdenovska E, Verger A, Girard N, et al. Right Brodmann area 18 predicts tremor arrest after Vim radiosurgery: a voxel-based morphometry study. Acta Neurochir 2018b; 160: 603–9.