

Adiabatic Passage through Level Anticrossings in Systems of Chemically Inequivalent Protons Incorporating Parahydrogen: Theory, Experiment, and Prospective Applications

Maria-Jose Ferrer, Erin L. Kuker, Evgeniya Semenova, Anghelo Josh Gangano, Michelle P. Lapak, Alexander J. Grenning, Vy M. Dong, and Clifford R. Bowers*



Cite This: *J. Am. Chem. Soc.* 2022, 144, 20847–20853



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Level anticrossings (LACs) are ubiquitous in quantum systems and have been exploited for spin-order transfer in hyperpolarized nuclear magnetic resonance spectroscopy. This paper examines the manifestations of adiabatic passage through a specific type of LAC found in homonuclear systems of chemically inequivalent coupled protons incorporating parahydrogen (pH_2). Adiabatic passage through such a LAC is shown to elicit translation of the pH_2 spin order. As an example, with prospective applications in biomedicine, proton spin polarizations of at least $19.8 \pm 2.6\%$ on the methylene protons and $68.7 \pm 0.5\%$ on the vinylic protons of selectively deuterated allyl pyruvate ester are demonstrated experimentally. After ultrasonic spray injection of a precursor solution containing propargyl pyruvate and a dissolved Rh catalyst into a chamber pressurized with 99% para-enriched H_2 , the products are collected and transported to a high magnetic field for NMR detection. The LAC-mediated hyperpolarization of the methylene protons is significant because of the stronger spin coupling to the pyruvate carbonyl ^{13}C , setting up an ideal initial condition for subsequent coherence transfer by selective INEPT. Furthermore, the selective deuteration of the propargyl side arm increases the efficiency and polarization level. LAC-mediated translation of parahydrogen spin order completes the first step toward a new and highly efficient route for the ^{13}C NMR signal enhancement of pyruvate via side-arm hydrogenation with parahydrogen.



INTRODUCTION

Avoided crossings of eigenvalue curves, or level anticrossings (LACs), are ubiquitous in quantum systems and underpin diverse physical phenomena.^{1–5} For instance, LACs of spin states dressed by interaction with resonant radiofrequency (RF) fields provide a means to manipulate singlet-triplet imbalances,⁶ and LACs induced by heteronuclear spin couplings have been extensively exploited for conversion of parahydrogen (pH_2) singlet order to ^{13}C magnetization at ultralow magnetic fields,^{7–14} thereby extending the applicability of parahydrogen-enhanced NMR.^{15–18} Apparently unnoticed until now is the significance of LACs in systems of chemically inequivalent protons incorporating pH_2 spin order. It will be shown here that adiabatic passage through such LACs can elicit translation of bilinear spin order to initially unpolarized protons in the molecule with an efficiency that depends on the spin couplings and relative proton isotropic chemical shifts. This quantum effect will enable a new and practical route for spin order transfer to ^{13}C for the NMR signal enhancement of pyruvate and related pH_2 adducts with applications in biomedicine.^{18–21}

THEORY

The essential aspects of the spin dynamics are illustrated in a three-proton AMX spin system (Pople Notation²²) formed by hydrogenation with pH_2 , such as in the catalytic hydrogenation of a selectively deuterated propargyl ester to the allyl ester, as shown in Scheme 1.


Protons H^1 and H^2 are assumed to originate from pH_2 , and H^3 is an initially unpolarized proton, e.g., the allyl methylene proton. The importance of the two different R groups in Scheme 1 derives from the perturbation on the isotropic chemical shifts δ_i of the three protons: for $\text{R} = \text{D}$, $\delta_2 > \delta_1 > \delta_3$, while for $\text{R} = \text{CD}_3$, $\delta_2 > \delta_3 > \delta_1$. For the present purposes, the spin couplings to ^2H and nuclei in the R' group are unimportant and will be ignored. The isotropic spin Hamiltonian pertinent to nonviscous solutions has the form

Received: August 23, 2022

Published: November 4, 2022



Scheme 1. Hydrogenation of Selectively Deuterated Propargyl Esters with Parahydrogen



$$\hat{H}(B) = \sum_{i < j}^3 J_{ij} \hat{\mathbf{I}}_i \cdot \hat{\mathbf{I}}_j + \sum_{i=1}^3 \gamma B(1 - \delta_i) \hat{I}_{zi} \quad (1)$$

where γ is the proton gyromagnetic ratio, B is the applied static magnetic field, \hat{I}_{zi} is the z -component of the i th spin operator $\hat{\mathbf{I}}_i$, and $J_{ij} \hat{\mathbf{I}}_i \cdot \hat{\mathbf{I}}_j$ is the isotropic spin–spin coupling between protons H^i and H^j . For the analytical treatment, we assume idealized Hamiltonians with

$$|J_{12}| \gg |J_{23}| \text{ and } J_{13} = 0$$

and we consider three different constructs for an AMX spin system, designated C1, C2, and C3, defined in **Table 1**. The three constructs differ only in the relative order of the isotropic chemical shifts.

Table 1. Density Operators Resulting from Adiabatic Passage from Strong to Weak Coupling of Idealized AMX Spin Systems, Assuming $J_{12} \gg J_{23}$ and $J_{13} = 0$

construct	Hamiltonian parameters	LAC?	final density operator, $\hat{\rho}_f$
C1	$\delta_2 > \delta_1 > \delta_3$ or $\delta_3 > \delta_1 > \delta_2$	YES	$\frac{1}{8} - \frac{1}{2} \hat{I}_{z2} \hat{I}_{z3} \pm \frac{1}{4} (\hat{I}_{z2} - \hat{I}_{z3})$
C2	$\delta_1 > \delta_2 > \delta_3$ or $\delta_3 > \delta_2 > \delta_1$	YES	$\frac{1}{8} - \frac{1}{2} \hat{I}_{z1} \hat{I}_{z3} \pm \frac{1}{4} (\hat{I}_{z1} - \hat{I}_{z3})$
C3	$\delta_2 > \delta_3 > \delta_1$ or $\delta_1 > \delta_3 > \delta_2$	NO	$\frac{1}{8} - \frac{1}{2} \hat{I}_{z1} \hat{I}_{z2} \pm \frac{1}{4} (\hat{I}_{z1} - \hat{I}_{z2})$

At the instant of hydrogenation, the projection operator for the pH_2 singlet state is

$$\hat{\rho}_s = \frac{1}{4} (|S_0\alpha\rangle\langle S_0\alpha| + |S_0\beta\rangle\langle S_0\beta|) = \frac{1}{2} \left(\frac{1}{4} - \hat{\mathbf{I}}_1 \cdot \hat{\mathbf{I}}_2 \right) \quad (2)$$

If $|J_{12}| \gg |J_{23}|$, then at a sufficiently low magnetic field B_{low} , the commutator $[\hat{H}(B_{low}), \hat{\rho}_s] \approx 0$, the singlet order will remain essentially intact after averaging over the kinetic time distribution of individual hydrogenation events, and the time-averaged density operator for the ensemble of hydrogenation adducts will be $\langle \hat{\rho}_s \rangle \approx \hat{\rho}_s$. In the ALTADENA (Adiabatic Longitudinal Transport After Dissociation Engenders Net Alignment) effect,²³ the hydrogenation adducts formed at $B_{low} \approx 0$ are adiabatically transported to a high magnetic field where weak coupling prevails and the NMR spectrum is acquired. Adiabatic passage, where individual spin states evolve without a change in their occupancy, occurs when the change in the spin Hamiltonian is sufficiently slow to avoid stimulating transitions between eigenstates yet fast enough to avert significant spin relaxation.^{5,13} Under such conditions, the density operator resulting from adiabatic transport over a time τ_{tr} that is short compared to the spin–lattice relaxation times can be approximated by unitary evolution:

$$\hat{\rho}_f(\tau_{tr}) = \hat{U}(\tau_{tr}) \langle \hat{\rho}_s \rangle \hat{U}^{-1}(\tau_{tr}) \quad (3)$$

where $\hat{U}(\tau_{tr}) = \exp(-i \int_0^{\tau_{tr}} \hat{H}(t') dt')$.

In the arrow notation, this transformation is expressed as $\langle \hat{\rho}_s \rangle \xrightarrow{\hat{H}(B)} \hat{\rho}_f$. In the trivial case where $J_{23} \rightarrow 0$ and $J_{13} \rightarrow 0$, H^3 is isolated, and the density operator $\hat{\rho}_f$ is like that of the original ALTADENA effect for two weakly coupled protons:²³

$$\left(\frac{1}{4} - \hat{\mathbf{I}}_1 \cdot \hat{\mathbf{I}}_2 \right) \otimes \frac{1}{2} \hat{I} \xrightarrow{\text{adiabatic}} \frac{1}{8} - \frac{1}{2} \hat{I}_{z1} \hat{I}_{z2} \pm \frac{1}{4} (\hat{I}_{z1} - \hat{I}_{z2}) \quad (4)$$

The sign of the third term in **eq 4** depends on the relative signs of J_{12} and $\delta_1 - \delta_2$.

Rotating frame eigenvalue correlation diagrams for the C1- and C3-type constructs are shown at the top of **Figure 1**. A LAC is apparent for C1, but not for C3 where δ_3 is bracketed by δ_1 and δ_2 . The initial and final state populations resulting from adiabatic passage are indicated by the blue disks, and the hyperpolarized single-quantum NMR transitions involving each of the three protons in the weak-coupling regime are indicated by the color-coded arrows. Because there is no LAC for C3, the transitions of H^3 are unpolarized. In contrast, when δ_3 lies outside the range spanned by δ_1 and δ_2 (either above or below) as in C1 or C2, the transitions of H^3 are hyperpolarized (burgundy dashed arrows).

The corresponding analytical calculations of the final density operator $\hat{\rho}_f$ resulting from unitary adiabatic propagation from the strong to weak coupling regimes for the idealized constructs C1, C2, and C3, according to **eq 3**, are reported in **Table 1**.

For C1 and C2, the final density operators are identical in form to **eq 4** but with proton labels 1 and 3 or 2 and 3 interchanged, respectively. For C3, $\hat{\rho}_f$ is the same as in **eq 4**. The quantity $\langle \hat{I}_{z3} \rangle = Tr(\hat{I}_{z3} \cdot \hat{\rho}_f)$ provides a measure of the efficiency of the conversion of scalar pH_2 spin order (**eq 1**) into H^3 polarization. As seen in **Tables 1** and **2**, adiabatic passage through the LAC results in spin exchange involving both the linear and bilinear terms of $\hat{\rho}_s$ leading to the complete transfer of parahydrogen-derived spin order to the initially unpolarized proton in the molecule. The density operator calculations confirm that when the chemical shift of H^3 is intermediate to those of H^1 and H^2 , H^3 remains unpolarized, irrespective of J_{23} .

Let us now consider the implications of the LAC-mediated proton spin exchange of parahydrogen spin order for producing ^{13}C hyperpolarization in pyruvate and other carboxylic acids. While pyruvate is not directly producible by hydrogenation, attachment of the propargyl side arm provides an unsaturated bond for addition of pH_2 ,²⁴ as in **Scheme 1**. In this side-arm hydrogenation (SAH) methodology, the pH_2 spin order needs to be transferred across the ester linkage to the carbonyl ^{13}C , and then the side arm is removed by hydrolysis to afford pyruvate. Propargylic side arms offer greater chemical stability and higher rates of hydrogenation^{7,25} than vinylic side arms. Disadvantageously, the pH_2 proton pair in allyl ester adducts is further away from the ^{13}C target due to the intervening CH_2 group in the former, and the spin–spin couplings to the nearest pyruvate ^{13}C are too small to mediate efficient coherence transfer in a single step.²⁷ This motivated the development of various two-step relayed coherence transfer pulse sequences^{25,28–30} that operate on PASADENA-derived $\hat{I}_{z1} \hat{I}_{z2}$ spin order generated at a high magnetic field.^{15,16} Advantageously, LAC-mediated translation to the H^3 (methylene) proton(s) in allyl ester side arms of type C1 is significant because of the stronger spin coupling to the




Figure 1. Rotating frame eigenvalue correlation diagrams for model Hamiltonians of types C1 and C3 as a function of the static magnetic field with $\bar{J}_{12} = 25$ Hz and $\bar{J}_{23} = 5$ Hz (upper diagrams) or $\bar{J}_{23} = 0$ Hz (lower diagrams). (a) C1, $\{\delta_1, \delta_2, \delta_3\} = \{0, +10, -10\}$ ppm. (b) C3, $\{\delta_1, \delta_2, \delta_3\} = \{+10, -10, 0\}$ ppm. The LAC is seen in the dashed red box. Orange, blue, and burgundy arrows indicate polarized single spin transitions of H^1 , H^2 , and H^3 , respectively. In the absence of a LAC, transitions of H^3 remain unpolarized. The correlation diagram for C2 is analogous to C1. Curves were calculated using the SpinDynamica package²⁶ in Wolfram's Mathematica.

Table 2. Zeeman Order Produced by Adiabatic Passage from Strong to Weak Coupling for Idealized Constructs

Hamiltonian	$J_{23} = 0$			$J_{23} > 0$		
Chemical Shifts	$\langle I_{z1} \rangle$	$\langle I_{z2} \rangle$	$\langle I_{z3} \rangle$	$\langle I_{z1} \rangle$	$\langle I_{z2} \rangle$	$\langle I_{z3} \rangle$
C1 $\delta_2 > \delta_1 > \delta_3$	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$
C3 $\delta_2 > \delta_3 > \delta_1$	$\frac{1}{2}$	$-\frac{1}{2}$	0	$\frac{1}{2}$	$-\frac{1}{2}$	0
C4 $\delta_1 = \delta_2$ $\neq \delta_3 = \delta_4$	0	0	0	$\frac{1}{4}$	$\frac{1}{4}$	$-\frac{1}{2}$


$\langle I_{z3} \rangle + \langle I_{z4} \rangle$.

pyruvate carbonyl ^{13}C , setting up an ideal initial condition for subsequent coherence transfer to the carbonyl ^{13}C by selective INEPT.³¹

The spin polarizations after adiabatic transport were evaluated for several pyruvate esters of interest (Scheme 2) by numerical density matrix simulations, and these are reported in Table 3. First, the results confirm the importance of the relative ordering of the chemical shifts (and hence the Zeeman energies) of the two pH_2 -sourced protons and the relay proton(s). While (*Z*)-allyl-3-*d* 2-oxopropanoate-1- ^{13}C (APd) is a type C1 molecule, 3-*d*₃-1-*d*-buten-2-yl (BPd₄)³² has the chemical shift ordering of C3 and is thus predicted to not exhibit a LAC. Even though APd and BPd₄ differ only in the attachment of a $-\text{CD}_3$ group at the methylene carbon, the methylation has the effect of changing the chemical shift ordering, thus switching off the LAC-mediated spin exchange process.

Second, the results in Table 3 show that selective deuteration can increase the efficiency of symmetry breaking conversion of

Scheme 2. Molecular Embodiments under Study^a



^aAP = allyl 2-oxopropanoate-1- ^{13}C ; APd = (*Z*)-allyl-3-*d* 2-oxopropanoate-1- ^{13}C ; APd₂ = (*Z*)-allyl-1,3-*d* 2-oxopropanoate-1- ^{13}C ; BPd₄ = 3-*d*₃-1-*d*-buten-2-yl; DMM = dimethyl maleate.

Table 3. Numerical Simulations of Zeeman Order Produced in the Molecular Embodiments

Construct	Molecule	$\langle I_{z1} \rangle$	$\langle I_{z2} \rangle$	$\langle I_{z3} \rangle$
C1	AP	-0.095	+0.28	-0.27 ^a
	APd, APd ₂	-0.0028	+0.42	-0.42 ^a
C3	BPd ₄	-0.46	+0.46	0.0024
C4	DMM	0.24	0.24	-0.48 ^b

^a $\sum \langle I_z \rangle$ over both CH₂ relay protons in AP and APd. ^b $\sum \langle I_z \rangle$ over all methyl protons.

pH₂ spin order into NMR-observable Zeeman order.^{7,29} In nondeuterated AP or BP, *J* couplings to H^s incur substantial irreversible loss of pH₂ spin order upon hydrogenation. Because $J_{25} > J_{12}$ and J_{23} , there is a large admixture of the H^s spin state in the low field eigenstates of the adduct. The destructive effect of J_{25} in the nondeuterated form is evident by evaluating $\text{Tr}\{\hat{\rho}_s \cdot \langle \hat{\rho}_s \rangle\}$. Recall that for a pure state, $\text{Tr}\{\hat{\rho}^2\} = 1$, but since H³ and H^s are initially unpolarized, $\text{Tr}\{\hat{\rho}^2\} = 1/4$ maximally. If




Figure 2. (a) Block diagram of the ultrasonic spray injection reactor system interfaced to the flow NMR spectrometer. The tubing connecting the withdrawal syringe to the reaction chamber outlet, including the $60 \mu\text{L}$ flow cell, is initially pre-filled with solvent, and the reaction chamber is pressurized to 6 bars with 99% pH_2 . The precursor solution is infused through an ultrasonic nozzle (3.5 W, 120 kHz) into the chamber at a flow rate of 5 mL/min. Products accumulate at the bottom of the funnel-shaped chamber and are drawn into the NMR probe at precisely controlled syringe pump flow rates. BPR = back-pressure regulator (5.17 bars). Further details are provided in the Supporting Information. (b) Numerical density matrix calculations of the spin polarization (neglecting spin relaxation) of individual protons in APd_2 as a function of the syringe pump withdrawal rate, estimated from the transport time τ_{tr} used in the simulation, the volume (0.33 mL), and the measured field profile of the Bruker 9.4 T Ultrashield magnet along the flow trajectory. Spin polarizations were computed using Spinach.³²




Figure 3. (a, b) 400 MHz ${}^1\text{H}$ ALTADENA experimental spectra (in red) and numerically simulated ${}^1\text{H}$ ALTADENA spectra (in blue) of AP and APd, respectively, obtained by hydrogenation of 8 mM propargyl pyruvate precursors with pH_2 using 2 mM Rh catalyst in the 2.4 mT fringe field followed by transport into the flow probe at a flow rate of 3 mL/min for detection at 9.4 T using a 90° RF pulse. The spectra have been normalized to the H^2 peak. (c) 400 MHz ${}^1\text{H}$ ALTADENA spectra of AP (in blue) and APd (in red) obtained by hydrogenation as in panels (a) and (b) but with 40 mM propargyl pyruvate and 10 mM Rh catalyst. The H^2 peak in the spectrum of AP has been scaled to match the H^2 peak of the APd spectrum. (d) Thermally polarized spectra of the reaction product solutions of panel (c), acquired by accumulation of four transients and plotted on the same vertical axis as panel (c).

$J_{1S} \rightarrow 0$, one obtains $\text{Tr}\{\hat{\rho}_s \cdot \langle \hat{\rho}_s \rangle\} = 0.21$, but if $\bar{J}_{1S} = 17.2$ Hz (as in AP), $\text{Tr}\{\hat{\rho}_s \cdot \langle \hat{\rho}_s \rangle\} = 0.122$. The loss can be mitigated by selective deuteration at H⁵, since $J_{\text{HD}}/J_{\text{HH}} \approx 1/7$.

Moreover, if hydrogenation is performed in a small but finite magnetic field (e.g., Earth's field) where deuterons and protons are only weakly coupled, losses due to state mixing are negligible. The numerical simulation results in **Table 3**, calculated assuming perfect adiabatic passage, show that deuteration at H⁵ affords an ~50% increase of $\langle I_{z3} \rangle$ after adiabatic passage through the LAC. The theoretical maximum achievable polarization of H³, neglecting spin relaxation, is 84% in APd₂ (in APd, the polarization is divided equally between the two methylene protons). Simulation details and the MATLAB/Spinach source code are provided in the *Supporting Information*.

EXPERIMENTAL SECTION

The LAC-mediated proton spin exchange in AP and APd (after their formation by Rh-catalyzed hydrogenation with pH₂ at 2.4 mT) was studied with the apparatus diagrammed in **Figure 2a**. A d₆-acetone solution of the alkyne precursor and dissolved Rh catalyst was infused through an ultrasonically vibrated nozzle into a funnel-shaped reaction chamber pressurized to 6 bars with 99% enriched pH₂. Experiments were performed with a fixed 4:1 substrate:catalyst ratio with either 2 mM or 10 mM Rh(cod)(dppb)BF₄ catalyst concentration. Immediately after collecting 0.5 mL of the pH₂ adduct solution, the liquid was drawn into the Varian 400 MHz flow NMR probe at precisely controlled withdrawal rates varying from 0.2 to 5 mL/min, corresponding to transport times of $\tau_{\text{tr}} = 100$ to 4 s, respectively, using a Chemixy 4000 dual syringe pump. The high-resolution ¹H NMR spectra were acquired on a Varian VNMRS spectrometer with a 2.25 μ s 90° flip-angle pulse. This choice of flip angle suppresses the bilinear operator terms in $\hat{\rho}_f$ as well as the possible PASADENA signals stemming from any ongoing hydrogenation after transport to a high magnetic field. Complete experimental details, including the synthesis of the isotopically labeled propargyl pyruvate and acetate substrates, are provided in the *Supporting Information*.

Figure 2b plots the numerically simulated spin polarization, $P_i = 2\langle \hat{I}_{zi} \rangle$, of each proton in APd₂ as a function of flow rate. These simulations indicate that in our system, we can expect to approach the adiabatic regime at flow rates slower than about 2 mL/min ($\tau_{\text{tr}} > 10$ s), which is easily accessible by our syringe pump, probe, and fluid handling system. The polarization of the H² proton is expected to be invariant of the flow rate. However, it is important to note that these simulations do not account for the spin–lattice relaxation during transport. High field inversion-recovery T_1 measurements on thermally polarized APd reaction products (with a catalyst present) indicate that the H³ T_1 is significantly shorter at 6.4 s compared to the 9.2 s and 9.6 s relaxation times of the H¹ or H² protons, respectively (see the *Supporting Information*).

RESULTS AND DISCUSSION

The 400 MHz ¹H ALTADENA NMR spectra of AP and APd acquired after ultrasonic spray injection of the propargyl pyruvate precursor solutions containing 2 mM Rh catalyst into the reaction chamber pressurized with 99% pH₂ followed by infusion into the NMR flow probe at 3 mL/min ($\tau_{\text{tr}} = 6.6$ s) are presented in **Figure 3a,b**. Superimposed on these spectra are the corresponding numerical simulations based on the measured magnetic field profile of the 9.4 T Bruker Ultrashield magnet. Each spectrum has been normalized to its H² peak amplitude. Without any fitting parameters, excellent agreement between theory and experiment is obtained except for the H³ resonance, which is noticeably lower in intensity than in the simulated spectra of both AP and APd. The discrepancy is attributed to the shorter spin–lattice relaxation time of the H³ methylene protons

(see the *Supporting Information*). The effects of differential spin–lattice relaxation losses during transport become even more pronounced at lower flow rates (see the *Supporting Information*). **Figure 3c** presents the spectrum obtained using the same flow rate of 3 mL/min but with a 5× more concentrated precursor solution (10 mM Rh and 40 mM AP or APd). For APd, a LAC-mediated polarization of $P_3 = 19.8 \pm 2.6\%$ was obtained for the initially unpolarized relay (methylene) protons, while a remarkable $P_2 = 68.7 \pm 0.5\%$ polarization (averaged over three trials) was obtained for the vinylic H² proton originating from pH₂. Interestingly, the H¹ signal in the spectrum of APd obtained using the more concentrated precursor solution exhibits a clear antiphase doublet that defied numerical simulation. This contrasts with the net emission doublet observed with the APd precursor solution of lower concentration (**Figure 3b**), consistent with theory. Signal enhancement and polarization values were calculated using the signals in the thermally polarized spectra shown in **Figure 3d**, and these are reported in **Table S1** of the *Supporting Information*.

Closely related to the LAC-mediated spin exchange in AMX spin systems is the net alignment spin order produced in C_s-symmetric AA'XX' adducts of pH₂. The archetypical example is dimethyl maleate (DMM), a system addressed in several previous accounts^{33,34} prior to the understanding of the role of LACs in ALTADENA experiments.³⁵ For completeness, the analytical form of the density operator for the associated idealized construct C4, defined in **Table 2**, and a numerical spectral simulation for the DMM embodiment are provided in the *Supporting Information*.

CONCLUSIONS

In summary, parahydrogen-induced hyperpolarization levels of up to 19.8% (and 68.7% on the H² proton) on the methylene relay protons (H³) of (Z)-allyl-3-d 2-oxopropanoate-1-¹³C (APd) were demonstrated using the LAC-mediated proton spin exchange method introduced herein. Such high polarization levels are remarkable considering the relatively short relaxation time of these protons ($T_1 = 6.4$ s). Even higher polarization levels might be achieved by reducing T_1 losses during adiabatic transport as may be realized using a variable syringe pump flow rate to maintain the constant-adiabaticity¹³ criterion or by employing a bench-top NMR spectrometer with a reduced flow path length. A further increase in the H³ proton polarization level is anticipated in (Z)-allyl-1,3-d 2-oxopropanoate-1-¹³C (APd₂) adducts where the spin–lattice relaxation time is likely to be increased by replacement of the –CH₂– with –CHD– on the allyl side arm. An important implication of these results is that the LAC-mediated proton spin exchange of pH₂ spin order in the AMX spin systems completes the first step toward spin order transfer to ¹³C in pyruvate and other key metabolites. The deuteration pattern in APd₂ also lends itself to a higher efficiency (by a factor of 2) in the selective coherence transfer from H³ to the pyruvate carbonyl ¹³C,^{7,36,37} providing an overall maximum theoretical efficiency of 84% for the conversion of pH₂ spin order to ¹³C magnetization on the metabolite. This is similar to the theoretical maximum efficiency of the high-field relayed-coherence transfer RF pulse sequences based on high-field pH₂ addition to propargyl side arms as well as the slower reacting and less stable vinyl ester side arms (which lack relay protons). Advantageously, the proposed LAC-INEPT approach requires only a single coherence transfer step, thereby reducing T_2 decoherence losses. Moreover, the approach is compatible

with hydrogenation in a high-performance hydrogenation reactor (such as our ultrasonic spray injection system) operating outside of the NMR magnet. Polarization levels achievable by our LAC-INEPT approach are competitive with the best results obtained by dissolution DNP^{19,38} but with substantially lower cost, higher reproducibility, shorter polarization times, and perhaps most importantly, compatibility with continuous-flow chemistry and separations.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.2c09000>.

Details of the experimental instrumentation, protocols, synthesis and characterization of isotopically labeled compounds, equations used to evaluate signal enhancements and spin polarizations, spin interaction parameters for the studied molecules, analytical derivation details, Mathematica/SpinDynamica source code, Spinach/Matlab source code, and additional spectral simulations ([PDF](#))

■ AUTHOR INFORMATION

Corresponding Author

Clifford R. Bowers — Department of Chemistry, University of Florida, Gainesville, Florida 32611-7200, United States;
National High Magnetic Field Lab, Tallahassee, Florida 32310, United States;  orcid.org/0000-0001-6155-5163; Email: bowers@chem.ufl.edu

Authors

Maria-Jose Ferrer — Department of Chemistry, University of Florida, Gainesville, Florida 32611-7200, United States;  orcid.org/0000-0002-8971-8849

Erin L. Kuker — Department of Chemistry, University of California, Irvine, California 92697-2025, United States

Evgeniya Semenova — Department of Chemistry, University of Florida, Gainesville, Florida 32611-7200, United States

Anghelo Josh Gangano — Department of Chemistry, University of Florida, Gainesville, Florida 32611-7200, United States;  orcid.org/0000-0002-5577-5503

Michelle P. Lapak — Department of Chemistry, University of Florida, Gainesville, Florida 32611-7200, United States;  orcid.org/0000-0001-7809-3276

Alexander J. Grenning — Department of Chemistry, University of Florida, Gainesville, Florida 32611-7200, United States;  orcid.org/0000-0002-8182-9464

Vy M. Dong — Department of Chemistry, University of California, Irvine, California 92697-2025, United States;  orcid.org/0000-0002-8099-1048

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/jacs.2c09000>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by NSF grants CHE-2108306 and CBET-1933723 and the National High Magnetic Field Laboratory User Collaborative Grants Program, which is supported by NSF DMR-1644779 and the State of Florida.

A.G. and V.M.D. thank the National Institutes of Health, grants R35GM137893-01 and R35GM127071, respectively.

■ REFERENCES

- (1) Zhou, J.; Huang, P.; Zhang, Q.; Wang, Z.; Tan, T.; Xu, X.; Shi, F.; Rong, X.; Ashhab, S.; Du, J. Observation of Time-Domain Rabi Oscillations in the Landau-Zener Regime with a Single Electronic Spin. *Phys. Rev. Lett.* **2014**, *112*, No. 010503.
- (2) Rubbmark, J. R.; Kash, M. M.; Littman, M. G.; Kleppner, D. Dynamical Effects at Avoided Level Crossings: A Study of the Landau-Zener Effect Using Rydberg Atoms. *Phys. Rev. A* **1981**, *23*, 3107–3117.
- (3) Zener, C. Non-Adiabatic Crossing of Energy Levels. *Proc. R. Soc. London, Ser. A* **1932**, *137*, 696–702.
- (4) Corzilius, B. High-Field Dynamic Nuclear Polarization. *Annu. Rev. Phys. Chem.* **2020**, *71*, 143–170.
- (5) Ivanov, K. L.; Pravdivtsev, A. N.; Yurkovskaya, A. V.; Vieth, H. M.; Kaptein, R. The Role of Level Anti-Crossings in Nuclear Spin Hyperpolarization. *Prog. Nucl. Magn. Reson. Spectrosc.* **2014**, *81*, 1–36.
- (6) DeVience, S. J.; Walsworth, R. L.; Rosen, M. S. Preparation of Nuclear Spin Singlet States Using Spin-Lock Induced Crossing. *Phys. Rev. Lett.* **2013**, *111*, No. 173002.
- (7) Marshall, A.; Salhov, A.; Gierse, M.; Müller, C.; Keim, M.; Lucas, S.; Parker, A.; Scheuer, J.; Vassiliou, C.; Neumann, P.; Jelezko, F.; Retzker, A.; Blanchard, J. W.; Schwartz, I.; Knecht, S. Radio-Frequency Sweeps at μT Fields for Parahydrogen-Induced Polarization of Biomolecules. *arXiv: 2205.15709*, **2022**, DOI: [10.48550/arxiv.2205.15709](https://doi.org/10.48550/arxiv.2205.15709) (accessed June 22, 2022).
- (8) Goldman, M.; Jóhannesson, H. Conversion of a Proton Pair Para Order into ^{13}C Polarization by Rf Irradiation, for Use in MRI. *C. R. Phys.* **2005**, *6*, 575–581.
- (9) Ripka, B.; Eills, J.; Kouřilová, H.; Leutzsch, M.; Levitt, M. H.; Münnemann, K. Hyperpolarized Fumarate via Parahydrogen. *Chem. Commun.* **2018**, *54*, 12246–12249.
- (10) Pravdivtsev, A. N.; Yurkovskaya, A. V.; Lukzen, N. N.; Ivanov, K. L.; Vieth, H.-M. Highly Efficient Polarization of Spin-1/2 Insensitive NMR Nuclei by Adiabatic Passage through Level Anticrossings. *J. Phys. Chem. Lett.* **2014**, *5*, 3421–3426.
- (11) Jóhannesson, H.; Axelsson, O.; Karlsson, M. Transfer of Para-Hydrogen Spin Order into Polarization by Diabatic Field Cycling. *C. R. Phys.* **2004**, *5*, 315–324.
- (12) Joalland, B.; Schmidt, A. B.; Kabir, M. S. H.; Chukanov, N. V.; Kovtunov, K. V.; Koptyug, I. V.; Hennig, J.; Hövener, J.-B.; Chekmenev, E. Y. Pulse-Programmable Magnetic Field Sweeping of Parahydrogen-Induced Polarization by Side Arm Hydrogenation. *Anal. Chem.* **2020**, *92*, 1340–1345.
- (13) Rodin, B. A.; Eills, J.; Picazo-Frutos, R.; Sheberstov, K. F.; Budker, D.; Ivanov, K. L. Constant-Adiabaticity Ultralow Magnetic Field Manipulations of Parahydrogen-Induced Polarization: Application to an AA'X Spin System. *Phys. Chem. Chem. Phys.* **2021**, *23*, 7125–7134.
- (14) Knecht, S.; Blanchard, J. W.; Barskiy, D.; Cavallari, E.; Dagys, L.; van Dyke, E.; Tsukanov, M.; Bliemel, B.; Münnemann, K.; Aime, S.; Reineri, F.; Levitt, M. H.; Buntkowsky, G.; Pines, A.; Blümlein, P.; Budker, D.; Eills, J. Rapid Hyperpolarization and Purification of the Metabolite Fumarate in Aqueous Solution. *Proc. Natl. Acad. Sci. U. S. A.* **2021**, *118*, No. e2025383118.
- (15) Bowers, C. R.; Weitekamp, D. P. Transformation of Symmetrization Order to Nuclear-Spin Magnetization by Chemical Reaction and Nuclear Magnetic Resonance. *Phys. Rev. Lett.* **1986**, *57*, 2645–2648.
- (16) Bowers, C. R.; Weitekamp, D. P. Parahydrogen and Synthesis Allow Dramatically Enhanced Nuclear Alignment. *J. Am. Chem. Soc.* **1987**, *109*, 5541–5542.
- (17) Schmidt, A. B.; Bowers, C. R.; Buckenmaier, K.; Chekmenev, E. Y.; de Maissin, H.; Eills, J.; Ellermann, F.; Glöggler, S.; Gordon, J. W.; Knecht, S.; Koptyug, I. v.; Kuhn, J.; Pravdivtsev, A. N.; Reineri, F.; Theis, T.; Them, K.; Hövener, J. B. Instrumentation for Hydrogenative Parahydrogen-Based Hyperpolarization Techniques. *Anal. Chem.* **2022**, *94*, 479–502.

- (18) Hovener, J. B.; Pravditzsev, A. N.; Kidd, B.; Bowers, C. R.; Gloggler, S.; Kovtunov, K. v.; Plaumann, M.; Katz-Bruell, R.; Buckenmaier, K.; Jerschow, A.; Reineri, F.; Theis, T.; Shchepin, R. v.; Wagner, S.; Bhattacharya, P.; Zacharias, N. M.; Chekmenev, E. Y. Parahydrogen-Based Hyperpolarization for Biomedicine. *Angew. Chem., Int. Ed.* **2018**, *57*, 11140–11162.
- (19) Ardenkjaer-Larsen, J. H. On the Present and Future of Dissolution-DNP. *J. Magn. Reson.* **2016**, *264*, 3–12.
- (20) Vander Heiden, M. G.; Cantley, L. C.; Thompson, C. B. Understanding the Warburg Effect: The Metabolic Requirements of Cell Proliferation. *Science* **2009**, *324*, 1029–1033.
- (21) Kurhanewicz, J.; Vigneron, D. B.; Brindle, K.; Chekmenev, E. Y.; Comment, A.; Cunningham, C. H.; DeBerardinis, R. J.; Green, G. G.; Leach, M. O.; Rajan, S. S.; Rizi, R. R.; Ross, B. D.; Warren, W. S.; Malloy, C. R. Analysis of Cancer Metabolism by Imaging Hyperpolarized Nuclei: Prospects for Translation to Clinical Research. *Neoplasia* **2011**, *13*, 81–97.
- (22) Bernstein, H. J.; Pople, J. A.; Schneider, W. G. The Analysis of Nuclear Magnetic Resonance Spectra: I. System of Two and Three Nuclei. *Can. J. Chem.* **1957**, *35*, 67–83.
- (23) Pravica, M. G.; Weitekamp, D. P. Net NMR Alignment by Adiabatic Transport of Parahydrogen Addition Products to High Magnetic Field. *Chem. Phys. Lett.* **1988**, *145*, 255–258.
- (24) Reineri, F.; Boi, T.; Aime, S. ParaHydrogen Induced Polarization of ¹³C Carboxylate Resonance in Acetate and Pyruvate. *Nat. Commun.* **2015**, *6*, 5858.
- (25) Syvatova, A.; Kozinenko, V. P.; Chukanov, N. V.; Burueva, D. B.; Chekmenev, E. Y.; Chen, Y.-W.; Hwang, D. W.; Kovtunov, K. V.; Koptyug, I. V. PHIP Hyperpolarized [¹-¹³C]Pyruvate and [¹-¹³C]Acetate Esters via PH-INEPT Polarization Transfer Monitored by ¹³C NMR and MRI. *Sci. Rep.* **2021**, *11*, 5646.
- (26) Bengs, C.; Levitt, M. H. SpinDynamica: Symbolic and Numerical Magnetic Resonance in a Mathematica Environment. *Magn. Reson. Chem.* **2018**, *56*, 374–414.
- (27) Stewart, N. J.; Kumeta, H.; Tomohiro, M.; Hashimoto, T.; Hatae, N.; Matsumoto, S. Long-Range Heteronuclear J-Coupling Constants in Esters: Implications for ¹³C Metabolic MRI by Side-Arm Parahydrogen-Induced Polarization. *J. Magn. Reson.* **2018**, *296*, 85–92.
- (28) Korchak, S.; Yang, S.; Mamone, S.; Glöggler, S. Pulsed Magnetic Resonance to Signal-Enhance Metabolites within Seconds by Utilizing Para-Hydrogen. *ChemistryOpen* **2018**, *7*, 344–348.
- (29) Dagys, L.; Jagtap, A. P.; Korchak, S.; Mamone, S.; Saul, P.; Levitt, M. H.; Glöggler, S. Nuclear Hyperpolarization of (¹-¹³C)-Pyruvate in Aqueous Solution by Proton-Relayed Side-Arm Hydrogenation. *Analyst* **2021**, *146*, 1772–1778.
- (30) Ding, Y.; Korchak, S.; Mamone, S.; Jagtap, A. P.; Stevanato, G.; Sternkopf, S.; Moll, D.; Schroeder, H.; Becker, S.; Fischer, A.; Gerhardt, E.; Outeiro, T. F.; Opazo, F.; Griesinger, C.; Glöggler, S. Rapidly Signal-Enhanced Metabolites for Atomic Scale Monitoring of Living Cells with Magnetic Resonance. *Chemistry-Methods* **2022**, No. e202200023.
- (31) Bax, A. D. Structure Determination and Spectral Assignment by Pulsed Polarization Transfer via Long-Range ¹H—¹³C Couplings. *J. Magn. Reson.* **1984**, *57*, 314–318.
- (32) Hogben, H. J.; Krzyszyniak, M.; Charnock, G. T. P.; Hore, P. J.; Kuprov, I. Spinach – A Software Library for Simulation of Spin Dynamics in Large Spin Systems. *J. Magn. Reson.* **2011**, *208*, 179–194.
- (33) Haake, M.; Barkemeyer, J.; Bargon, J. Symmetry Breakdown during Parahydrogen-Labeling of Symmetric Substrates: Proton Spin Polarization and Singlet/Triplet Mixing Due to ¹³C-Isotopes in Natural Abundance. *J. Phys. Chem.* **1995**, *99*, 17539–17543.
- (34) Aime, S.; Gobetto, R.; Canet, D. Longitudinal Nuclear Relaxation in an A 2 Spin System Initially Polarized through Para-Hydrogen. *J. Am. Chem. Soc.* **1998**, *120*, 6770–6773.
- (35) Buljubasic, L.; Franzoni, M. B.; Spiess, H. W.; Münnemann, K. Level Anti-Crossings in ParaHydrogen Induced Polarization Experiments with Cs-Symmetric Molecules. *J. Magn. Reson.* **2012**, *219*, 33–40.
- (36) Levitt, M. H. Symmetry Constraints on Spin Dynamics: Application to Hyperpolarized NMR. *J. Magn. Reson.* **2016**, *262*, 91–99.
- (37) Nielsen, N. C.; Schulte-Herbrüggen, T.; Sørensen, O. W. Bounds on Spin Dynamics Tightened by Permutation Symmetry Application to Coherence Transfer in I2S and I3S Spin Systems. *Mol. Phys.* **1995**, *85*, 1205–1216.
- (38) Ardenkjaer-Larsen, J. H.; Fridlund, B.; Gram, A.; Hansson, G.; Hansson, L.; Lerche, M. H.; Servin, R.; Thaning, M.; Golman, K. Increase in Signal-to-Noise Ratio of > 10,000 Times in Liquid-State NMR. *Proc. Natl. Acad. Sci. U. S. A.* **2003**, *100*, 10158–10163.

□ Recommended by ACS

Distinguishing “Through-Space” from “Through-Bonds” Contribution in Indirect Nuclear Spin–Spin Coupling: General Approaches Applied to Complex J_{pp} and J_{pSe} ...

Olga L. Malkina, Vladimir G. Malkin, et al.

JUNE 10, 2022

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ ▶

Decoupling of Spin Decoherence Paths near Zero Magnetic Field

Sven Bodenstedt, Michael C. D. Taylor, et al.

DECEMBER 28, 2021

THE JOURNAL OF PHYSICAL CHEMISTRY LETTERS

READ ▶

Frémy’s Salt as a Low-Persistence Hyperpolarization Agent: Efficient Dynamic Nuclear Polarization Plus Rapid Radical Scavenging

Mattia Negroni, Dennis Kurzbach, et al.

NOVEMBER 02, 2022

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ ▶

Illuminating Ligand Field Contributions to Molecular Qubit Spin Relaxation via T_1 Anisotropy

Nathanael P. Kazmierczak and Ryan G. Hadt

NOVEMBER 03, 2022

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ ▶

Get More Suggestions >