


# Faraday Discussions

Accepted Manuscript



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

This article can be cited before page numbers have been issued, to do this please use: C. Fleischer, S. T. Holmes, K. Levin, S. L. Veinberg and R. Schurko, *Faraday Discuss.*, 2024, DOI: 10.1039/D4FD00089G.

# Characterization of Ephedrine HCl and Pseudoephedrine HCl Using Quadrupolar NMR Crystallography Guided Crystal Structure Prediction

Carl H. Fleischer III,<sup>1,2</sup> Sean T. Holmes,<sup>1,2</sup> Kirill Levin<sup>3</sup>

Stanislav L. Veinberg,<sup>3</sup> Robert W. Schurko<sup>1,2\*</sup>

<sup>1</sup> Department of Chemistry & Biochemistry, Florida State University, Tallahassee, FL 32306

<sup>2</sup> National High Magnetic Field Laboratory, Tallahassee, FL, 32310

<sup>3</sup> Department of Chemistry & Biochemistry, University of Windsor, Windsor, ON, N9B 3P4

\*Author to whom correspondence should be addressed.

E-mail: [rschurko@fsu.edu](mailto:rschurko@fsu.edu)



## Abstract

Quadrupolar NMR crystallography guided crystal structure prediction (QNMRX-CSP) is a nascent protocol for predicting, solving, and refining crystal structures. QNMRX-CSP employs a combination of solid-state NMR data from quadrupolar nuclides (*i.e.*, nuclear spin  $> 1/2$ ), static lattice energies and electric field gradient (EFG) tensors from dispersion-corrected density functional theory (DFT-D2\*) calculations, and powder X-ray diffraction (PXRD) data; however, it has so far been applied only to organic HCl salts with small and rigid organic components, using  $^{35}\text{Cl}$  EFG tensor data for both structural refinement and validation. Herein, the QNMRX-CSP protocol is extended to ephedrine HCl (Eph) and pseudoephedrine HCl (Pse), which are diastereomeric compounds that feature distinct space groups and organic components that are larger and more flexible. A series of benchmarking calculations are used to generate structural models that can be validated against experimental data, and to explore the impacts of the (i) starting structural models (*i.e.*, geometry-optimized fragments based on either a known crystal structure or an isolated gas-phase molecule) and (ii) selection of unit cell parameters and space groups. Finally, we use QNMRX-CSP to predict the structure of Pse in the dosage form Sudafed using only  $^{35}\text{Cl}$  SSNMR data as experimental input. This proof-of-concept work suggests the possibility of employing QNMRX-CSP protocols to solve the structures of organic HCl salts in dosage forms – something which is often beyond the capabilities of conventional, diffraction-based characterization methods.



## 1. Introduction

Advances in theory and computational power have enabled the rise of crystal structure prediction (CSP) methods for the discovery of new materials and refinement of their solid-state structures,<sup>1–7</sup> including active pharmaceutical ingredients (APIs),<sup>8–11</sup> metal-organic frameworks,<sup>12,13</sup> and organic semi-conductors.<sup>14–16</sup> There are numerous commercial and open-access CSP software packages available,<sup>17–23</sup> many of which are used in combination with experimental data and/or other computational methods. For the CSP of organic compounds, the Cambridge Crystallographic Data Center<sup>24</sup> (CCDC) has run six CSP “blind tests” (a seventh is currently underway), where participants are tasked with predicting the crystal structures of various organic molecules, ranging from those with small organic components to intricate multicomponent cocrystals.<sup>25–30</sup> Limitations of CSP methods include, but are not limited to: (i) the need for advanced algorithms for searching the complex structure spaces,<sup>4,31</sup> especially for systems that have high molecular weights, high flexibility,<sup>32,33</sup> and/or multiple independent molecules,<sup>34</sup> all of which can increase the computational cost; and (ii) the difficulties in the ranking the relative energies of candidate structures.<sup>35–37</sup>

NMR crystallography (NMRX), which utilizes a combination of solid-state NMR (SSNMR), quantum chemical computations, and in some cases, powder X-ray diffraction (PXRD) data, is widely used for the prediction, refinement, and validation of structures,<sup>38–42</sup> and has emerged as a means of improving CSP methods.<sup>1–7</sup> Notably, NMRX-CSP can greatly decrease the computational costs associated with CSP methods, while also providing metrics for validation of structural models (*i.e.*, via comparison of experimentally measured and computationally derived NMR parameters and crystallographic data).<sup>43–48</sup> While PXRD provides invaluable information on long-range order, the space group, and the unit cell parameters,<sup>49,50</sup>



SSNMR adds information on local atomic environments through chemical shifts,<sup>51–53</sup> interatomic proximities through dipolar couplings,<sup>54–56</sup> and confirmation of the number of asymmetric units.<sup>57</sup> These data reduce the search space of computational methods, which refine candidate crystal structures, determine relative energy rankings, and calculate NMR interactions for comparison with experiment.<sup>35,36,58</sup>

To date, the vast majority of NMRX-CSP studies on organic systems rely upon the comparisons of experimental and calculated  $^1\text{H}$  and  $^{13}\text{C}$  chemical shifts,<sup>39–42,51–53</sup> however, given the ubiquity of elements with quadrupolar nuclei in organic solids (*i.e.*, spin  $I > 1/2$ , including  $^{14}\text{N}$ ,  $^{17}\text{O}$ ,  $^{23}\text{Na}$ , and  $^{35}\text{Cl}$ ), it seems their exploitation would be of great benefit. The quadrupolar interaction, which is the interaction between the nuclear quadrupole moment and the electric field gradients (EFGs) at the nuclear origin, can be directly measured from SSNMR spectra of quadrupolar nuclides, in the form of the quadrupolar coupling constant,  $C_Q$ , and asymmetry parameter,  $\eta_Q$  (see **Table 1** for definitions). The EFGs, which are described by symmetric, second-rank tensors, are exquisitely sensitive to their local electronic environments, as well as longer-range electrostatic interactions that often do not have significant influences on chemical shifts. The EFG tensor principal components and its orientation in the molecular/crystal frame depend only upon the ground-state electron density; as such, first principles calculations of EFG tensors are very efficient in comparison to those of chemical shielding.<sup>59,60</sup> Despite these advantages, the use of quadrupolar nuclei in NMRX-CSP studies is limited to date, with a few reports of  $^{14}\text{N}$ - $^1\text{H}$  correlation experiments and/or  $^{13}\text{C}$ - $^{14}\text{N}$  residual dipolar couplings to determine the proximities of these atom pairs.<sup>61–63</sup> We suspect that the two main reasons that quadrupolar guided NMRX-CSP has largely gone underutilized are: (i) the acquisition of high quality SSNMR spectra of quadrupolar nuclides has traditionally been regarded as challenging and



outside of the interest of many practitioners of advanced SSNMR methods; and (ii) there have been issues with calculations of EFG tensors in organic solids that match well with experiment.<sup>59,60,64,65</sup> Fortunately, the former issue has largely been resolved, with a wide range of hardware and pulse sequences available to the end user for the investigation of quadrupoles; however, the latter issue has only recently been addressed by including reparametrized semi-empirical dispersion corrections in geometry optimizations.<sup>66,67</sup>

Our group has recently developed a quadrupolar guided NMR crystallographic crystal structure prediction (QNMRX-CSP) protocol for the prediction, validation, and refinement of structures of organic HCl salts using the comparison of experimental and calculated <sup>35</sup>Cl EFG tensors.<sup>68</sup> In this initial work, QNMRX-CSP was benchmarked using five HCl salts and successfully used in two blind tests, all of which have known crystal structures featuring small unit cells and simple, rigid organic molecules. The ultimate objective of QNMRX-CSP is to predict and refine high-quality crystal structures, starting only with a molecular formula and <sup>35</sup>Cl EFG tensor data (and in some cases, PXRD data).

In this work, we extend QNMRX-CSP benchmarking calculations to organic HCl salts featuring larger unit cells and organic molecules of increased conformational complexity. Specifically, we investigate solid forms of (1*R*,2*S*)-(-)-ephedrine HCl (Eph) and (1*S*,2*S*)-(+)-pseudoephedrine HCl (Pse), which have different space groups and unit cell parameters (*N.B.*; it is emphasized that these are not enantiomorphs),<sup>69,70</sup> as well as <sup>35</sup>Cl SSNMR data that were acquired in our laboratory. Four aspects of QNMRX-CSP benchmarking are considered, featuring different starting points and conditions, with the objective of obtaining structural candidates that agree well with the known crystal structures. The final aspect involves applying QNMRX-CSP to predict the structure of Pse in the dosage form Sudafed, using only <sup>35</sup>Cl



SSNMR data of this solid form. Finally, we discuss the expansion of QNMRX-CSP to incorporate other quadrupolar nuclides (*e.g.*,  $^{14}\text{N}$ ,  $^{17}\text{O}$ ), as well as tandem operation with conventional NMRX-CSP methods featuring  $^{13}\text{C}$  and  $^{15}\text{N}$  chemical shifts; this may enable applications to molecules of greater size and complexity, and possibly the characterization of microcrystalline forms of APIs in dosage forms that undergo structural changes, including hydration, disproportionation, amorphization, and phase changes due to tableting.

## 2. Methods

### 2.1 Chemicals

Ephedrine HCl, pseudoephedrine HCl (**Scheme 1**), and Sudafed® Sinus Congestion 30 mg Nasal Decongestant (Sudafed), which contains 19.5 wt-% Pse, were purchased and ground with a pestle to be used in all subsequent experiments.

### 2.2 Powder X-ray Diffraction

PXRD patterns of Eph and Pse were acquired with a Proto AXRD benchtop X-ray diffractometer operating with Bragg-Brentano geometry and featuring a Cu  $K\alpha$  radiation source and a Proto DECTRIS hybrid pixel detector. The X-ray tube voltage and amperage were 30 kV and 20 mA, respectively. Diffraction patterns were acquired with a detector scanning  $2\theta$  from  $5^\circ$  to  $40^\circ$  with a step size of  $0.015^\circ$  and a dwell time of 4 s. The PXRD pattern of Sudafed was acquired using a Rigaku Miniflex X-ray diffractometer operating with Bragg-Brentano geometry and featuring a Cu  $K\alpha$  radiation source and a D/tex Ultra 250 1D silicon strip detector. The X-ray tube voltage and amperage were 40 kV and 15 mA, respectively. Diffraction patterns were acquired with a detector scanning  $2\theta$  from  $2.5^\circ$  to  $50^\circ$  with a step size of  $0.03^\circ$  and at a rate of  $0.5^\circ \text{ min}^{-1}$ .



## 2.3 Solid-State NMR Spectroscopy

**Overview.** SSNMR spectra were acquired at 9.4 T using a Bruker Avance III HD spectrometer and an Oxford wide bore magnet at the University of Windsor (Windsor, ON), where the relevant Larmor frequencies are  $\nu_0(^1\text{H}) = 400.23$  MHz,  $\nu_0(^{13}\text{C}) = 100.65$  MHz,  $\nu_0(^{15}\text{N}) = 40.54$  MHz, and  $\nu_0(^{35}\text{Cl}) = 39.21$  MHz. Static spectra were acquired with a Revolution 5.0 mm o.d. HX static probe with samples packed into glass tubes sealed with Teflon tape, whereas MAS spectra were acquired with a Varian/Chemagnetics 4.0 mm o.d. HX MAS probe with samples packed into zirconia rotors. Additional SSNMR spectra were acquired at 14.1 T using a Bruker Avance NEO spectrometer and an Oxford wide bore magnet at the National high Magnetic Field Laboratory (Tallahassee, FL), where the relevant Larmor frequencies are  $\nu_0(^1\text{H}) = 600.07$  MHz and  $\nu_0(^{13}\text{C}) = 150.89$  MHz. MAS spectra were acquired with a NHMFL-built 3.2 mm o.d. HXY probe with samples packed into 3.2 o.d. zirconia rotors. High-field spectra were acquired at 21.1 T using a Bruker Avance II console and an Oxford standard bore magnet at the National Ultrahigh-Field NMR Facility for Solids (Ottawa, ON), where the relevant Larmor frequencies are  $\nu_0(^1\text{H}) = 899.53$  MHz and  $\nu_0(^{35}\text{Cl}) = 88.13$  MHz. Static and MAS experiments used a Bruker 4.0 mm HX MAS probe with samples packed into 4.0 mm o.d. zirconia rotors. Additional high-field spectra were acquired at 18.8 T using a Bruker Avance NEO spectrometer and an Oxford wide bore magnet at the National high Magnetic Field Laboratory (Tallahassee, FL), where the relevant Larmor frequencies are  $\nu_0(^1\text{H}) = 799.71$  MHz and  $\nu_0(^{35}\text{Cl}) = 78.35$  MHz. Static and MAS experiments used a NHMFL-built 3.2 mm HXY MAS probe with samples packed into a 3.2 mm o.d. zirconia rotors. A listing of all acquisition parameters is provided in **Tables S1 – S2**. All pulse sequences and recommended calibration parameters and standards are available from the authors at <https://github.com/rschurko>.





**$^1\text{H}\rightarrow^{13}\text{C}$  and  $^1\text{H}\rightarrow^{15}\text{N}$  CP/MAS Experiments.** The  $^1\text{H}\rightarrow^{13}\text{C}$  and  $^1\text{H}\rightarrow^{15}\text{N}$  ramped-amplitude CP/MAS pulse sequence was used to obtain the  $^{13}\text{C}$  and  $^{15}\text{N}$  SSNMR spectra.<sup>71–75</sup>  $^{13}\text{C}$  chemical shifts were referenced to TMS at  $\delta_{\text{iso}}(^{13}\text{C}) = 0.0$  ppm using the frequency peak of  $^{13}\text{C}$  labeled  $\alpha$ -glycine at  $\delta_{\text{iso}}(^{13}\text{C}) = 176.5$  ppm as a secondary reference,<sup>76</sup> whereas  $^{15}\text{N}$  chemical shifts were referenced directly to nitromethane at  $\delta_{\text{iso}}(^{15}\text{N}) = 0.0$  ppm.<sup>77</sup>

**$^{35}\text{Cl}\{^1\text{H}\}$  Experiments.**  $^{35}\text{Cl}\{^1\text{H}\}$  spectra were acquired under static conditions at 9.4 T, and under both static and MAS conditions at 21.1 T or 18.8 T. Static spectra at 9.4 T and 18.8 T were acquired using the Hahn-echo sequence with CT-selective  $\pi/2$  pulses and a decoupling field of  $\nu_2(^1\text{H}) = 30 - 50$  kHz.<sup>78,79</sup> Static spectra at 21.1 T were acquired using the quadrupolar echo pulse sequence with CT-selective  $\pi/2$  pulses and a decoupling field of  $\nu_2(^1\text{H}) = 60$  kHz. MAS spectra at 18.8 T and 21.1 T were acquired using the Bloch decay experiment with CT-selective  $\pi/2$  pulses, a decoupling field of  $\nu_2(^1\text{H}) = 50 - 60$  kHz, and MAS rates of  $\nu_{\text{rot}} = 5$  kHz for Eph and  $\nu_{\text{rot}} = 10$  kHz for Pse and Sudafed.  $^{35}\text{Cl}$  chemical shifts were referenced to 0.1 M NaCl at  $\delta_{\text{iso}}(^{35}\text{Cl}) = 0.0$  ppm using NaCl (s) at  $\delta_{\text{iso}}(^{35}\text{Cl}) = -41.11$  ppm as a secondary reference.<sup>80</sup> Spectra were processed using the Bruker TopSpin 4.3 software package and fit using the ssNake v1.4 package.<sup>81</sup> To ensure proper expression of the Euler angles in the ZY'Z'' convention, the results of iterative simulations in ssNake, which uses the ZX'Z'' convention and different definitions for anisotropic chemical shift and quadrupolar parameters, were verified in WSolids1.<sup>82</sup>

## 2.4 Computational Details

**Overview.** QNMRX-CSP has been designed and benchmarked for the crystal structure prediction of small organic HCl salts.<sup>68</sup> The protocol (outlined in Scheme S1 and detailed in §3.2) combines three complementary methods: (i) PXRD (§2.2), (ii)  $^{35}\text{Cl}$  SSNMR (§2.3), and (iii) calculations (*vide infra*). For calculations, two graphical user interfaces are used: (i)



BIOVIA Materials Studio 2020 R3 is used to interface with Polymorph<sup>83</sup> and CASTEP<sup>84</sup> and (ii) CASTEP Data Manager (developed in our laboratory) is used to automate QNMRX-CSP.<sup>68</sup> CASTEP Data Manager takes advantage of a standalone academic version of CASTEP 2020 and is available at <https://github.com/rschurko>.

**Polymorph.** Polymorph<sup>83</sup> is used to explore the conformational space of Eph and Pse. Polymorph requires the specification of a space group, motion groups, and atomic charges. In this context, motion groups are defined as a geometry optimized organic cation and a Cl<sup>-</sup> anion, with Hirshfeld charges assigned to each atom. Subsequently, these inputs are used in the Polymorph routine which employs a four-step approach to generate a maximum of 10,000 candidate crystal structures per trial, where a trial is defined as one complete iteration of the four-step Polymorph routine — this includes the following steps: (i) packing, (ii) clustering, (iii) force-field geometry optimization, and (iv) clustering. Packing uses a Monte-Carlo simulated annealing algorithm to generate the candidate crystal structures with a maximum and minimum temperature of  $1.5 \times 10^5$  K and 300 K, heating and cooling factors of 0.025 and 0.0005, and a minimum move factor of  $1.0 \times 10^{-10}$ . Clustering removes duplicate structures that are generated during the packing step based on a radial distribution cutoff of 7.0 Å, a tolerance of 0.13, and 140 bins. Dreiding force-field<sup>85</sup> geometry optimizations are used to refine the candidate structure and calculate their static lattice energies. Convergence for the force-field geometry optimization is reached after a maximum change in energy of  $2 \times 10^{-5}$  kcal mol<sup>-1</sup>, in force of  $10^{-3}$  kcal mol<sup>-1</sup> Å<sup>-1</sup>, in stress of  $10^{-3}$  GPa, and in atomic displacement of  $10^{-5}$  Å. Candidate structures are clustered again to remove any duplicate structures following force-field geometry optimization. A final round of clustering is performed following all trials of Polymorph.



**CASTEP.** CASTEP<sup>84</sup> is used to conduct plane-wave DFT-D2\*<sup>67</sup> geometry optimizations and subsequent calculations of NMR interaction tensors. Three types of geometry optimizations are used: (i) truncated, (ii) convergent, and (iii) full volume convergent. All optimizations use the RPBE functional with a plane-wave energy cutoff of 800 eV, the ultrasoft pseudopotentials generated on-the-fly<sup>86</sup>, the zeroth-order regular approximation<sup>87</sup>, and a  $k$ -point spacing of 0.05 Å<sup>-1</sup>. The three approaches differ in the implementation of the low memory BFGS scheme,<sup>88</sup> where for truncated geometry optimizations the maximum BFGS cycles is set to 5, whereas the convergent and full volume convergent geometry optimizations are refined until the convergence thresholds. The convergence thresholds are set to be reached after a maximum change in energy of  $5 \times 10^{-6}$  eV atom<sup>-1</sup>, in displacement of  $5 \times 10^{-4}$  Å, and a maximum force of  $10^{-2}$  eV Å<sup>-1</sup>. Full volume convergent calculations have the added convergence threshold of a maximum stress tolerance of 0.02 GPa and use a fixed basis quality with soft compressibility. A nuclear quadrupole moment of  $Q(^{35}\text{Cl}) = -8.112$  fm<sup>2</sup> is used for the calculation of the <sup>35</sup>Cl EFG tensors.<sup>89</sup>

## 2.5 Metrics for Assessing Candidate Structures.

**Overview.** QNMRX-CSP uses metrics that are designed to retain the best candidate structures. Three metrics rely on the comparison of unit cell parameters, static lattice energies, and <sup>35</sup>Cl EFG tensors.

**Unit Cell Parameters.** Candidate structures are compared to the indexed unit cell parameters of the known crystal structures; structures are retained if they fall within  $\pm 20\%$  of the known values.

**Static Lattice Energies.** Calculated static lattice energies of candidate structural models ( $E_{\text{lat}}$ ) are compared to that with the lowest overall static lattice energy ( $E_{\text{low}}$ ). In M2, all structural models that are in the bottom 13.5% of the  $E_{\text{lat}}$  are retained, according to:



$$(i) E_{\text{low}} \leq E_{\text{lat}} \leq 0.865 * E_{\text{low}}$$

In M3, candidate structural models are retained if they have an  $E_{\text{lat}}$  that is less than or equal to a cutoff static lattice energy,  $E_{\text{cutoff}}$ , such that:

$$(ii) E_{\text{lat}} - E_{\text{low}} \leq E_{\text{cutoff}}$$

Here,  $E_{\text{cutoff}}$  is a relative static lattice energy determined to be 135, 50, or 1 kJ mol<sup>-1</sup> greater than  $E_{\text{low}}$ , which are applied in M3 after Steps 1, 2, and 3. The choice of 13.5% in M2 and selections of  $E_{\text{cutoff}}$  in M3 have been determined by extensive benchmarking calculations.<sup>68</sup>

**<sup>35</sup>Cl EFG Tensors.** The similarity of calculated and experimental <sup>35</sup>Cl EFG tensors is assessed using the EFG distance:<sup>67</sup>

$$\Gamma_{\text{EFG}} = \left( \frac{1}{15} [3\Delta_{11}^2 + 3\Delta_{22}^2 + 3\Delta_{33}^2 + 2\Delta_{11}\Delta_{22} + 2\Delta_{22}\Delta_{33} + 2\Delta_{33}\Delta_{11}] \right)^{1/2}$$

$$\Delta_{kk} = |V_{kk}^{\text{calc}} - V_{kk}^{\text{exp}}|$$

where differences between the two sets of principal components of the EFG tensors ( $V_{kk}$ ,  $k = 1, 2, 3$ ) provide the degree of similarity between two tensors (*i.e.*, a value of  $\Gamma_{\text{EFG}} = 0$  MHz indicates that the two tensors are identical). This metric is based on the chemical shift distance proposed by Alderman and Grant.<sup>90</sup>

## 2.6 Structural Validation.

**Overview.** QNMRX-CSP uses two structural validation terms,  $R$ -factor and root mean square deviation (RMSD) of atomic positions, to compare candidate structures predicted by the protocol to the convergent geometry optimized known crystal structures. Recent CCDC<sup>24</sup> blind tests have set a threshold of RMSDs  $\leq 0.80$  Å, and for the latter, an  $R$ -factor  $\leq 10\%$  as acceptable.<sup>91–94</sup>

**$R$ -factor.** PXRD patterns are simulated using the default settings in the Powder Pattern tool in Mercury 2022.3.0.<sup>95</sup> Agreement between the simulated PXRD patterns of the convergent



geometry optimized crystal structures and candidate structures obtained from QNMRX-CSP is assessed by the  $R$ -factor:

$$R\% = \frac{\sum |F_o - F_c|}{\sum |F_o|} \times 100 \%$$

where  $F_o$  is the calculated signal amplitude of the convergent geometry optimized crystal structures and  $F_c$  is the calculated signal amplitude of the candidate structure.

**RMSD in atomic positions.** Candidate structural models in the validation step of QNMRX-CSP are compared to their respective convergent geometry optimized crystal structures via calculation of RMSDs from relative atomic positions using the CSD-Materials Crystal Packing Similarity tool in Mercury 2022.3.0 with a 15-molecule aggregate and a distance and angle tolerance of 20% and 20°, respectively. Isolated organic fragments generated in QNMRX-CSP are compared with a 1-molecule aggregate and a distance and angle tolerance of 20% and 20°, respectively.

### 3. Research and Discussion

#### 3.1 Characterization with Powder X-ray Diffraction and Solid-State NMR Spectroscopy

**PXRD.** The PXRD patterns of Eph and Pse match simulated patterns based on previously reported SCXRD structures (EPHECL02 and PEPHCL, respectively),<sup>69,70</sup> with no indications of impurity phases (**Fig. S1A, S1B; Table S3**). The reported SCXRD structure for Pse acquired at room temperature. No room temperature SCXRD structures have been reported for Eph with either a low  $R$ -factor (*i.e.*, <5%) or all of the atom positions; instead, a simulated PXRD pattern based on a SCXRD crystal structure of Eph determined at 150 K is used for comparison;<sup>69</sup> which accounts for the slight differences between the experimental and simulated patterns. Since the  $R$ -factors of the EPHECHL02 and PEPHCL structures are a bit high (6.57% and 7.5%,



respectively), these make interesting cases for structural prediction and refinement using our protocol.

**$^1\text{H}\rightarrow^{13}\text{C}$  and  $^1\text{H}\rightarrow^{15}\text{N}$  CP/MAS NMR Spectra.** The  $^1\text{H}\rightarrow^{13}\text{C}\{^1\text{H}\}$  CP/MAS spectra of Eph and Pse (**Fig. 1A, 1B**) are consistent with those from a previous report, in which all peaks were assigned to their respective atomic sites,<sup>96</sup> with the exceptions of C7 and C8 (**Scheme 1, Table S4**). The  $^1\text{H}\rightarrow^{15}\text{N}$  CP/MAS spectra for Eph and Pse each feature a single peak corresponding to the nitrogen atoms in the charged secondary amine groups (**Figure S2**); in this instance, the small difference between the  $^{15}\text{N}$  chemical shifts, along with high uncertainties, limits their use for NMRX.

**$^{35}\text{Cl}\{^1\text{H}\}$  NMR Spectra.** The  $^{35}\text{Cl}\{^1\text{H}\}$  spectra of Eph and Pse (**Fig. 2**) feature central transition (CT,  $+1/2 \leftrightarrow -1/2$ ) powder patterns that are influenced by second-order quadrupolar interactions and chemical shift anisotropy. These data are used to extract values of the quadrupolar and anisotropic chemical shift tensor parameters, as well as the relative orientations of the EFG and chemical shift tensors (as described by Euler angles, see **Table 1**). At a base level, the unique  $^{35}\text{Cl}$  powder patterns of Eph and Pse, which arise from unique sets of NMR interaction tensors, serve as spectral fingerprints for each solid form.

The  $\text{Cl}^-$  ion environments of Eph and Pse each feature three  $\text{H}\cdots\text{Cl}$  hydrogen bonds (*i.e.*,  $r(\text{H}\cdots\text{Cl}) \lesssim 2.6 \text{ \AA}$ )<sup>97</sup> involving one alcohol and two charged secondary amine moieties (**Fig. 3; Table 2**). In Eph, the three hydrogen bonds are classified as *short contacts* (*i.e.*,  $r(\text{H}\cdots\text{Cl}) \lesssim 2.2 \text{ \AA}$ ), which are those hydrogen bonds having the highest impact on the  $^{35}\text{Cl}$  EFG tensors.<sup>98–101</sup> From the large number of  $^{35}\text{Cl}$  EFG tensors that have been measured and calculated for organic HCl salts, it is well known that the presence of one or two short contacts typically results in



sizeable magnitudes of  $C_Q$  (e.g., from 6.0 to 10.5 MHz);<sup>80,98,99,102–117</sup> however, in the case of Eph, the magnitude of  $C_Q$  is small (1.23 MHz).

In order to understand the origins of this small  $C_Q$  value in Eph, the local geometry of the  $\text{Cl}^-$  ion is considered. There are clear guidelines for predicting arrangements of point charges that result in zero EFGs at any point in space;<sup>118</sup> for instance, these have been applied by Bryce and co-workers in computational models of a  $\text{ClH}_4^{3-}$  ion, where the magnitudes of  $C_Q$  and  $\eta_Q$  are observed to change as a function of distortion away from tetrahedral symmetry.<sup>80</sup> The most well-known examples include the placement of  $N$  equal point charges at the  $N$  vertices of polyhedra, like tetrahedra, cubes, and octahedra, which yields zero EFGs at a point  $S_0$  at their centers (i.e., the centroid). Interestingly, no EFGs occur at  $S_0$  if  $N/2$  equal point charges are positioned at  $N/2$  vertices such that none are related by inversion through  $S_0$ . The  $\text{ClH}_3^-$  “fragment” in Eph has a trigonal pyramidal arrangement, with all the  $\text{H}\cdots\text{Cl}\cdots\text{H}$  bond angles,  $\angle(\text{HClH})$ , near  $90^\circ$ ; hence, its geometry resembles this latter configuration, with the hydrogen atoms arranged approximately near the three orthogonal vertices of a fictitious cube. However, since the symmetry is not perfectly cubic, there are non-zero EFGs, accounting in part for the small magnitude of  $C_Q$ .

A similar hydrogen bonding arrangement is observed for Pse; however, one of the  $\text{H}\cdots\text{Cl}^-$  hydrogen bonds in the  $\text{ClH}_3^-$  fragment involving a secondary amine group is significantly longer than the other two short contacts. This likely accounts for the  $C_Q = 2.20$  MHz, which is larger than that of Eph.

### 3.2 QNMRX-CSP

QNMRX-CSP was designed to predict the crystal structures of small organic HCl salts.<sup>68</sup> This method is comprised of three *modules*, each of which features distinct *steps* and *metrics* (a



flowchart depicting this protocol is shown in **Scheme S1**). The functions of the three modules are as follows: Module 1 (M1) selects molecular fragments that correspond to “chemically sensible” arrangements of atoms defined in terms of their individual conformations, followed by the assignment of motion groups and atomic Hirshfeld charges. Module 2 (M2) generates thousands of unique candidate structures, which exhibit distinct unit cells and packing arrangements (see §2.4). Finally, Module 3 (M3) refines the structural models through plane-wave DFT-D2\* geometry optimizations and comparison of calculated and experimental  $^{35}\text{Cl}$  EFG tensors. *Steps* are actions in each module applied to candidate structures (*vide infra*). *Metrics* are parameters used to retain the best candidate structures, which include unit cell parameters ( $a$ ,  $b$ ,  $c$ ,  $\alpha$ ,  $\beta$ , and  $\gamma$ ), static lattice energies ( $E_{\text{lat}}$ , along with  $E_{\text{low}}$  and  $E_{\text{cutoff}}$ ), and  $^{35}\text{Cl}$  EFG distances ( $\Gamma_{\text{EFG}}$ ) (see §2.5 for definitions). Candidate structures are validated via comparison to convergent geometry-optimized structural models based on known crystal structures (Eph: OPT-EPHECL02, and Pse: OPT-PEPHCL) by calculating  $R$ -factors (of PXRD patterns) and atomic position RMSDs; those with values falling below the thresholds described in §2.5 are considered valid structural models.

In the ensuing sections, we describe the use of QNMRX-CSP for benchmarking calculations on Eph and Pse (**Table 3**), including calculations using: (i) fragments obtained from a geometry optimized structural model based on a known crystal structure as a starting point (§3.2.1); (ii) geometry optimized fragments based on isolated, gas-phase molecules as a starting point (§3.2.2); (iii) fragments from (i) above for Pse, but with the space group, unit cell parameters, and  $^{35}\text{Cl}$  quadrupolar parameters for Eph (and *vice versa*, §3.2.3); and (iv) fragments from (ii), but only with  $^{35}\text{Cl}$  quadrupolar parameters (XRD data is absent, §3.2.4). The purposes of (iii) and (iv) are to explore the outcome of QNMRX-CSP calculations when experimental data





is limited or unavailable, with the hope of developing general applications to *de novo* crystal structure predictions.

**3.2.1 Benchmarking starting from a refined crystal structure.** In order to benchmark the QNMRX-CSP protocol and its metrics, the first sets of calculations use the known crystal structures as starting points. The application of the protocol to Eph is considered first (see **Table 4** for an overview of modules, steps, metrics, and numbers of candidate structures).

The starting point is Module 1 (M1): (i) in M1 Step 1, the crystal structure of Eph (EPHECL02) is obtained from the CCDC; (ii) in M1 Step 2, a convergent geometry optimization is conducted to refine the atomic coordinates; (iii) in M1 Step 3, Hirshfeld charges (**Table S5**), as determined from population analysis, are assigned to the atoms; and (iv) in M1 Step 4, the organic cation and chloride anion are assigned as motion groups (**Fig. 4**).

In Module 2 (M2), the motion groups from M1 are used in the four-step Polymorph routine (see **§2.4**), which consists of packing, clustering, force-field geometry optimization, and clustering, leading to thousands of candidate structures. This routine is repeated 10 times (each iteration is defined as a *trial*, meaning that there are 10 trials), after which a final clustering of candidate structures is performed, leading to 46,889 candidates. The best candidate structures are retained using the unit cell parameters and energy metrics (unit cell parameters within  $\pm 20\%$  of the experimental values, and the bottom 13.5% of the  $E_{\text{lat}}$ ; see **§2.5** for explanations), retaining 702 structures.

Module 3 (M3) consists of four steps: (i) in M3 Step 1, truncated geometry optimizations (*i.e.*, those that are halted after five BFGS iterations, see **§2.5**) are conducted on the candidate structures followed by application of energy and EFG distance metrics ( $E_{\text{cutoff}} = 135 \text{ kJ mol}^{-1}$  and  $\Gamma_{\text{EFG}} \leq 0.70 \text{ MHz}$ ), resulting in 25 candidate structures being retained; (ii) in M3 Step 2,



convergent geometry optimizations are conducted on the candidate structures, leading to the retention of 19 structures using energy and EFG distance metrics ( $E_{\text{cutoff}} = 135 \text{ kJ mol}^{-1}$  and  $\Gamma_{\text{EFG}} \leq 0.49 \text{ MHz}$ ); (iii) in M3 Step 3, the unit cell parameters of the candidate structures are adjusted to match those of EPHECL02 and subsequent convergent geometry optimizations are conducted, followed by the application of energy and EFG distance metrics ( $E_{\text{cutoff}} = 1 \text{ kJ mol}^{-1}$  and  $\Gamma_{\text{EFG}} \leq 0.49 \text{ MHz}$ ), which retains 6 candidate structures; and (iv) in M3 Step 4, the final 6 structures are assessed for validation (**Fig. 5A**; **Table 5**). In this case, only 1 structure has RMSDs and  $R$ -factors that are below the CCDC thresholds (see §2.6), though the other 5 candidate structures exhibit similar packing motifs (**Fig. S3**; **Table 5**) but fail structural validation.

QNMRIX-CSP is similarly applied to predict the structure of Pse starting from the PEPHCL structure, resulting in 2 candidate structures (**Fig. 6A**; **Table 5**). Both candidate structures have RMSDs and  $R$ -factors well below the CCDC thresholds.

**3.2.2 Benchmarking starting from an isolated molecule.** These calculations are conducted as follows: (i) in M1 Step 1, a molecular structure of the Eph organic cation is built in Materials Studio and centered in a  $30 \times 30 \times 30 \text{ \AA}^3 P1$  unit cell; (ii) in M1 Step 2, the organic cation is subjected to a convergent geometry optimization; (iii) in M1 Step 3, Hirshfeld charges, as determined from a charge database (**Table S5**), are assigned to the organic cation and the  $\text{Cl}^-$  anion; (iv) in M1 Step 4, the organic cation and  $\text{Cl}^-$  anion are assigned as motion groups (**Fig. 4**). The remainder of the QNMRIX-CSP protocol (M2 and M3) is applied as described above. In this case, 10 trials of the Polymorph routine are necessary. Following M3, 1 candidate structure is passed to validation in M3 Step 4 (**Fig. 5B**; **Table 5**) that has RMSDs and  $R$ -factors that are below the CCDC thresholds.



QNMRIX-CSP is similarly applied to Pse; however, a total of 50 trials of the Polymorph routine are necessary, resulting in 1 structure passing validation in M3 Step 4 (**Fig. 6B**; **Table 5**). Comparison of the isolated organic fragments from each of the predicted candidate structures for Eph and Pse to their respective gas-phase refined organic fragments is achieved with the RMSDs; here, the RMSDs of the Eph organic fragments are higher than those of Pse (0.233 Å and 0.073 Å, respectively).

**3.2.3 Benchmarking starting from a molecular fragment that is mismatched with experimental data.** It is of interest to see if the application of QNMRIX-CSP to one molecular fragment, but using the space group, unit cell parameters, and  $^{35}\text{Cl}$  EFG tensors associated with the other, yields structural models that pass the metrics and CCDC thresholds. In M2, the Polymorph routine was used to pack the motion groups corresponding to the Pse cation (obtained from OPT-PEPHCL) and  $\text{Cl}^-$  ions into unit cells with the space group of Eph (*i.e.*,  $P2_1$ ). Candidate structures are retained using the unit cell parameters and  $^{35}\text{Cl}$  EFG tensors of Eph as metrics (**Table 6**). After 100 trials of the Polymorph routine, and the generation of over half a million candidate structures, 2,993 were passed to M3, from which no viable candidate structures were generated for validation. This result could be interpreted as meaning that there is no form of Pse that crystallizes in the  $P2_1$  space group with  $^{35}\text{Cl}$  EFG tensor parameters matching those of Eph.

QNMRIX-CSP was similarly applied to the Eph cation (OPT-EPHECL02) and  $\text{Cl}^-$  anion in the  $P2_12_12_1$  space group of Pse. After only 30 trials in M2, 6 structures were passed to M3 Step 4 for validation. In comparison to OPT-PEPHCL, all 6 structures have RMSDs and *R*-factors above the CCDC threshold (**Fig. 7**; **Table 7**). The 6 structures each feature three  $\text{H}\cdots\text{Cl}$  hydrogen bonds (one alcohol and two secondary amine moieties, **Fig. 3**, **Table 2**), similar to the



Cl<sup>-</sup> environment of Pse (*vide supra*). This might indicate that Eph has a theoretical polymorph with a structure in the  $P2_12_12_1$  space group with <sup>35</sup>Cl EFG tensors similar to those of Pse, though this is beyond the scope of the current study.

**3.2.4 Benchmarking starting from an isolated molecule but in the absence of XRD data.** The motivation for these calculations is to explore the possibility of CSP in instances where <sup>35</sup>Cl EFG tensors are available, but XRD is not. This would be of great value for the CSP of micro- and nanocrystalline HCl APIs in dosage forms. Eph, Pse, and the dosage form Sudafed, which contains Pse, represent an ideal set of samples for a proof-of-concept study.

Since Eph and Pse are readily differentiated by their distinct <sup>35</sup>Cl CT patterns, it is possible to use <sup>35</sup>Cl SSNMR to determine which, if either, solid form is present in Sudafed. The <sup>35</sup>Cl{<sup>1</sup>H} MAS and static NMR spectra of Sudafed at 18.8 T (**Fig. 8**) have powder patterns matching that of bulk Pse, confirming its presence. In addition, there is a small peak *ca.* -41.1 ppm that indicates the presence of NaCl, which is not detected in the PXRD pattern (**Fig. S4**). The <sup>13</sup>C spectrum of Sudafed (**Fig. 1C**) features peaks corresponding to excipients in the range of *ca.* 60 – 110 ppm; however, the outer peaks match well with those in the spectrum of bulk Pse. The PXRD pattern of Sudafed (**Fig. S1C**) also clearly indicates the presence of Pse, based on comparison to its simulated PXRD pattern. However, indexing the PXRD data to obtain information on the space group and unit cell parameters is nontrivial due to the interfering signals from the excipients.

There are several considerations for using QNMRX-CSP in the absence of XRD data: (i) the choice of space group(s) that should be searched using the Polymorph routine (M2); (ii) determination of valid unit cell parameters for selecting candidate structures (M2); and (iii) the choice of unit cell parameters that should be used in the last stages of structural refinement (M3).



To address the first consideration, the CCDC database was data mined to determine the number of crystal structures with the following criteria: (i) C, H, O, N, and Cl atoms only; (ii) the Cl atom is an anion; and (iii)  $Z' = 1$ . We identified 4,713 structures meeting these criteria, whose space groups were ranked according to their frequency. Five space groups (*i.e.*,  $P2_1/c$ ,  $P\bar{1}$ ,  $P2_12_12_1$ ,  $P2_1$ , and  $Pna2_1$ ) with  $Z \leq 4$  account for 81.4 % of structures (**Table S6**). Herein, the Polymorph routine is applied only to structural models in these five space groups to limit the computational cost. To address the second consideration, the 1,000 candidate structures with the lowest energies resulting from 10 trials of the Polymorph routine are retained, regardless of the predicted unit cell parameters (*i.e.*, structural models are retained based only on relative energies). Finally, the third consideration is addressed by conducting a full volume convergent geometry optimization (§2.4) to refine the unit cell parameters and atomic coordinates of the candidate structures, as opposed to adjusting the former to match those of the indexed PXRD pattern.

With these considerations in mind, QNMRX-CSP was performed to predict the structure of Pse in Sudafed (**Table 8**) using only  $^{35}\text{Cl}$  quadrupolar parameters. Starting with the refined gas-phase motion group of Pse described in §3.2.2, 10 trials of the Polymorph routine were conducted for each of the five space groups. The 1,000 candidate structures with the lowest static lattice energies in each space group were retained and clustered to remove duplicates. The remaining candidate structures were subjected to truncated geometry optimizations, followed by retention of structures using only the EFG distance metric (*i.e.*,  $\Gamma_{\text{EFG}} \leq 0.70$  MHz). This resulted in the elimination of all candidate structures with the space groups  $P\bar{1}$ ,  $P2_1$ , and  $Pna2_1$ . Convergent geometry optimization and application of the second EFG distance metric (*i.e.*,  $\Gamma_{\text{EFG}} \leq 0.49$  MHz) led to the retention of 6 structural models, all in the  $P2_12_12_1$  space group. Final



volume convergent geometry optimizations resulted in 6 candidate structures that were retained for structural validation.

The final structural models have similar packing motifs and unit cell volumes (**Fig. 9**) with RMSDs below the CCDC thresholds (**Table 9**). However, they have unit cell volumes that are overestimated relative to that of the known crystal structure by *ca.* 1.2 – 1.5%, which leads to poor agreement between their simulated PXRD patterns and that of the known crystal structure (**Fig. S5**). This is not unexpected, as it has been demonstrated that full volume convergent DFT-D2\* geometry optimizations tend to overestimate unit cell volumes.<sup>67</sup>

#### 4. Conclusions

Herein, QNMRX-CSP has been extended to systems featuring crystal structures of greater complexity than those investigated to date (*i.e.*, higher molecular weights, greater conformational flexibility in the organic components, and larger unit cells). First, it was demonstrated that <sup>35</sup>Cl SSNMR spectra of Eph and Pse provide distinct spectral fingerprints for each crystalline phase, with the <sup>35</sup>Cl EFG tensors lending insight into the distinct hydrogen bonding arrangements of the Cl<sup>-</sup> ions. Second, QNMRX-CSP benchmarking calculations were conducted for structural models of Eph and Pse. Two sets of calculations with different geometry-optimized organic fragments as starting points (*i.e.*, one from the known crystal structure and the other from the gas phase) were carried out, yielding structural models in good agreement with known structures, featuring RMSDs  $\leq 0.80$  Å and *R*-factor  $\leq 10\%$ , which are below recommended CCDC thresholds.<sup>91–94</sup> Third, QNMRX-CSP benchmarking calculations were conducted using a geometry-optimized motion group of Pse packed into unit cells of the space group of Eph, where the unit cell parameters and <sup>35</sup>Cl EFG tensors of Eph were used to



retain the best candidate structures (and *vice versa*). In both cases, it was found that no viable candidate structures emerged. Finally, using only the  $^{35}\text{Cl}$  EFG tensors determined from the  $^{35}\text{Cl}$  SSNMR spectra of a sample of Sudafed, we confirmed that Pse is the solid form therein, and conducted QNMRX-CSP benchmarking calculations that yield candidate structural models that passed most metrics, with the exception of the *R*-factors – this is due to small overestimations of the unit cell volumes in the DFT-D2\* calculations.

QNMRX-CSP, while still in the early stages of development, shows much promise; however, it still faces challenges, many of which can lead to future extensions of the protocol. As systems grow in complexity, the use of other quadrupolar nuclides (*i.e.*,  $^{14}\text{N}$  and  $^{17}\text{O}$ ) could prove useful for predicting the best structural models – their involvement in both covalent and hydrogen bonds would provide a different perspective on structure. The tandem use of QNMRX-CSP with methods employing either first principles calculations of chemical shielding or rapid prediction of chemical shifts could also be beneficial (*i.e.*,  $^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{15}\text{N}$ ). For instance, determination of the best starting fragments with the aid of DFT calculations of chemical shifts would be relatively inexpensive, since these could serve to limit the number of possible molecular conformations.<sup>43,119,120</sup> Conversely, the use of artificial intelligence methods (*e.g.*, ShiftML<sup>121,122</sup>) could be advantageous for filtering candidate structures. This increased efficiency could lead in several directions. First, it may be possible to predict previously unknown polymorphs, subject to modification of the metrics for choosing candidate structures (*e.g.*, multiple NMR interaction tensors with benchmarked cutoff values). Second, this would permit an exploration of an increased number of space groups (as opposed to the five discussed herein) and a more extensive conformational space. Finally, difficulties related to geometry optimizations remain. We currently use the DFT-D2\* method, which is known to slightly



overestimate unit cell volumes. It is possible that higher-order dispersion corrections (*e.g.*, DFT-D3 or DFT-D3/BJ)<sup>123–125</sup> can be reparametrized for organic solids in a similar manner to DFT-D2 – this would greatly benefit from the increased efficiency in calculations described above. These considerations can open doors for the discovery of new solid forms, including polymorphs, and perhaps even solvates and hydrates – all of which are relevant for *de novo* CSP of APIs *in situ* in complex dosage forms.





## Conflicts of Interest

There are no conflicts to declare.

## Data availability

The pulse sequences used and the CASTEP data manager (CDM), an executable application developed in-house, are available on the web from <https://github.com/rschurko>.

Additional data are available in the ESI.†

## Author Contributions

C.H. Fleischer III, and S.T. Veinberg acquired the PXRD data. C.H. Fleischer III, S.T. Holmes, K. Levin, and S.T. Veinberg participated in the acquisition of the SSNMR data. C.H. Fleischer III conducted QNMRX-CSP. C.H. Fleischer III, S.T. Holmes, and R.W. Schurko contributed to the writing and editing of the manuscript.

## Acknowledgements

We are grateful for research support from The Florida State University and the National High Magnetic Field Laboratory (NHMFL), which is funded by the National Science Foundation Cooperative Agreement (DMR-1644779, DMR-2128556) and by the State of Florida. This work was supported in part by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Award Number DE-SC0022310, covering expenses related to postdoctoral wages. We also thank Genentech and the Natural Sciences and Engineering Research Council of Canada (NSERC, RGPIN-2016\_06642 Discovery Grant) for funding early stages of this research. We thank Dr. Victor Terskikh for acquiring  $^{35}\text{Cl}$  SSNMR spectra at 21.1



T at the National Ultrahigh-Field NMR Facility for Solids (Ottawa, Canada), a national research facility funded by the Canada Foundation for Innovation, the Ontario Innovation Trust, Recherche Quebec, the National Research Council of Canada, and Bruker BioSpin (<http://www.nmr900.ca>).



## References

- (1) Woodley, S. M.; Catlow, R. Crystal Structure Prediction from First Principles. *Nat. Mater.* **2008**, *7*, 937–946. <https://doi.org/10.1038/nmat2321>.
- (2) Day, G. M. Current Approaches to Predicting Molecular Organic Crystal Structures. *Crystallogr. Rev.* **2011**, *17*, 3–52. <https://doi.org/10.1080/0889311X.2010.517526>.
- (3) Price, S. L. Predicting Crystal Structures of Organic Compounds. *Chem. Soc. Rev.* **2014**, *43*, 2098–2111. <https://doi.org/10.1039/c3cs60279f>.
- (4) Oganov, A. R.; Pickard, C. J.; Zhu, Q.; Needs, R. J. Structure Prediction Drives Materials Discovery. *Nat. Rev. Mater.* **2019**, *4*, 331–348. <https://doi.org/10.1038/s41578-019-0101-8>.
- (5) Bowskill, D. H.; Sugden, I. J.; Konstantinopoulos, S.; Adjiman, C. S.; Pantelides, C. C. Crystal Structure Prediction Methods for Organic Molecules: State of the Art. *Annu. Rev. Chem. Biomol. Eng.* **2021**, *12*, 593–623. <https://doi.org/10.1146/annurev-chembioeng-060718-030256>.
- (6) Wang, Y.; Lv, J.; Gao, P.; Ma, Y. Crystal Structure Prediction via Efficient Sampling of the Potential Energy Surface. *Acc. Chem. Res.* **2022**, *55*, 2068–2076. <https://doi.org/10.1021/acs.accounts.2c00243>.
- (7) Beran, G. J. O. Frontiers of Molecular Crystal Structure Prediction for Pharmaceuticals and Functional Organic Materials. *Chem. Sci.* **2023**, *14*, 13290–13312. <https://doi.org/10.1039/d3sc03903j>.
- (8) Baias, M.; Widdifield, C. M.; Dumez, J. N.; Thompson, H. P. G.; Cooper, T. G.; Salager, E.; Bassil, S.; Stein, R. S.; Lesage, A.; Day, G. M.; Emsley, L. Powder Crystallography of Pharmaceutical Materials by Combined Crystal Structure Prediction and Solid-State <sup>1</sup>H NMR Spectroscopy. *Phys. Chem. Chem. Phys.* **2013**, *15*, 8069–8080.



- <https://doi.org/10.1039/c3cp41095a>.
- (9) Neumann, M. A.; Van De Streek, J.; Fabbiani, F. P. A.; Hidber, P.; Grassmann, O. Combined Crystal Structure Prediction and High-Pressure Crystallization in Rational Pharmaceutical Polymorph Screening. *Nat. Commun.* **2015**, *6*. <https://doi.org/10.1038/ncomms8793>.
- (10) Leblanc, L. M.; Johnson, E. R. Crystal-Energy Landscapes of Active Pharmaceutical Ingredients Using Composite Approaches. *CrystEngComm* **2019**, *21*, 5995–6009. <https://doi.org/10.1039/c9ce00895k>.
- (11) Sun, G.; Jin, Y.; Li, S.; Yang, Z.; Shi, B.; Chang, C.; Abramov, Y. A. Virtual Cofomer Screening by Crystal Structure Predictions: Crucial Role of Crystallinity in Pharmaceutical Cocrystallization. *J. Phys. Chem. Lett.* **2020**, *11*, 8832–8838. <https://doi.org/10.1021/acs.jpcclett.0c02371>.
- (12) Darby, J. P.; Arhangel'skis, M.; Katsenis, A. D.; Marrett, J. M.; Frišćić, T.; Morris, A. J. Ab Initio Prediction of Metal-Organic Framework Structures. *Chem. Mater.* **2020**, *32*, 5835–5844. <https://doi.org/10.1021/acs.chemmater.0c01737>.
- (13) Xu, Y.; Marrett, J. M.; Titi, H. M.; Darby, J. P.; Morris, A. J.; Frišćić, T.; Arhangel'skis, M. Experimentally Validated Ab Initio Crystal Structure Prediction of Novel Metal-Organic Framework Materials. *J. Am. Chem. Soc.* **2023**, *145*, 3515–3525. <https://doi.org/10.1021/jacs.2c12095>.
- (14) Sokolov, A. N.; Atahan-Evrenk, S.; Mondal, R.; Akkerman, H. B.; Sánchez-Carrera, R. S.; Granados-Focil, S.; Schrier, J.; Mannsfeld, S. C. B.; Zoombelt, A. P.; Bao, Z.; Aspuru-Guzik, A. From Computational Discovery to Experimental Characterization of a High Hole Mobility Organic Crystal. *Nat. Commun.* **2011**, *2*. <https://doi.org/10.1038/ncomms1451>.



- (15) Cheng, C. Y.; Campbell, J. E.; Day, G. M. Evolutionary Chemical Space Exploration for Functional Materials: Computational Organic Semiconductor Discovery. *Chem. Sci.* **2020**, *11*, 4922–4933. <https://doi.org/10.1039/d0sc00554a>.
- (16) Schmidt, J. A.; Weatherby, J. A.; Sugden, I. J.; Santana-Bonilla, A.; Salerno, F.; Fuchter, M. J.; Johnson, E. R.; Nelson, J.; Jelfs, K. E. Computational Screening of Chiral Organic Semiconductors: Exploring Side-Group Functionalization and Assembly to Optimize Charge Transport. *Cryst. Growth Des.* **2021**, *21*, 5036–5049. <https://doi.org/10.1021/acs.cgd.1c00473>.
- (17) Gale, J. D. GULP: A Computer Program for the Symmetry-Adapted Simulation of Solids. *J. Chem. Soc. - Faraday Trans.* **1997**, *93*, 629–637. <https://doi.org/10.1039/a606455h>.
- (18) Glass, C. W.; Oganov, A. R.; Hansen, N. USPEX—Evolutionary Crystal Structure Prediction. *Comput. Phys. Commun.* **2006**, *175*, 713–720. <https://doi.org/10.1016/j.cpc.2006.07.020>.
- (19) Lonie, D. C.; Zurek, E. XtalOpt: An Open-Source Evolutionary Algorithm for Crystal Structure Prediction. *Comput. Phys. Commun.* **2011**, *182*, 372–387. <https://doi.org/10.1016/j.cpc.2010.07.048>.
- (20) Pickard, C. J.; Needs, R. J. Ab Initio Random Structure Searching. *J. Phys. Condens. Matter* **2011**, *23*. <https://doi.org/10.1088/0953-8984/23/5/053201>.
- (21) Wang, Y.; Lv, J.; Zhu, L.; Ma, Y. CALYPSO: A Method for Crystal Structure Prediction. *Comput. Phys. Commun.* **2012**, *183*, 2063–2070. <https://doi.org/10.1016/j.cpc.2012.05.008>.
- (22) Yamashita, T.; Kanehira, S.; Sato, N.; Kino, H.; Terayama, K.; Sawahata, H.; Sato, T.; Utsuno, F.; Tsuda, K.; Miyake, T.; Oguchi, T. CrySPY: A Crystal Structure Prediction Tool Accelerated by Machine Learning. *Sci. Technol. Adv. Mater. Methods* **2021**, *1*, 87–97.



<https://doi.org/10.1080/27660400.2021.1943171>.

- (23) Wang, J.; Gao, H.; Han, Y.; Ding, C.; Pan, S.; Wang, Y.; Jia, Q.; Wang, H. T.; Xing, D.; Sun, J. MAGUS: Machine Learning and Graph Theory Assisted Universal Structure Searcher. *Natl. Sci. Rev.* **2023**, *10*. <https://doi.org/10.1093/nsr/nwad128>.
- (24) Groom, C. R.; Bruno, I. J.; Lightfoot, M. P.; Ward, S. C. The Cambridge Structural Database. *Acta Crystallogr. Sect. B Struct. Sci. Cryst. Eng. Mater.* **2016**, *72*, 171–179. <https://doi.org/10.1107/S2052520616003954>.
- (25) Lommerse, J. P. M.; Motherwell, W. D. S.; Ammon, H. L.; Dunitz, J. D.; Gavezzotti, A.; Hofmann, D. W. M.; Leusen, F. J. J.; Mooij, W. T. M.; Price, S. L.; Schweizer, B.; Schmidt, M. U.; van Eijck, B. P.; Verwer, P.; Williams, D. E. A Test of Crystal Structure Prediction of Small Organic Molecules. *Acta Crystallogr. Sect. B Struct. Sci.* **2000**, *56*, 697–714. <https://doi.org/10.1107/S0108768100004584>.
- (26) Motherwell, W. D. S.; Ammon, H. L.; Dunitz, J. D.; Dzyabchenko, A.; Erk, P.; Gavezzotti, A.; Hofmann, D. W. M.; Leusen, F. J. J.; Lommerse, J. P. M.; Mooij, W. T. M.; Price, S. L.; Scheraga, H.; Schweizer, B.; Schmidt, M. U.; van Eijck, B. P.; Verwer, P.; Williams, D. E. Crystal Structure Prediction of Small Organic Molecules: A Second Blind Test. *Acta Crystallogr. Sect. B Struct. Sci.* **2002**, *58*, 647–661. <https://doi.org/10.1107/S0108768102005669>.
- (27) Day, G. M.; Motherwell, W. D. S.; Ammon, H. L.; Boerrigter, S. X. M.; Della Valle, R. G.; Venuti, E.; Dzyabchenko, A.; Dunitz, J. D.; Schweizer, B.; Van Eijck, B. P.; Erk, P.; Facelli, J. C.; Bazterra, V. E.; Ferraro, M. B.; Hofmann, D. W. M.; Leusen, F. J. J.; Liang, C.; Pantelides, C. C.; Karamertzanis, P. G.; Price, S. L.; Lewis, T. C.; Nowell, H.; Torrisi, A.; Scheraga, H. A.; Arnautova, Y. A.; Schmidt, M. U.; Verwer, P. A Third Blind Test of



- Crystal Structure Prediction. *Acta Crystallogr. Sect. B Struct. Sci.* **2005**, *61*, 511–527.  
<https://doi.org/10.1107/S0108768105016563>.
- (28) Day, G. M.; Cooper, T. G.; Cruz-Cabeza, A. J.; Hejczyk, K. E.; Ammon, H. L.; Boerrigter, S. X. M.; Tan, J. S.; Della Valle, R. G.; Venuti, E.; Jose, J.; Gadre, S. R.; Desiraju, G. R.; Thakur, T. S.; van Eijck, B. P.; Facelli, J. C.; Bazterra, V. E.; Ferraro, M. B.; Hofmann, D. W. M.; Neumann, M. A.; Leusen, F. J. J.; Kendrick, J.; Price, S. L.; Misquitta, A. J.; Karamertzanis, P. G.; Welch, G. W. A.; Scheraga, H. A.; Arnautova, Y. A.; Schmidt, M. U.; van de Streek, J.; Wolf, A. K.; Schweizer, B. Significant Progress in Predicting the Crystal Structures of Small Organic Molecules – a Report on the Fourth Blind Test. *Acta Crystallogr. Sect. B Struct. Sci.* **2009**, *65*, 107–125.  
<https://doi.org/10.1107/S0108768109004066>.
- (29) Bardwell, D. A.; Adjiman, C. S.; Arnautova, Y. A.; Bartashevich, E.; Boerrigter, S. X. M.; Braun, D. E.; Cruz-Cabeza, A. J.; Day, G. M.; Della Valle, R. G.; Desiraju, G. R.; van Eijck, B. P.; Facelli, J. C.; Ferraro, M. B.; Grillo, D.; Habgood, M.; Hofmann, D. W. M.; Hofmann, F.; Jose, K. V. J.; Karamertzanis, P. G.; Kazantsev, A. V.; Kendrick, J.; Kuleshova, L. N.; Leusen, F. J. J.; Maleev, A. V.; Misquitta, A. J.; Mohamed, S.; Needs, R. J.; Neumann, M. A.; Nikylov, D.; Orendt, A. M.; Pal, R.; Pantelides, C. C.; Pickard, C. J.; Price, L. S.; Price, S. L.; Scheraga, H. A.; van de Streek, J.; Thakur, T. S.; Tiwari, S.; Venuti, E.; Zhitkov, I. K. Towards Crystal Structure Prediction of Complex Organic Compounds – a Report on the Fifth Blind Test. *Acta Crystallogr. Sect. B Struct. Sci.* **2011**, *67*, 535–551.  
<https://doi.org/10.1107/S0108768111042868>.
- (30) Reilly, A. M.; Cooper, R. I.; Adjiman, C. S.; Bhattacharya, S.; Boese, A. D.; Brandenburg, J. G.; Bygrave, P. J.; Bylsma, R.; Campbell, J. E.; Car, R.; Case, D. H.; Chadha, R.; Cole,



J. C.; Cosburn, K.; Cuppen, H. M.; Curtis, F.; Day, G. M.; DiStasio Jr, R. A.; Dzyabchenko, A.; van Eijck, B. P.; Elking, D. M.; van den Ende, J. A.; Facelli, J. C.; Ferraro, M. B.; Fusti-Molnar, L.; Gatsiou, C.-A.; Gee, T. S.; de Gelder, R.; Ghiringhelli, L. M.; Goto, H.; Grimme, S.; Guo, R.; Hofmann, D. W. M.; Hoja, J.; Hylton, R. K.; Iuzzolino, L.; Jankiewicz, W.; de Jong, D. T.; Kendrick, J.; de Klerk, N. J. J.; Ko, H.-Y.; Kuleshova, L. N.; Li, X.; Lohani, S.; Leusen, F. J. J.; Lund, A. M.; Lv, J.; Ma, Y.; Marom, N.; Masunov, A. E.; McCabe, P.; McMahon, D. P.; Meekes, H.; Metz, M. P.; Misquitta, A. J.; Mohamed, S.; Monserrat, B.; Needs, R. J.; Neumann, M. A.; Nyman, J.; Obata, S.; Oberhofer, H.; Oganov, A. R.; Orendt, A. M.; Pagola, G. I.; Pantelides, C. C.; Pickard, C. J.; Podeszwa, R.; Price, L. S.; Price, S. L.; Pulido, A.; Read, M. G.; Reuter, K.; Schneider, E.; Schober, C.; Shields, G. P.; Singh, P.; Sugden, I. J.; Szalewicz, K.; Taylor, C. R.; Tkatchenko, A.; Tuckerman, M. E.; Vacarro, F.; Vasileiadis, M.; Vazquez-Mayagoitia, A.; Vogt, L.; Wang, Y.; Watson, R. E.; de Wijs, G. A.; Yang, J.; Zhu, Q.; Groom, C. R. Report on the Sixth Blind Test of Organic Crystal Structure Prediction Methods. *Acta Crystallogr. Sect. B Struct. Sci. Cryst. Eng. Mater.* **2016**, *72*, 439–459.  
<https://doi.org/10.1107/S2052520616007447>.

- (31) Atahan-Evrenk, Ş.; Aspuru-Guzik, A. *Prediction and Calculation of Crystal Structures: Methods and Applications*; Atahan-Evrenk, S., Aspuru-Guzik, A., Eds.; Topics in Current Chemistry; Springer International Publishing: Cham, 2014; Vol. 345.  
<https://doi.org/10.1007/978-3-319-05774-3>.
- (32) Braun, D. E.; McMahon, J. A.; Bhardwaj, R. M.; Nyman, J.; Neumann, M. A.; Van De Streek, J.; Reutzel-Edens, S. M. Inconvenient Truths about Solid Form Landscapes Revealed in the Polymorphs and Hydrates of Gandotinib. *Cryst. Growth Des.* **2019**, *19*,





- 2947–2962. <https://doi.org/10.1021/acs.cgd.9b00162>.
- (33) Dudek, M. K.; Druzbecki, K. Along the Road to Crystal Structure Prediction (CSP) of Pharmaceutical-like Molecules. *CrystEngComm* **2022**, *24*, 1665–1678. <https://doi.org/10.1039/d1ce01564h>.
- (34) Abramov, Y. A.; Iuzzolino, L.; Jin, Y.; York, G.; Chen, C.-H.; Shultz, C. S.; Yang, Z.; Chang, C.; Shi, B.; Zhou, T.; Greenwell, C.; Sekharan, S.; Lee, A. Y. Cocrystal Synthesis through Crystal Structure Prediction. *Mol. Pharm.* **2023**, *20*, 3380–3392. <https://doi.org/10.1021/acs.molpharmaceut.2c01098>.
- (35) Day, G. M.; Chisholm, J.; Shan, N.; Motherwell, W. D. S.; Jones, W. An Assessment of Lattice Energy Minimization for the Prediction of Molecular Organic Crystal Structures. *Cryst. Growth Des.* **2004**, *4*, 1327–1340. <https://doi.org/10.1021/cg0498148>.
- (36) Nyman, J.; Day, G. M. Static and Lattice Vibrational Energy Differences between Polymorphs. *CrystEngComm* **2015**, *17*, 5154–5165. <https://doi.org/10.1039/c5ce00045a>.
- (37) Taylor, C. R.; Day, G. M. Evaluating the Energetic Driving Force for Cocrystal Formation. *Cryst. Growth Des.* **2018**, *18*, 892–904. <https://doi.org/10.1021/acs.cgd.7b01375>.
- (38) Martineau, C.; Senker, J.; Taulelle, F. NMR Crystallography. In *Annual Reports on NMR Spectroscopy*; Harris, R. K., Wasylishen, R. E., Duer, M. J., Eds.; John Wiley & Sons Ltd.: New York, NY, 2014; Vol. 82, pp 1–57. <https://doi.org/10.1016/B978-0-12-800184-4.00001-1>.
- (39) Ashbrook, S. E.; McKay, D. Combining Solid-State NMR Spectroscopy with First-Principles Calculations – a Guide to NMR Crystallography. *Chem. Commun.* **2016**, *52*, 7186–7204. <https://doi.org/10.1039/C6CC02542K>.
- (40) Bryce, D. L. NMR Crystallography: Structure and Properties of Materials from Solid-State



- Nuclear Magnetic Resonance Observables. *IUCrJ* **2017**, *4*, 350–359.  
<https://doi.org/10.1107/S2052252517006042>.
- (41) Hodgkinson, P. NMR Crystallography of Molecular Organics. *Prog. Nucl. Magn. Reson. Spectrosc.* **2020**, *118–119*, 10–53. <https://doi.org/10.1016/j.pnmrs.2020.03.001>.
- (42) Harris, K. D. M. NMR Crystallography as a Vital Tool in Assisting Crystal Structure Determination from Powder XRD Data. *CRYSTALS* **2022**, *12*.  
<https://doi.org/10.3390/cryst12091277>.
- (43) Harper, J. K.; Grant, D. M. Enhancing Crystal-Structure Prediction with NMR Tensor Data. *Cryst. Growth Des.* **2006**, *6*, 2315–2321. <https://doi.org/10.1021/cg060244g>.
- (44) Salager, E.; Day, G. M.; Stein, R. S.; Pickard, C. J.; Elena, B.; Emsley, L. Powder Crystallography by Combined Crystal Structure Prediction and High-Resolution  $^1\text{H}$  Solid-State NMR Spectroscopy. *J. Am. Chem. Soc.* **2010**, *132*, 2564–2566.  
<https://doi.org/10.1021/ja909449k>.
- (45) Baias, M.; Dumez, J. N.; Svensson, P. H.; Schantz, S.; Day, G. M.; Emsley, L. De Novo Determination of the Crystal Structure of a Large Drug Molecule by Crystal Structure Prediction-Based Powder NMR Crystallography. *J. Am. Chem. Soc.* **2013**, *135*, 17501–17507. <https://doi.org/10.1021/ja4088874>.
- (46) Hofstetter, A.; Balodis, M.; Paruzzo, F. M.; Widdifield, C. M.; Stevanato, G.; Pinon, A. C.; Bygrave, P. J.; Day, G. M.; Emsley, L. Rapid Structure Determination of Molecular Solids Using Chemical Shifts Directed by Unambiguous Prior Constraints. *J. Am. Chem. Soc.* **2019**, *141*, 16624–16634. <https://doi.org/10.1021/jacs.9b03908>.
- (47) Dudek, M. K.; Paluch, P.; Śniechowska, J.; Nartowski, K. P.; Day, G. M.; Potrzebowski, M. J. Crystal Structure Determination of an Elusive Methanol Solvate-Hydrate of Catechin



- Using Crystal Structure Prediction and NMR Crystallography. *CrystEngComm* **2020**, *22*, 4969–4981. <https://doi.org/10.1039/d0ce00452a>.
- (48) Pawlak, T.; Sudgen, I.; Bujacz, G.; Iuga, D.; Brown, S. P.; Potrzebowski, M. J. Synergy of Solid-State NMR, Single-Crystal X-Ray Diffraction, and Crystal Structure Prediction Methods: A Case Study of Teriflunomide (TFM). *Cryst. Growth Des.* **2021**, *21*, 3328–3343. <https://doi.org/10.1021/acs.cgd.1c00123>.
- (49) Woolfson, M. M. *An Introduction to X-Ray Crystallography*, 2nd ed.; Cambridge University Press, 1997.
- (50) Shankland, K. An Overview of Powder X-Ray Diffraction and Its Relevance to Pharmaceutical Crystal Structures; 2016; pp 293–314. [https://doi.org/10.1007/978-1-4939-4029-5\\_8](https://doi.org/10.1007/978-1-4939-4029-5_8).
- (51) Harris, R. K.; Cadars, S.; Emsley, L.; Yates, J. R.; Pickard, C. J.; Jetti, R. K. R.; Griesser, U. J. NMR Crystallography of Oxybuprocaine Hydrochloride, Modification II°. *Phys. Chem. Chem. Phys.* **2007**, *9*, 360–368. <https://doi.org/10.1039/b614318k>.
- (52) Harper, J. K.; Iulicci, R.; Gruber, M.; Kalakewich, K. Refining Crystal Structures with Experimental <sup>13</sup>C NMR Shift Tensors and Lattice-Including Electronic Structure Methods. *CrystEngComm* **2013**, *15*, 8693–8704. <https://doi.org/10.1039/c3ce40108a>.
- (53) Widdifield, C. M.; Farrell, J. D.; Cole, J. C.; Howard, J. A. K.; Hodgkinson, P. Resolving Alternative Organic Crystal Structures Using Density Functional Theory and NMR Chemical Shifts. *Chem. Sci.* **2020**, *11*, 2987–2992. <https://doi.org/10.1039/c9sc04964a>.
- (54) Chierotti, M. R.; Gobetto, R. NMR Crystallography: The Use of Dipolar Interactions in Polymorph and Co-Crystal Investigation. *CrystEngComm* **2013**, *15*, 8599–8612. <https://doi.org/10.1039/c3ce41026a>.



- (55) Thureau, P.; Sturniolo, S.; Zilka, M.; Ziarelli, F.; Viel, S.; Yates, J. R.; Mollica, G. Reducing the Computational Cost of NMR Crystallography of Organic Powders at Natural Isotopic Abundance with the Help of  $^{13}\text{C}$ - $^{13}\text{C}$  Dipolar Couplings. *Magn. Reson. Chem.* **2019**, *57*, 256–264. <https://doi.org/10.1002/mrc.4848>.
- (56) Struppe, J.; Quinn, C. M.; Sarkar, S.; Gronenborn, A. M.; Polenova, T. Ultrafast  $^1\text{H}$  MAS NMR Crystallography for Natural Abundance Pharmaceutical Compounds. *Mol. Pharm.* **2020**, *17*, 674–682. <https://doi.org/10.1021/acs.molpharmaceut.9b01157>.
- (57) Webber, A. L.; Emsley, L.; Claramunt, R. M.; Brown, S. P. NMR Crystallography of Campho[2,3-c]Pyrazole ( $Z' = 6$ ): Combining High-Resolution  $^1\text{H}$ - $^{13}\text{C}$  Solid-State MAS NMR Spectroscopy and GIPAW Chemical-Shift Calculations. *J. Phys. Chem. A* **2010**, *114*, 10435–10442. <https://doi.org/10.1021/jp104901j>.
- (58) Price, S. L. Computed Crystal Energy Landscapes for Understanding and Predicting Organic Crystal Structures and Polymorphism. *Acc. Chem. Res.* **2009**, *42*, 117–126. <https://doi.org/10.1021/ar800147t>.
- (59) Vega, A. J. Quadrupolar Nuclei in Solids. In *Encyclopedia of Magnetic Resonance*; Grant, D. M., Harris, R. K., Eds.; John Wiley & Sons, Ltd: Chichester, UK, 2007; p 3869–3888. <https://doi.org/10.1002/9780470034590.emrstm0431>.
- (60) Autschbach, J.; Zheng, S.; Schurko, R. W. Analysis of Electric Field Gradient Tensors at Quadrupolar Nuclei in Common Structural Motifs. *Concepts Magn. Reson. Part A* **2010**, *36A*, 84–126. <https://doi.org/10.1002/cmr.a.20155>.
- (61) Bravetti, F.; Russo, R. E.; Bordignon, S.; Gallo, A.; Rossi, F.; Nervi, C.; Gobetto, R.; Chierotti, M. R. Zwitterionic or Not? Fast and Reliable Structure Determination by Combining Crystal Structure Prediction and Solid-State NMR. *Molecules* **2023**, *28*, 1876.



- <https://doi.org/10.3390/molecules28041876>.
- (62) Widdifield, C. M.; Nilsson Lill, S. O.; Broo, A.; Lindkvist, M.; Pettersen, A.; Svensk Ankarberg, A.; Aldred, P.; Schantz, S.; Emsley, L. Does  $Z'$  Equal 1 or 2? Enhanced Powder NMR Crystallography Verification of a Disordered Room Temperature Crystal Structure of a P38 Inhibitor for Chronic Obstructive Pulmonary Disease. *Phys. Chem. Chem. Phys.* **2017**, *19*, 16650–16661. <https://doi.org/10.1039/c7cp02349a>.
- (63) Khalaji, M.; Paluch, P.; Potrzebowski, M. J.; Dudek, M. K. Narrowing down the Conformational Space with Solid-State NMR in Crystal Structure Prediction of Linezolid Cocrystals. *Solid State Nucl. Magn. Reson.* **2022**, *121*, 101813. <https://doi.org/10.1016/j.ssnmr.2022.101813>.
- (64) Chapman, R. P.; Bryce, D. L. Application of Multinuclear Magnetic Resonance and Gauge-Including Projector-Augmented-Wave Calculations to the Study of Solid Group 13 Chlorides. *Phys. Chem. Chem. Phys.* **2009**, *11*, 6987. <https://doi.org/10.1039/b906627f>.
- (65) Charpentier, T. The PAW/GIPAW Approach for Computing NMR Parameters: A New Dimension Added to NMR Study of Solids. *Solid State Nucl. Magn. Reson.* **2011**, *40*, 1–20. <https://doi.org/https://doi.org/10.1016/j.ssnmr.2011.04.006>.
- (66) Holmes, S. T.; Schurko, R. W. Refining Crystal Structures with Quadrupolar NMR and Dispersion-Corrected Density Functional Theory. *J. Phys. Chem. C* **2018**, *122*, 1809–1820. <https://doi.org/10.1021/acs.jpcc.7b12314>.
- (67) Holmes, S. T.; Vojvodin, C. S.; Schurko, R. W. Dispersion-Corrected DFT Methods for Applications in Nuclear Magnetic Resonance Crystallography. *J. Phys. Chem. A* **2020**, *124*, 10312–10323. <https://doi.org/10.1021/acs.jpca.0c06372>.
- (68) Peach, A. A.; Fleischer, C. H. I.; Levin, K.; Holmes, S. T.; Sanchez, J. E.; Schurko, R. W.



Quadrupolar NMR Crystallography Guided Crystal Structure Prediction (QNMRX-CSP).  
*CrystEngComm* In review.

- (69) Collier, E. A.; Davey, R. J.; Black, S. N.; Roberts, R. J. 17 Salts of Ephedrine: Crystal Structures and Packing Analysis. *Acta Crystallogr. Sect. B Struct. Sci.* **2006**, *62*, 498–505. <https://doi.org/10.1107/S0108768106012018>.
- (70) Mathew, M.; Palenik, G. J. The Crystal and Molecular Structures of (+)-Pseudoephedrine and (+)-Pseudoephedrine Hydrochloride. *Acta Crystallogr. Sect. B Struct. Crystallogr. Cryst. Chem.* **1977**, *33*, 1016–1022. <https://doi.org/10.1107/S0567740877005287>.
- (71) Pines, A.; Gibby, M. G.; Waugh, J. S. Proton-Enhanced NMR of Dilute Spins in Solids. *J. Chem. Phys.* **1973**, *59*, 569–590. <https://doi.org/10.1063/1.1680061>.
- (72) Peersen, O. .; Wu, X. L.; Kustanovich, I.; Smith, S. O. Variable-Amplitude Cross-Polarization MAS NMR. *J. Magn. Reson. Ser. A* **1993**, *104*, 334–339. <https://doi.org/https://doi.org/10.1006/jmra.1993.1231>.
- (73) Metz, G.; Wu, X. L.; Smith, S. O. Ramped-Amplitude Cross Polarization in Magic Angle Spinning NMR. *J. Magn. Reson. Ser. A1* **1994**, *110*, 219–227. <https://doi.org/https://doi.org/10.1006/jmra.1994.1208>.
- (74) Schaefer, J.; Stejskal, E. O. Carbon-13 Nuclear Magnetic Resonance of Polymers Spinning at the Magic Angle. *J. Am. Chem. Soc.* **1976**, *98*, 1031–1032. <https://doi.org/10.1021/ja00420a036>.
- (75) Pines, A.; Gibby, M. G.; Waugh, J. S. Proton-Enhanced Nuclear Induction Spectroscopy 13C Chemical Shielding Anisotropy in Some Organic Solids. *Chem. Phys. Lett.* **1972**, *15*, 373–376. [https://doi.org/10.1016/0009-2614\(72\)80191-X](https://doi.org/10.1016/0009-2614(72)80191-X).
- (76) Taylor, R. E. 13C CP/MAS: Application to Glycine. *Concepts Magn. Reson.* **2004**, *22A*,



- 79–89. <https://doi.org/10.1002/cmr.a.20015>.
- (77) Bertani, P.; Raya, J.; Bechinger, B. <sup>15</sup>N Chemical Shift Referencing in Solid State NMR. *Solid State Nucl. Magn. Reson.* **2014**, *61–62*, 15–18. <https://doi.org/10.1016/j.ssnmr.2014.03.003>.
- (78) Hahn, E. L. Spin Echoes. *Phys. Rev.* **1950**, *80*, 580–594. <https://doi.org/10.1103/PhysRev.80.580>.
- (79) Chan, J. C. C. Spin Echoes in Half-Integer Quadrupole Systems. *Concepts Magn. Reson.* **1999**, *11*, 363–377. [https://doi.org/10.1002/\(SICI\)1099-0534\(1999\)11:6<363::AID-CMR2>3.0.CO;2-A](https://doi.org/10.1002/(SICI)1099-0534(1999)11:6<363::AID-CMR2>3.0.CO;2-A).
- (80) Bryce, D. L.; Sward, G. D. Chlorine-35/37 NMR Spectroscopy of Solid Amino Acid Hydrochlorides: Refinement of Hydrogen-Bonded Proton Positions Using Experiment and Theory. *J. Phys. Chem. B* **2006**, *110*, 26461–26470. <https://doi.org/10.1021/jp065878c>.
- (81) van Meerten, S. G. J.; Franssen, W. M. J.; Kentgens, A. P. M. SsNake: A Cross-Platform Open-Source NMR Data Processing and Fitting Application. *J. Magn. Reson.* **2019**, *301*, 56–66. <https://doi.org/10.1016/j.jmr.2019.02.006>.
- (82) Eichele, K. WSOLIDS1: Solid-State NMR Simulated (Version 1.21.7). Universität Tübingen, 2021.
- (83) Akkermans, R. L. C.; Spenley, N. A.; Robertson, S. H. Monte Carlo Methods in Materials Studio. *Mol. Simul.* **2013**, *39*, 1153–1164. <https://doi.org/10.1080/08927022.2013.843775>.
- (84) Clark, S. J.; Segall, M. D.; Pickard, C. J.; Hasnip, P. J.; Probert, M. I. J.; Refson, K.; Payne, M. C. First Principles Methods Using CASTEP. *Zeitschrift für Krist. - Cryst. Mater.* **2005**, *220*, 567–570. <https://doi.org/10.1524/zkri.220.5.567.65075>.
- (85) Mayo, S. L.; Olafson, B. D.; Goddard, W. A. DREIDING: A Generic Force Field for





- Molecular Simulations. *J. Phys. Chem.* **1990**, *94*, 8897–8909.  
<https://doi.org/10.1021/j100389a010>.
- (86) Yates, J. R.; Pickard, C. J.; Mauri, F. Calculation of NMR Chemical Shifts for Extended Systems Using Ultrasoft Pseudopotentials. *Phys. Rev. B* **2007**, *76*, 024401.  
<https://doi.org/10.1103/PhysRevB.76.024401>.
- (87) van Lenthe, E.; Snijders, J. G.; Baerends, E. J. The Zero-Order Regular Approximation for Relativistic Effects: The Effect of Spin–Orbit Coupling in Closed Shell Molecules. *J. Chem. Phys.* **1996**, *105*, 6505–6516. <https://doi.org/10.1063/1.472460>.
- (88) Pfrommer, B. G.; Côté, M.; Louie, S. G.; Cohen, M. L. Relaxation of Crystals with the Quasi-Newton Method. *J. Comput. Phys.* **1997**, *131*, 233–240.  
<https://doi.org/10.1006/jcph.1996.5612>.
- (89) Pyykkö, P. Year-2017 Nuclear Quadrupole Moments. *Mol. Phys.* **2018**, *116*, 1328–1338.  
<https://doi.org/10.1080/00268976.2018.1426131>.
- (90) Alderman, D. W.; Sherwood, M. H.; Grant, D. M. Comparing, Modeling, and Assigning Chemical-Shift Tensors in the Cartesian, Irreducible Spherical, and Icosahedral Representations. *J. Magn. Reson. Ser. A* **1993**, *101*, 188–197.  
<https://doi.org/10.1006/jmra.1993.1029>.
- (91) McREE, D. E. COMPUTATIONAL TECHNIQUES. In *Practical Protein Crystallography*; Elsevier, 1999; pp 91–cp1. <https://doi.org/10.1016/B978-012486052-0/50005-1>.
- (92) Rondeau, J.-M.; Schreuder, H. Protein Crystallography and Drug Discovery. In *The Practice of Medicinal Chemistry*; Elsevier, 2015; pp 511–537.  
<https://doi.org/10.1016/B978-0-12-417205-0.00022-5>.
- (93) Allen, F. H. The Cambridge Structural Database: A Quarter of a Million Crystal Structures





- and Rising. *Acta Crystallogr. Sect. B Struct. Sci.* **2002**, *58*, 380–388.  
<https://doi.org/10.1107/S0108768102003890>.
- (94) Barr, G.; Dong, W.; Gilmore, C. J.; Kern, A.; Parkin, A.; Wilson, C. C. Using the Cambridge Structural Database to Validate Powder Structures. *Zeitschrift für Krist.* **2007**, *2007*, 209–214. [https://doi.org/10.1524/zkri.2007.2007.suppl\\_26.209](https://doi.org/10.1524/zkri.2007.2007.suppl_26.209).
- (95) Macrae, C. F.; Sovago, I.; Cottrell, S. J.; Galek, P. T. A.; McCabe, P.; Pidcock, E.; Platings, M.; Shields, G. P.; Stevens, J. S.; Towler, M.; Wood, P. A. Mercury 4.0 : From Visualization to Analysis, Design and Prediction. *J. Appl. Crystallogr.* **2020**, *53*, 226–235.  
<https://doi.org/10.1107/S1600576719014092>.
- (96) Lee, G. S. H.; Taylor, R. C.; Dawson, M.; Kannangara, G. S. K.; Wilson, M. A. High-Resolution Solid State <sup>13</sup>C Nuclear Magnetic Resonance Spectra of 3,4-Methylenedioxyamphetamine Hydrochloride and Related Compounds and Their Mixtures with Lactose. *Solid State Nucl. Magn. Reson.* **2000**, *16*, 225–237.  
[https://doi.org/10.1016/S0926-2040\(00\)00071-0](https://doi.org/10.1016/S0926-2040(00)00071-0).
- (97) Desiraju, G.; Steiner, T. *The Weak Hydrogen Bond*; Oxford University Press: Oxford, England, 2001; Vol. 9. <https://doi.org/10.1093/acprof:oso/9780198509707.001.0001>.
- (98) Hildebrand, M.; Hamaed, H.; Namespetra, A. M.; Donohue, J. M.; Fu, R.; Hung, I.; Gan, Z.; Schurko, R. W. <sup>35</sup>Cl Solid-State NMR of HCl Salts of Active Pharmaceutical Ingredients: Structural Prediction, Spectral Fingerprinting and Polymorph Recognition. *CrystEngComm* **2014**, *16*, 7334–7356. <https://doi.org/10.1039/c4ce00544a>.
- (99) Hamaed, H.; Pawlowski, J. M.; Cooper, B. F. T.; Fu, R.; Eichhorn, S. H.; Schurko, R. W. Application of Solid-State <sup>35</sup>Cl NMR to the Structural Characterization of Hydrochloride Pharmaceuticals and Their Polymorphs. *J. Am. Chem. Soc.* **2008**, *130*, 11056–11065.



<https://doi.org/10.1021/ja802486q>.

- (100) Holmes, S. T.; Vojvodin, C. S.; Veinberg, N.; Iacobelli, E. M.; Hirsh, D. A.; Schurko, R. W. Hydrates of Active Pharmaceutical Ingredients: A  $^{35}\text{Cl}$  and  $^2\text{H}$  Solid-State NMR and DFT Study. *Solid State Nucl. Magn. Reson.* **2022**, *122*, 101837. <https://doi.org/10.1016/j.ssnmr.2022.101837>.
- (101) Vojvodin, C. S.; Holmes, S. T.; Watanabe, L. K.; Rawson, J. M.; Schurko, R. W. Multi-Component Crystals Containing Urea: Mechanochemical Synthesis and Characterization by  $^{35}\text{Cl}$  Solid-State NMR Spectroscopy and DFT Calculations. *CrystEngComm* **2022**, *24*, 2626–2641. <https://doi.org/10.1039/D1CE01610E>.
- (102) Namespetra, A. M.; Hirsh, D. A.; Hildebrand, M. P.; Sandre, A. R.; Hamaed, H.; Rawson, J. M.; Schurko, R. W.  $^{35}\text{Cl}$  Solid-State NMR Spectroscopy of HCl Pharmaceuticals and Their Polymorphs in Bulk and Dosage Forms. *CrystEngComm* **2016**, *18*, 6213–6232. <https://doi.org/10.1039/C6CE01069E>.
- (103) Pandey, M. K.; Kato, H.; Ishii, Y.; Nishiyama, Y. Two-Dimensional Proton-Detected  $^{35}\text{Cl}/^1\text{H}$  Correlation Solid-State NMR Experiment under Fast Magic Angle Sample Spinning: Application to Pharmaceutical Compounds. *Phys. Chem. Chem. Phys.* **2016**, *18*, 6209–6216. <https://doi.org/10.1039/c5cp06042g>.
- (104) Peach, A. A.; Hirsh, D. A.; Holmes, S. T.; Schurko, R. W. Mechanochemical Syntheses and  $^{35}\text{Cl}$  Solid-State NMR Characterization of Fluoxetine HCl Cocrystals. *CrystEngComm* **2018**, *20*, 2780–2792. <https://doi.org/10.1039/c8ce00378e>.
- (105) Hirsh, D. A.; Holmes, S. T.; Chakravarty, P.; Peach, A. A.; Dipasquale, A. G.; Nagapudi, K.; Schurko, R. W. In Situ Characterization of Waters of Hydration in a Variable-Hydrate Active Pharmaceutical Ingredient Using  $^{35}\text{Cl}$  Solid-State NMR and X-Ray Diffraction.



- Cryst. Growth Des.* **2019**, *19*, 7349–7362. <https://doi.org/10.1021/acs.cgd.9b01218>.
- (106) Wijesekara, A. V.; Venkatesh, A.; Lampkin, B. J.; VanVeller, B.; Lubach, J. W.; Nagapudi, K.; Hung, I.; Gor'kov, P. L.; Gan, Z.; Rossini, A. J. Fast Acquisition of Proton-Detected HETCOR Solid-State NMR Spectra of Quadrupolar Nuclei and Rapid Measurement of NH Bond Lengths by Frequency Selective HMQC and RESPDOR Pulse Sequences. *Chem. - A Eur. J.* **2020**, *26*, 7881–7888. <https://doi.org/10.1002/chem.202000390>.
- (107) Iuga, D.; Corlett, E. K.; Brown, S. P. <sup>35</sup>Cl–<sup>1</sup>H Heteronuclear Correlation Magic-Angle Spinning Nuclear Magnetic Resonance Experiments for Probing Pharmaceutical Salts. *Magn. Reson. Chem.* **2021**, *59*, 1089–1100. <https://doi.org/10.1002/mrc.5188>.
- (108) Holmes, S. T.; Hook, J. M.; Schurko, R. W. Nutraceuticals in Bulk and Dosage Forms: Analysis by <sup>35</sup>Cl and <sup>14</sup>N Solid-State NMR and DFT Calculations. *Mol. Pharm.* **2022**, *19*, 440–455. <https://doi.org/10.1021/acs.molpharmaceut.1c00708>.
- (109) Abdulla, L. M.; Peach, A. A.; Holmes, S. T.; Dowdell, Z. T.; Watanabe, L. K.; Iacobelli, E. M.; Hirsh, D. A.; Rawson, J. M.; Schurko, R. W. Synthesis and Characterization of Xylazine Hydrochloride Polymorphs, Hydrates, and Cocrystals: A <sup>35</sup>Cl Solid-State NMR and DFT Study. *Cryst. Growth Des.* **2023**, *23*, 3412–3426. <https://doi.org/10.1021/acs.cgd.2c01539>.
- (110) Szell, P. M. J.; Rehman, Z.; Tatman, B. P.; Hughes, L. P.; Blade, H.; Brown, S. P. Exploring the Potential of Multinuclear Solid-State <sup>1</sup>H, <sup>13</sup>C, and <sup>35</sup>Cl Magnetic Resonance To Characterize Static and Dynamic Disorder in Pharmaceutical Hydrochlorides. *ChemPhysChem* **2023**, *24*. <https://doi.org/10.1002/cphc.202200558>.
- (111) Bryce, D. L.; Gee, M.; Wasylishen, R. E. High-Field Chlorine NMR Spectroscopy of Solid Organic Hydrochloride Salts: A Sensitive Probe of Hydrogen Bonding Environment. *J. Phys. Chem. A* **2001**, *105*, 10413–10421. <https://doi.org/10.1021/jp011962a>.



- (112) Azäs, T.; Bonhomme, C.; Smith, M. E.  $^{35}\text{Cl}$  Quadrupolar Constants Obtained by Solid-State NMR: Study of Chlorinated Al-O-P Clusters, Involving OH  $\cdots$  Cl Hydrogen Bonds. *Solid State Nucl. Magn. Reson.* **2003**, *23*, 14–27. [https://doi.org/10.1016/S0926-2040\(02\)00012-7](https://doi.org/10.1016/S0926-2040(02)00012-7).
- (113) Gervais, C.; Dupree, R.; Pike, K. J.; Bonhomme, C.; Profeta, M.; Pickard, C. J.; Mauri, F. Combined First-Principles Computational and Experimental Multinuclear Solid-State NMR Investigation of Amino Acids. *J. Phys. Chem. A* **2005**, *109*, 6960–6969. <https://doi.org/10.1021/jp0513925>.
- (114) Chapman, R. P.; Bryce, D. L. A High-Field Solid-State  $^{35}/^{37}\text{Cl}$  NMR and Quantum Chemical Investigation of the Chlorine Quadrupolar and Chemical Shift Tensors in Amino Acid Hydrochlorides. *Phys. Chem. Chem. Phys.* **2007**, *9*, 6219–6230. <https://doi.org/10.1039/b712688c>.
- (115) Penner, G. H.; Webber, R.; O'Dell, L. A. A Multinuclear NMR and Quantum Chemical Study of Solid Trimethylammonium Chloride. *Can. J. Chem.* **2011**, *89*, 1036–1046. <https://doi.org/10.1139/v11-034>.
- (116) Vogt, F. G.; Williams, G. R.; Copley, R. C. B. Solid-State NMR Analysis of a Boron-Containing Pharmaceutical Hydrochloride Salt. *J. Pharm. Sci.* **2013**, *102*, 3705–3716. <https://doi.org/10.1002/jps.23679>.
- (117) Vogt, F. G.; Williams, G. R.; Strohmeier, M.; Johnson, M. N.; Copley, R. C. B. Solid-State NMR Analysis of a Complex Crystalline Phase of Ronacaleret Hydrochloride. *J. Phys. Chem. B* **2014**, *118*, 10266–10284. <https://doi.org/10.1021/jp505061j>.
- (118) Akitt, J. W.; McDonald, W. S. Arrangements of Ligands Giving Low Electric Field Gradients. *J. Magn. Reson.* **1984**, *58*, 401–412. <https://doi.org/10.1016/0022->



2364(84)90144-6.

- (119) Harper, J. K.; Mulgrew, A. E.; Li, J. Y.; Barich, D. H.; Strobel, G. A.; Grant, D. M. Characterization of Stereochemistry and Molecular Conformation Using Solid-State NMR Tensors. *J. Am. Chem. Soc.* **2001**, *123*, 9837–9842. <https://doi.org/10.1021/ja0109971>.
- (120) Harper, J. K.; Barich, D. H.; Hu, J. Z.; Strobel, G. A.; Grant, D. M. Stereochemical Analysis by Solid-State NMR: Structural Predictions in Ambuic Acid. *J. Org. Chem.* **2003**, *68*, 4609–4614. <https://doi.org/10.1021/jo020377i>.
- (121) Paruzzo, F. M.; Hofstetter, A.; Musil, F.; De, S.; Ceriotti, M.; Emsley, L. Chemical Shifts in Molecular Solids by Machine Learning. *Nat. Commun.* **2018**, *9*, 4501. <https://doi.org/10.1038/s41467-018-06972-x>.
- (122) Cordova, M.; Engel, E. A.; Stefaniuk, A.; Paruzzo, F.; Hofstetter, A.; Ceriotti, M.; Emsley, L. A Machine Learning Model of Chemical Shifts for Chemically and Structurally Diverse Molecular Solids. *J. Phys. Chem. C* **2022**, *126*, 16710–16720. <https://doi.org/10.1021/acs.jpcc.2c03854>.
- (123) Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. A Consistent and Accurate Ab Initio Parametrization of Density Functional Dispersion Correction (DFT-D) for the 94 Elements H-Pu. *J. Chem. Phys.* **2010**, *132*. <https://doi.org/10.1063/1.3382344>.
- (124) Grimme, S.; Ehrlich, S.; Goerigk, L. Effect of the Damping Function in Dispersion Corrected Density Functional Theory. *J. Comput. Chem.* **2011**, *32*, 1456–1465. <https://doi.org/10.1002/jcc.21759>.
- (125) Caldeweyher, E.; Bannwarth, C.; Grimme, S. Extension of the D3 Dispersion Coefficient Model. *J. Chem. Phys.* **2017**, *147*, 0–7. <https://doi.org/10.1063/1.4993215>.



**Table 1.** Experimental and calculated  $^{35}\text{Cl}$  EFG and chemical shift tensor parameters.<sup>a-f</sup>

		$C_Q$ (MHz)	$\eta_Q$	$\delta_{\text{iso}}$ (ppm)	$\Omega$ (ppm)	$\kappa$	$\alpha$ (°)	$\beta$ (°)	$\gamma$ (°)
Eph	Exp.	1.23(2)	0.43(3)	43(1)	65(5)	-0.7(1)	22(5)	50(5)	2(20)
	Calc.	1.110	0.44	42	65	-0.13	237	62	78
Pse	Exp.	2.20(2)	0.85(3)	41(1)	70(10)	0.3(2)	30(30)	30(5)	150(30)
	Calc.	-2.774	0.45	36	74	0.89	128	83	237

<sup>a</sup>Theoretical EFG tensor parameters were obtained from calculations on XRD-derived structural models that were refined at the RPBE-D2\* level.

<sup>b</sup>The experimental uncertainties in the last digit for each value are indicated in parentheses.

<sup>c</sup>The principal components of the EFG tensors are defined such that  $|V_{33}| \geq |V_{22}| \geq |V_{11}|$ . The quadrupolar coupling constant and asymmetry parameter are given by  $C_Q = eQV_{33}/h$ , and  $\eta_Q = (V_{11} - V_{22})/V_{33}$ , respectively. The sign of  $C_Q$  cannot be determined from the experimental  $^{35}\text{Cl}$  spectra.

<sup>d</sup>The principal components of the chemical shift tensors are defined using the frequency-ordered convention, with  $\delta_{11} \geq \delta_{22} \geq \delta_{33}$ . The isotropic chemical shift, span, and skew are given by  $\delta_{\text{iso}} = (\delta_{11} + \delta_{22} + \delta_{33})/3$ ,  $\Omega = \delta_{11} - \delta_{33}$ , and  $\kappa = 3(\delta_{22} - \delta_{\text{iso}})/\Omega$ , respectively.

<sup>e</sup>The Euler angles  $\alpha$ ,  $\beta$ , and  $\gamma$  define the relative orientation of the EFG and chemical shift tensors using the ZY'Z'' convention for rotation.

<sup>f</sup>Calculated  $^{35}\text{Cl}$  magnetic shielding constants were converted to the chemical shift scale through a separate calculation on a geometry optimized structural model of L-histidine HCl·H<sub>2</sub>O (based on a structure from the CSD, code HSTCM01) for which the experimental chemical shift is set to 34.5 ppm (see ref #1).



**Table 2.** H...Cl hydrogen bond networks for the DFT-D2\* refined crystal structures of Eph and Pse, as shown in **Fig. 3**.

	$r_1$ (Å) <sup>a</sup>	$r_2$ (Å) <sup>a</sup>	$r_3$ (Å) <sup>a</sup>	$\varphi_{12}$ (°) <sup>a</sup>	$\varphi_{13}$ (°) <sup>a</sup>	$\varphi_{23}$ (°) <sup>a</sup>	Avg. $\varphi$
Eph	2.184	2.210	2.212	82.68	83.90	79.37	81.98
Pse	2.142	2.214	2.358	95.44	91.68	77.87	88.33
24-317 <sup>b</sup>	2.159	2.198	2.263	90.06	88.66	84.21	87.64

<sup>a</sup> For definitions of geometrical parameters, see **Fig. 3**.<sup>b</sup> Structure 24-317 is one of the six structures predicted by QNMRX-CSP; see §3.2.3 for relevant discussion.

**Table 3.** The benchmarking of QNMRX-CSP featuring different starting points and conditions.

Section #	Origin of Motion Group		PXR		SSNMR
	Known crystal structure	Isolated molecule	Unit cell Parameters	Space group	<sup>35</sup> Cl EFGs
3.2.1	✓	-	✓	✓	✓
3.2.2	-	✓	✓	✓	✓
3.2.3	✓	-	✓	✓	✓
3.2.4	-	✓	-	-	✓





**Table 4.** A summary of the initial and final numbers of structural models in each step of the QNMRX-CSP protocol, as applied to Eph and Pse.

QNMRX-CSP Protocol	Known crystal structures <sup>a</sup>			Refined isolated molecule <sup>b</sup>		
	Initial		Retained	Initial		Retained
<b>Eph</b>						
M2: $\pm 20\%$ UCPs; $E_{\text{low}} \leq 13.5\%$	46889	→	702	56673	→	901
M3 Step 1: $\Gamma_{\text{EFG}} \leq 0.70$ MHz; $E_{\text{lat}} \leq 135$ kJ mol <sup>-1</sup>	702	→	25	901	→	9
M3 Step 2: $\Gamma_{\text{EFG}} \leq 0.49$ MHz; $E_{\text{lat}} \leq 50$ kJ mol <sup>-1</sup>	25	→	19	9	→	9
M3 Step 3: $\Gamma_{\text{EFG}} \leq 0.49$ MHz; $E_{\text{lat}} \leq 1$ kJ mol <sup>-1</sup>	19	→	6	9	→	1
<b>Pse</b>						
M2: $\pm 20\%$ UCPs; $E_{\text{low}} \leq 13.5\%$	38718	→	242	177564	→	922
M3 Step 1: $\Gamma_{\text{EFG}} \leq 0.70$ MHz; $E_{\text{lat}} \leq 135$ kJ mol <sup>-1</sup>	242	→	4	922	→	22
M3 Step 2: $\Gamma_{\text{EFG}} \leq 0.49$ MHz; $E_{\text{lat}} \leq 50$ kJ mol <sup>-1</sup>	4	→	3	22	→	10
M3 Step 3: $\Gamma_{\text{EFG}} \leq 0.49$ MHz; $E_{\text{lat}} \leq 1$ kJ mol <sup>-1</sup>	3	→	2	10	→	1

<sup>a</sup> The starting point for these calculations were geometry optimized structures based on a known crystal structure.

<sup>b</sup> The starting point for these calculations were geometry optimized structures based on isolated, gas-phase molecules.



**Table 5.** Validation of the structural models of Eph and Pse obtained from QNMRX-CSP.

	Motion Group <sup>a</sup>	Structural Model	$\Gamma_{\text{EFG}}$ (MHz)	$E_{\text{lat}}$ (kJ mol <sup>-1</sup> )	$R$ (%)	RMSD (Å)
Eph	OPT-EPHECL02	5-98	0.102	0.000	2.26	0.026
	OPT-EPHECL02	3-12 <sup>b</sup>	0.086	0.008	17.75	0.169
	OPT-EPHECL02	3-2 <sup>b</sup>	0.092	0.019	18.03	0.174
	OPT-EPHECL02	6-11 <sup>b</sup>	0.085	0.023	17.97	0.167
	OPT-EPHECL02	5-3 <sup>b</sup>	0.087	0.030	17.97	0.173
	OPT-EPHECL02	5-7 <sup>b</sup>	0.070	0.082	18.22	0.174
Pse	Isolated molecule	10-1	0.113	0.000	2.33	0.027
	OPT-PEPHCL	5-307	0.387	0.000	1.19	0.011
	OPT-PEPHCL	8-117	0.393	0.029	1.94	0.014
	Isolated molecule	44-278	0.408	0.000	1.56	0.013

<sup>a</sup> The starting point for each motion group was either a known crystal structure (OPT-EPHECL02 or OPT-PEPHCL), or a refined isolated molecule.

<sup>b</sup> Structural models that failed validation because of poor agreement with PXR. D.



**Table 6.** A summary of the initial and final numbers of structural models in each step of the QNMRX-CSP protocol, as applied in the benchmarking starting from molecular fragments mismatched with experimental data.

QNMRX-CSP Protocol			
<b>Pse Molecular Fragments and Eph Metrics</b>	Initial		Retained
M2: $\pm 20\%$ UCPS; $E_{\text{low}} \leq 13.5\%$	505475	→	2993
M3 Step 1: $\Gamma_{\text{EFG}} \leq 0.70$ MHz; $E_{\text{lat}} \leq 135$ kJ mol <sup>-1</sup>	2993	→	3
M3 Step 2: $\Gamma_{\text{EFG}} \leq 0.49$ MHz; $E_{\text{lat}} \leq 50$ kJ mol <sup>-1</sup>	3	→	0
M3 Step 3: $\Gamma_{\text{EFG}} \leq 0.49$ MHz; $E_{\text{lat}} \leq 1$ kJ mol <sup>-1</sup>	0	→	0
<b>Eph Molecular Fragments and Pse Metrics</b>			
M2: $\pm 20\%$ UCPS; $E_{\text{low}} \leq 13.5\%$	115734	→	795
M3 Step 1: $\Gamma_{\text{EFG}} \leq 0.70$ MHz; $E_{\text{lat}} \leq 135$ kJ mol <sup>-1</sup>	795	→	74
M3 Step 2: $\Gamma_{\text{EFG}} \leq 0.49$ MHz; $E_{\text{lat}} \leq 50$ kJ mol <sup>-1</sup>	74	→	57
M3 Step 3: $\Gamma_{\text{EFG}} \leq 0.49$ MHz; $E_{\text{lat}} \leq 1$ kJ mol <sup>-1</sup>	57	→	6



**Table 7.** Validation of the structural models of Eph obtained from QNMRX-CSP when using the Eph fragment as a structural unit, but the XRD and SSNMR data for Pse as metrics.

Structural Model	$\Gamma_{\text{EFG}}$ (MHz)	$E_{\text{lat}}$ (kJ mol <sup>-1</sup> )	$R$ (%)	RMSD (Å)
24-317	0.481	0.000	95.87	0.776
14-129	0.283	0.010	88.14	0.787
17-119	0.282	0.029	87.90	0.788
19-89	0.283	0.193	86.26	0.787
14-92	0.271	0.251	86.21	0.785
24-152	0.287	0.357	88.32	0.788



**Table 8.** The number of structures at each step and for each space group searched in the QNMRX-CSP protocol as applied to Sudafed.

	$P2_1/c$	$P\bar{1}$	$P2_12_12_1$	$P2_1$	$Pna2_1$
# MC-SA <sup>a</sup>	1000	1000	1000	1000	1000
# Cluster <sup>b</sup>	362	314	190	190	190
# TGO <sup>c</sup>	1	0	6	0	0
# CGE <sup>d</sup>	0	0	6	0	0
#UCP CGE <sup>e</sup>	0	0	6	0	0
# Validation <sup>f</sup>	0	0	6	0	0

<sup>a</sup> Initial number of structures obtained from 10 trials of Monte-Carlo simulated annealing.

<sup>b</sup> Number of structures following clustering.

<sup>c</sup> Remaining structures following truncated geometry optimization and application of the <sup>35</sup>Cl EFG distance with a cut-off of 0.70 MHz.

<sup>d</sup> Remaining structures following convergent geometry optimization and application of the <sup>35</sup>Cl EFG distance with a cut-off of 0.49 MHz.

<sup>e</sup> Remaining structures following full volume convergent geometry optimization and application of the <sup>35</sup>Cl EFG distance with a cut-off of 0.49 MHz.


<sup>f</sup> Number of candidate structures matching of the refined crystal structure of Pse (CSD code PEPHCL).



**Table 9.** Validation of the structural models of Sudafed obtained from QNMRX-CSP.


	<sup>35</sup> Cl EFG Dist. (MHz)	Energy Diff. (kJ/mol)	Volume (Å <sup>3</sup> )	Volume Diff. (%)	RMSD (Å)	R% (%)
PEPHCL	-	-	1124.871	-	-	-
2-75	0.233	0	1138.202	1.19	0.073	89.61
7-62	0.204	0.029	1141.548	1.48	0.078	94.63
8-91	0.207	0.048	1139.949	1.34	0.076	94.54
8-71	0.191	0.067	1141.009	1.43	0.079	97.00
8-86	0.182	0.077	1140.464	1.39	0.080	91.70
8-85	0.202	0.096	1141.099	1.44	0.080	97.80





**Scheme 1.** Molecular diagrams and atomic numbering for (1*R*,2*S*)-(-)-ephedrine HCl (A) and (1*S*,2*S*)-(+)-pseudoephedrine HCl (B).






**Figure 1.**  $^1\text{H} \rightarrow ^{13}\text{C}$  CP/MAS NMR spectra of (A, red) Eph, (B, blue) Pse, and (C, green) Sudafed. Field strengths and MAS rates ( $\nu_{\text{rot}}$ ) are shown to the right of each spectrum.









**Figure 2.**  $^{35}\text{Cl}\{^1\text{H}\}$  NMR spectra of Eph (red) and Pse (blue) acquired at 21.1 T under MAS (Bloch decay) and static conditions (quadrupolar echo), and at 9.4 T under static conditions (Hahn echo). Spectral simulations are shown in black.






**Figure 3.** Schematic of the H $\cdots$ Cl hydrogen bond network for the DFT-D2\* refined crystal structures of Eph and Pse. From shortest to longest,  $r_1$ ,  $r_2$ , and  $r_3$  correspond to the H $\cdots$ Cl bond lengths with secondary amine, alcohol, and another secondary amine moieties, respectively. Angles  $\varphi_{ij}$  describe the interbond angles. Values are shown in **Table 3**.






**Figure 4.** Motion groups assigned in Module 1 of the QNMRX-CSP protocol for benchmarking calculations, with structural models for Eph (top) and Pse (bottom) based on the known crystal structures (left) and structural models obtained from refining isolated molecules (right).






**Figure 5.** A comparison of the convergent geometry optimized crystal structure of Eph to structures that were validated from benchmarking by (A) using known structures (5-98), or (B) starting from structural models obtained from geometry optimizations of isolated molecules (10-1).






**Figure 6.** A comparison of the convergent geometry optimized crystal structure of Pse to structures that were validated from benchmarking by (A) using known structures (5-307), or (B) starting from structural models obtained from isolated molecule geometry optimizations (44-278).





**Figure 7.** A comparison of the convergent geometry optimized known structure of Pse and the structural model with the lowest energy that reached validation of the using the Eph cation and the Pse unit cell parameters and  $^{35}\text{Cl}$  EFG tensors (24-317).






**Figure 8.**  $^{35}\text{Cl}\{^1\text{H}\}$  SSNMR spectra of Sudafed (green) acquired at 18.8 T under (A) MAS at  $\nu_{\text{rot}} = 10$  kHz using a Bloch decay experiment and (B) static conditions using a Hahn echo experiment. Simulated spectra (black) of Pse are displayed above each spectrum. A peak corresponding to NaCl(s) is indicated with an asterisk (\*).







**Figure 9.** A view along each crystallographic axis for a crystal structure of Pse obtained by QNMRX-CSP, 8-85, and the convergent geometry optimized crystal structure of Pse.





## Data availability

The pulse sequences used and the CASTEP data manager (CDM), an executable application developed in-house, are available on the web from <https://github.com/rschurko>.

Additional data are available in the ESI.†

