

Unconventional Parahydrogen-Induced Hyperpolarization Effects in Chemistry and Catalysis: From Photoreactions to Enzymes

Andrey N. Pravdivtsev,* Ben J. Tickner, Stefan Glöggler, Jan-Bernd Hövener, Gerd Buntkowsky, Simon B. Duckett, Clifford R. Bowers, and Vladimir V. Zhivonitko*



Cite This: *ACS Catal.* 2025, 15, 6386–6409



Read Online


ACCESS |

Metrics & More

Article Recommendations

ABSTRACT: Nuclear spin hyperpolarization utilizing parahydrogen has the potential for broad applications in chemistry, catalysis, biochemistry, and medicine. This review examines recent chemical and biochemical insights gained using parahydrogen-induced polarization (PHIP). We begin with photoinduced PHIP, which allows the investigation of short-lived and photoactivated catalysis. Next, we review the partially negative line effect, in which distinctive line shape helps to reveal information about rapid exchange with parahydrogen and the role of short-lived catalytic species. The NMR signal enhancement of a single proton in oneH-PHIP is discussed, challenging the underpinning concept of the necessity of pairwise hydrogenation. Furthermore, we examine metal-free PHIP facilitated by frustrated Lewis pair molecular tweezers and radicaloids, demonstrating alternative routes to hydrogenation. Although symmetric molecules incorporating parahydrogen are NMR silent, we showcase methods that reveal hyperpolarized states through post-hydrogenation reactions. We discuss chemical exchange processes that mediate polarization transfer between parahydrogen and a molecular target, expanding the reach of PHIP without synthesizing specialized precursors. We conclude this review by highlighting the role of PHIP in uncovering the H_2 activation mechanisms of hydrogenases. By providing a detailed review of these diverse phenomena, we aim to familiarize the reader with the versatility of PHIP and its potential applications for mechanistic studies and chemical analysis.

KEYWORDS: parahydrogen, catalysis, hyperpolarization, mechanisms, NMR



1. INTRODUCTION

Without hesitation, nuclear magnetic resonance (NMR) is one of the leading and most widely used spectroscopic methods in chemistry. This success is due to its broad analytical capabilities and noninvasive nature, enabled by the minuscule magnetic moments associated with the nuclear spins. Despite its ubiquitous role in molecular characterization and analysis, NMR suffers from low sensitivity, as only a tiny fraction of all available nuclear spins effectively contributes to the total NMR signal. Consequently, combinations of time-consuming signal averaging, concentrated samples (>mM), and elaborate equipment/hardware are often necessary. Addressing these challenges is the focus of much attention, and the scope of NMR applications is constantly expanding through the introduction of more sensitive NMR instruments with higher magnetic fields,¹ cryoprobes,² ultrafast sequences,³ and nuclear spin hyperpolarization methods.⁴

In this review, we focus on one of many hyperpolarization methods that increase nuclear spin polarization using various physical and chemical effects. Such processes create transient non-Boltzmann population distributions across closely spaced nuclear spin energy levels. This effect is highly beneficial as NMR signal enhancements of several orders of magnitude can

be obtained. For example, at a clinical field strength of 1.5 T, only 0.000125%, about one in 800,000 ^{13}C nuclei, is effectively detected (thermally polarized under ambient conditions). When the system is fully polarized, all ^{13}C spins contribute to the signal, enhancing the signal—and thus the sensitivity—by more than 5 orders of magnitude. Hence, hyperpolarization is a versatile tool that aids the magnetic resonance detection of molecules at low concentrations or short lifetime, with diverse applications in biomedical imaging,^{5–8} protein studies,^{9–11} composition analysis^{12,13} and catalysis.^{14–16}


Parahydrogen-induced polarization (PHIP), introduced in the 1980s,^{17–19} exploits the alignment of the nuclear spins in the singlet state spin isomer of molecular hydrogen, parahydrogen (pH_2).²⁰ pH_2 reflects the lowest energy spin isomer of dihydrogen and can be easily prepared in an almost pure state by cooling hydrogen gas to 20 K or below,^{21–24} though

Received: December 20, 2024

Revised: February 21, 2025

Accepted: February 25, 2025

Published: April 4, 2025



77 K is often used for practical simplicity and yields a pH_2 enrichment of about 50%.^{25–27} Para-enriched H_2 gas can typically be stored for weeks at ambient temperatures and used on demand to generate nuclear spin hyperpolarization. However, as the para-state is associated with singlet nuclear spin order, it does not yield any NMR signal and is considered “NMR silent”; further chemical reactivity breaking the symmetry of pH_2 is required to enhance the NMR signal.^{17–19}

The detailed spin dynamics-based theory underlying PHIP and related phenomena are well described elsewhere.^{20,28–31} Here, we aim to provide an accessible narrative that, while linking appropriate physicochemical descriptions, does not overburden the reader with overly advanced spin physics. In classical PHIP experiments,^{17–19} which are also termed hydrogenative PHIP (hPHIP), two protons of pH_2 are typically added at vicinal positions to double or triple CC bonds in an unsaturated substrate through catalytic hydrogenation (Figure 1a). As the spin states in the newly formed




Figure 1. hPHIP and SABRE. Schematics of (a) hPHIP, whereby parahydrogen (pH_2) is added to an unsaturated bond in the presence of a catalyst, [cat], and (b) SABRE, where a ligating substrate, Sub, and pH_2 undergo reversible exchange at a catalyst. The typical SABRE auxiliary ligand L is IMes = 1,3-bis(2,4,6-trimethylphenyl)-1,3-dihydro-2H-imidazol-2-ylidene.

molecule evolve, the 1H NMR signal can be significantly enhanced. In this regard, two experimental schemes are distinguished: parahydrogen and synthesis allow dramatically enhanced nuclear alignment (PASADENA), where the hydrogenation process takes place at a high magnetic field,¹⁸ and adiabatic longitudinal transport after dissociation engenders net alignment (ALTADENA), where hydrogenation occurs at low-field.³² In both cases, the resultant 1H hyperpolarization

can be used as is or transferred to another nucleus, such as ^{13}C and ^{15}N , e.g., for observation to benefit from longer relaxation times, no background signals, and greater chemical shift dispersion relative to 1H . While hPHIP has been most commonly achieved by homogeneous Rh or Ru catalysts,^{33–36} other metals^{37–42} and heterogeneous catalysts have also exhibited PHIP,^{43,44} although typically with smaller polarization than homogeneous systems.

For alkyne substrates, *cis*-vicinal hydrogenation is the most common type of hydrogenation achieved by homogeneous metal catalysts, and, therefore, it is a common reaction used in hPHIP (Figure 1a). Recently, however, *trans*-vicinal hydrogenation was turned into a dominant pathway to unlock direct hyperpolarization of fumarate^{45,46}—an intermediate in the citric acid cycle—and a marker of cell necrosis.⁴⁷ In contrast, heterogeneous catalysts (e.g., supported metal nanoparticles) typically yield low hydrogenation selectivity as the reaction mechanism can involve multiple metal sites that deliver their hydrogen atoms to various positions in the product, either directly or reversibly.^{15,44,50,51} The convenience of using heterogeneous catalysts is now facilitating the production of hyperpolarized gases as pH_2 -enhanced contrast agents for lung MRI.^{48,49}

In general, hPHIP experiments can also be used to quantify the precise stereo and regioselectivity in such processes through the signal amplification effect.^{50–53} Regardless of the hydrogenation mechanism, a crucial requirement to observe hPHIP is the pairwise addition of H_2 to a metal catalyst or unsaturated precursor. In other words, both 1H spins of the same pH_2 molecule must end up in the same product molecule to retain their spin correlation. This requirement has provided significant insight into metal-based oxidative addition reactions.⁵⁴

An alternative to hPHIP is non-hydrogenative PHIP (nhPHIP, Figure 2), with its most popular form called signal amplification by reversible exchange (SABRE, Figure 1b).⁵⁵ In SABRE, pH_2 and a substrate ligate transiently with a metal center. Depending on the experimental conditions, *J*-couplings or cross-relaxation can then drive the conversion of the spin alignment of the pH_2 -derived spins into the spin polarization of the transiently bound substrate in the resulting complex. Subsequent dissociation of this ligand results in a chemically

Type of chemical reaction used to generate PHIP		
Substrate- pH_2 -relationship	Catalytic hydrogenation of an unsaturated substrate with pH_2	Transient interaction of a substrate with pH_2 on the catalyst
	Pairwise pH_2 addition	- hPHIP (including PHIP-SAH) - photo-PHIP (hydrogenation of substrate) - Reversible exchange of two H atoms in the catalyst (photo-PHIP, MF-PHIP, SABRE, photo-SABRE) - Pairwise replacement of two H atoms in the substrate with pH_2 - PNL
	Non-pairwise pH_2 addition	- oneH-PHIP - MF-PHIP - NEPTUN - SWAMP*
	Relayed polarization transfer	- PHIP-X (also EX, RELAY) - SABRE-Relay

Figure 2. Classification of chemical effects that lead to PHIP. The primary scope of this review is to describe the pH_2 -derived hyperpolarization effects except those of routine hPHIP and SABRE, which are well-reviewed. *Preliminary assignment based on current data. Adapted with permission from Emondts et al.¹¹⁰ Copyright 2018 John Wiley and Sons.




Figure 3. Examples of photo-PHIP. (a) The outline of a photo-PHIP experiment in which light-induced ligand dissociation creates a metal complex [M] that can add pH_2 to create an inequivalent metal dihydride species with PHIP-enhanced hydride NMR signals. (b) Depiction of laser-pump NMR-probe spectroscopy in which laser irradiation stimulated $H_2 - pH_2$ exchange followed by NMR detection (where θ is typically 45° or 90°), with a variable delay between the two (top left). The NMR spectrum at the top right is a single-laser shot, single 90° RF pulse, pH_2 -enhanced $^1H\{^{31}P\}$ NMR spectrum of the hydride region of $[Ru(PPh_3)_3(CO)(H)_2]$ with $\tau = 0.05$ ms (broadband ^{31}P decoupling). When the delay is on the millisecond time scale, the zero quantum coherences of pH_2 in the chemically inequivalent dihydride complex can evolve due to the chemical shift difference from unobservable $\hat{I}_x\hat{S}_x + \hat{I}_y\hat{S}_y$ to observable $\hat{I}_y\hat{S}_x - \hat{I}_x\hat{S}_y$. The lower kinetic traces show the hydride signal integral at $\delta \sim -6.5$ ppm from $^1H\{^{31}P\}$ NMR spectra of $[Ru(PPh_3)_3(CO)(H)_2]$ (1×10^{-3} M, 3 bar pH_2) as a function of τ . Experimental points are shown in blue with fitted red lines. This spin evolution can give information about kinetic hydrogen addition rates. (c) Slow photo-PHIP leading to hyperpolarized styrene and ethylbenzene after $[Ru(H)_2(PPh_3)_3(CO)]$ and an iridium photosensitizer (PS) are irradiated at 420 nm in DCM- d_2 at 298 K.¹²⁸ In these cases, delays between laser irradiation and NMR pulse acquisition are on the order of seconds. Consequently, ZQC coherences are not observed, which is similar to the traditional PASADENA effect. (d) Depiction of photo-SABRE in which a SABRE-catalyst hyperpolarizes *cis*-azobenzene (ABZ) followed by light irradiation to switch between *cis* and *trans* isomers of ABZ (upper). Hyperpolarized molecules are shown with a gray background.¹²⁹ Example 1H (lower left) and ^{15}N (lower right) NMR spectra were recorded after 10 min of light irradiation of a sample containing 56 mM of $^{15}N_2$ -ABZ in CD_3OD with 1 mM of $[IrCl(COD)(IMes)]$ and 200 mM DMSO- d_6 at 9.4 T. SABRE polarization transfer to 1H was performed at 200 nT and that to ^{15}N , at 400 nT. Signals for *cis*-ABZ are shown with a red background, with those of *trans*-ABZ with a blue background. (a) Adapted with permission from Procacci et al.¹¹⁵ Copyright 2016 Royal Society of Chemistry. This publication is licensed under CC BY-NC 3.0. (b) Adapted from Torres et al.¹¹⁴ Copyright 2014 American Chemical Society. This publication is licensed under CC-BY. (c) Adapted from Brown et al.¹²⁸ Copyright 2022 American Chemical Society. This publication is licensed under CC-BY 4.0. (d) Adapted with permission from Kiryutin et al.¹²⁹ Copyright 2024 John Wiley and Sons.

unaltered but hyperpolarized free substrate in solution,⁵⁶ allowing for multiple contacts and continuous polarization of the same molecule.^{57–59} Typically, the most efficient polar-

ization transfer in SABRE is reached when there is a match between certain energy levels in the SABRE complex spin system, and this can be readily achieved by selecting

appropriate DC or AC magnetic fields for the experiment.^{60–63} Such a matching condition is more precisely termed level anticrossing (LAC), and its theoretical description in the context of hyperpolarization is well-reviewed.⁶⁴

Several specific requirements must be met for molecules to become hyperpolarized by PHIP. For hPHIP, these include the presence of an unsaturated functionality (typically alkene or alkyne groups) that can accept pH₂. The strategy to hyperpolarize molecules that do not possess an appropriate unsaturated bond^{65–72} or ligating functionality^{73,74} is to add such a function on a side arm. The resulting side arm is then released after hydrogenation and hyperpolarization transfer by hydrolysis (e.g., PHIP-SAH).^{66–72} For SABRE, the substrate and pH₂ must ligate to a metal center, form spin–spin interactions, and dissociate. Therefore, such substrates often contain an electron-donating nitrogen site.^{55,75} However, the ongoing design of new catalysts has significantly expanded the types of molecules amenable to SABRE to include biologically significant O-donor ketoacids.⁷⁶ Other studies have seen sulfur, phosphorus, and silicon donor sites employed.^{77–79}

Collectively, hPHIP,^{48,80–85} PHIP-SAH,^{86–88} and SABRE^{89,90} have led to *in vivo* metabolic imaging applications. The translation from optimization of PHIP to *in vitro* and *in vivo* studies is not straightforward but has accelerated in recent years, demonstrating PHIP as a viable route to produce preclinical and clinical hyperpolarized agents in liquid and gas phases competitive with other approaches such as dissolution dynamic nuclear polarization (dDNP)⁸⁶ or spin-exchange optical pumping (SEOP).^{91–93} Advances in PHIP for *in vivo* imaging have been well-reviewed.^{4,94–97} This progress is underpinned by several advancements in the instrumentation associated with PHIP delivery.⁹⁸

In addition to the relatively well-known protocols of hPHIP and SABRE, there are other exciting effects that, although less well appreciated, more generally illustrate the unique properties of pH₂, which in turn provide valuable insight into both chemical and catalytic reactivity.²⁹ For example, one less common type of hPHIP is geminal hydrogenation, where two protons of pH₂ bind to the same carbon.^{99,100} There are other variations of the PHIP effect, such as when the catalyst itself interacts with pH₂ and becomes transiently polarized.^{101–105} Other cases include pairwise replacement of two substrate protons with a proton pair from pH₂ with no net hydrogenation,^{16,50,106,107} the addition of only one proton from a pH₂ molecule to the substrate (oneH-PHIP)¹⁰⁸ (after the formation of an intermediate by pairwise addition) or even hydrogenation accompanied by oligomerization.¹⁰⁹

This review will focus on such more unusual PHIP effects beyond PASADENA, ALTADENA, and SABRE (Figure 2). We will discuss these phenomena and their mechanisms, seeking to promote new analytical applications and provide valuable insight into the underlying chemical interactions. In this regard, we will discuss photoinduced PHIP and SABRE, partially negative line (PNL) effects and their implications for the analysis of short-lived intermediates, oneH-PHIP effects in hydroformylation, the various mechanisms of hyperpolarization of water using pH₂, PHIP effects in metal-free hydrogenation (MF-PHIP) reactions, secondary transformation which reveal hidden PHIP, chemically relayed polarization transfer, and PHIP in enzymatically catalyzed hydrogenation reactions.

2. UNCONVENTIONAL VARIANTS OF PHIP

2.1. Photo-PHIP and Photo-SABRE. Breaking the bond in pH₂ is a critical step in any PHIP-type experiment, as it unlocks the spin order of pH₂, ideally without losing the initial spin alignment. This step is most commonly associated with the oxidative addition of pH₂ to a metal center. An essential feature of this step, therefore, is the involvement of metal complex species with appropriate electron configurations (preferably diamagnetic) that both prevent rapid nuclear spin relaxation and support the formation of two new metal–hydride bonds. Accordingly, many stable complexes must undergo ligand loss before such a process occurs. This has given rise to studies where light irradiation stimulates ligand loss from a stable metal complex to generate a reactive intermediate that can react rapidly with pH₂ (Figure 3a).^{111–117} Consequently, hyperpolarized metal complexes are formed in a process that is somewhat analogous to hPHIP, which involves pH₂ addition to a stable 16-electron transition metal precursor. However, combining photochemistry with PHIP, termed photo-PHIP, can create magnetic states that differ from those created under conventional PASADENA-type PHIP.^{18,118,119} This photochemical approach has been used to study rapid kinetics^{112,115} and detect reaction intermediates or products.^{111,115,120} The examples of photo-PHIP typically involve Ir and Ru carbonyl precursors containing phosphine, diphosphine, or diarsine ligands.^{116,117,120,121}

To introduce photo-PHIP in further detail, it is convenient to provide theoretical considerations that illustrate the basic differences between traditional and photochemistry-assisted approaches. For reference, pH₂ exists as a nuclear spin singlet state that is defined by the density matrix described in eq 1

$$\hat{\rho}_{\text{pH}_2} = \frac{1}{4} - (\hat{I}_x\hat{S}_x + \hat{I}_y\hat{S}_y + \hat{I}_z\hat{S}_z) \quad (1)$$

where $\hat{I}_i$ and $\hat{S}_i$ ($i = x, y, z$) are spin operators which refer to identical atoms in pH₂. This state consists of the longitudinal two-spin order $\hat{I}_z\hat{S}_z$ and the in-phase zero-quantum coherence (ZQC) $\hat{I}_x\hat{S}_x + \hat{I}_y\hat{S}_y$. In hPHIP, numerous catalytic hydrogenation steps, or reversible H₂ exchange in SABRE, commonly create weakly coupled metal-dihydrides. In this case, the in-phase ZQC oscillates due to the periodic transformation into the out-of-phase ZQC state, $\hat{I}_y\hat{S}_x - \hat{I}_x\hat{S}_y$. As a result, it averages to zero over the time of thermally initiated reactions,¹²² unless strong proton decoupling is applied^{83,123} or experiments are performed at low magnetic fields when metal-dihydrides are strongly coupled.¹²⁴

In contrast, the photo-PHIP approach involves short laser-induced photodissociation (on the order of nanoseconds) to create a vacant ligand site such that rapid reaction with pH₂ is possible (Figure 3a). As a consequence of the relatively facile pH₂ addition (faster than $1/\nu_{\text{ZQC}}$ such that ZQC does not have enough time to oscillate) all terms in eq 1 are preserved through the hydrogenation reaction even though the singlet state is not stationary in the dihydride product complex.¹¹⁷ Due to this, the ZQC oscillation is observable in the subsequent ¹H NMR measurements (Figure 3b). For two weakly coupled spins, this oscillation of ZQC occurs at a frequency equal to the difference between the Larmor frequencies of the two nuclei, $\nu_{\text{ZQC}} = |\nu_1 - \nu_2|$, where ν_1 and ν_2 are the Larmor frequencies of the respective spins I and S. Typically, in photo-PHIP systems, there are strong interactions

between hydride protons and ^{31}P of spectator phosphine ligands, which additionally contribute to ν_{ZQC} .

At the instant the symmetry of the initial pH_2 is broken in forming the newly created hydride ligands (the time after the laser pulse is essentially $\tau = 0$ for a rapid reaction), the newly formed spin system cannot be perturbed by a hard excitation RF pulse as it is instantly formed from the NMR silent singlet state (Figure 3b). Consequently, this situation does not lead to an observable NMR signal. However, as the transverse ZQC term (eq 1) evolves under chemical shift and coupling, an NMR-observable spin state arrangement is created that can be probed directly with hard pulses (typically 45° or 90° , Figure 3b).^{113,123,125} This approach was used to study the evolution of dihydride singlet order for a wide range of Ru and Ir complexes with a variable τ between laser irradiation and NMR detection.¹¹³ This type of laser-pump NMR-probe experiment, conducted over a microsecond-to-millisecond delay time scale, allowed the observation of reactivity in an approach analogous to other laser-based UV and IR time-resolved spectroscopies, which was only possible by marrying laser-induced ligand dissociation with a pH_2 addition step (Figure 3b).^{111–117} Multiple laser pulses or continuous wave laser irradiation over times much longer than $1/\nu_{\text{ZQC}}$ have also been employed.^{115,117} However, in these cases of slow hyperpolarization preparation, spin order is averaged across the light irradiation time, which leads to an effect analogous to the time-averaged PASADENA effect as the ZQC terms are lost (Figure 3c).

In contrast to hard pulse excitations, the selective RF excitation of one of the chemically inequivalent hydride ligands in the complex can lead to the observation of strong NMR signals without the need for further spin-state evolution after the laser pulse.¹²⁶ Alternatively, adiabatic RF pulses can convert the singlet spin order into observable magnetization of one or two protons.^{125,127}

The analysis process required to extract information about the kinetics of pH_2 addition in photo-PHIP experiments is complex.^{112,114,125} For example, in cases where the pH_2 addition rate is on the same order of magnitude as the frequency of ZQC evolution, ν_{ZQC} , only partial averaging occurs. This can lead to apparent phase shifts in the signal oscillation that contain quantitative information about the pH_2 addition rate.¹¹⁵ However, if product formation is much faster than the rate of ZQC evolution, then only an upper bound for the H_2 addition rate can be determined. On the other hand, if product formation is much slower than the characteristic frequencies of the spin system evolution, the ZQC oscillations are no longer observed due to complete averaging. The latter case is similar to traditional PHIP in which thermally controlled reactions (rather than laser-induced) build up the number of H_2 addition products over a longer time window (ca. seconds).¹¹²

The photo-PHIP method has been applied to measure the H_2 addition rate of $[\text{Ir}(\text{I})(\text{PPh}_3)_2(\text{CO})]$, which is formed from laser-induced H_2 dissociation from $[\text{Ir}(\text{I})(\text{H})_2(\text{PPh}_3)_2(\text{CO})]$ (Figure 3b).¹¹⁵ Notably, the obtained rate constants were comparable to those measured using flash photolysis coupled with optical spectroscopy.¹¹⁵ However, unlike flash photolysis, NMR has additional chemical resolution that enables a more detailed analysis of chemical reactions, which is especially important for mixtures of photoactive substrates.

Approaches of this type have also been used to study ruthenium arsine complexes that act as alkyne hydrogenation

catalysts as catalytic activity is observed after photolysis, and many intermediates and species involved in the catalytic cycle are detected and characterized with the help of photo-PHIP-enhanced signals.¹¹⁶ Due to the lower reaction rate of these catalysts, these types of examples can be called slow photo-PHIP. The delay between laser irradiation and NMR signal acquisition is much longer (on the order of seconds). ZQC completely decays, revealing enhanced antiphase NMR signals analogous to those in traditional PASADENA experiments without amplitude oscillations. Initiating hydrogenation catalysis with a laser pulse allows access to hyperpolarized species inaccessible in thermal reactions (without laser-induced dissociation), as thermal conditions are insufficient to enable ligand dissociation, which must occur before pH_2 addition for many metal complexes. One recent example also involved the use of an irradiated iridium photosensitizer to produce excited $[\text{Ru}(\text{H})_2(\text{PPh}_3)_3(\text{CO})]$, which in turn stimulated H_2 dissociation to form $[\text{Ru}(\text{PPh}_3)_3(\text{CO})]$. Subsequent hydrogenation of phenylacetylene with pH_2 yielded PHIP-enhanced styrene (single hydrogenation) and ethylbenzene (double hydrogenation) with ^1H NMR signals for these organic photoactivated hydrogenation products enhanced by up to 1630 times at 9.4 T (Figure 3c).¹²⁸

While photoactivated PHIP has several examples in general, few examples of photoactivated SABRE exist. SABRE typically relies on iridium catalysts that reversibly exchange H_2 under thermal conditions. Rational catalyst design is, however, often used to improve the efficiency of SABRE by tuning ligand exchange rates or controlling the binding of particular target substrate molecules.¹⁶ One example of a type of photo-SABRE involved azobenzene, with light irradiation of the target ligand rather than the metal catalyst. This approach used light irradiation to switch azobenzene between *cis* and *trans* conformations, controlling SABRE activity as only *cis*-azobenzene has appropriate geometry for ligation to the iridium SABRE catalyst (Figure 3d).¹²⁹ Upon hyperpolarization of its ^{15}N nuclei using a traditional SABRE approach, light-induced isomerization locked the polarization within *trans*-azobenzene as it does not interact with the metal center, allowing for prolonged polarization lifetimes. Another benefit was that the *trans* isomer also formed a long-lived spin state with a lifetime of ca. 25 min, which means it can act as a store of polarization. While this example utilized a photoswitchable target, in the future, light-controlled catalysts could be exploited for SABRE, where photoactivation initiates either H_2 or substrate exchange within the metal coordination sphere. Indeed, while this review was provisionally accepted, an example of the latter was reported.¹³⁰ Developing such photo-SABRE approaches would allow the precise tuning of the scalar coupling network within the transient active catalyst to match the ligand exchange rate, thereby optimizing polarization transfer to the bound substrate. By employing photochemical control, such systems could achieve polarization levels closer to the theoretical maximum than those currently achieved through thermally driven exchange. Moreover, rather than requiring a diverse range of catalysts to accommodate different substrates and conditions, a single, photoswitchable catalyst could potentially be used. This would enhance both efficiency and selectivity while simplifying synthetic needs. Given these potential advantages, developing new classes of photo-SABRE catalysts reflects a promising avenue for advancing this hyperpolarization technique.

2.2. Reversible H₂ Exchange and the Partial Negative Line (PNL) Effect. Intra- and intermolecular chemical exchange can have a strong influence on the appearance of the resonances in an NMR spectrum, inducing line broadening and line shape distortions. Consequently, NMR line shape analysis can be employed as an analytical tool to monitor the dynamics of transient catalytic species. In the context of PHIP, such behavior is generally relevant to molecular dihydrogen complexes interacting reversibly with pH₂ and hence dissolved dihydrogen (Figure 4a), as well as other weakly interacting




Figure 4. Partial Negative Line effect induced by a catalyst–pH₂ interaction. (a) Chemical exchange of pH₂ with the metal center ML_n of the catalyst results in the PNL effect on free H₂. (b) 1H PHIP NMR spectrum recorded during the hydrogenation of Fmoc-O-allyl-tyrosine (left) to Fmoc-O-propyl-tyrosine (right) with pH₂, showing a strong PNL signal for free H₂ at 4.53 ppm. Adapted with permission from Kiryutin et al.¹³² Copyright 2017 American Chemical Society.

ligands. At the same time, since these processes involve pH₂, unlocking its magnetism introduces novel phenomena that cannot be observed with normal H₂. One intriguing effect of this kind was independently observed as an enhanced antiphase signal for dissolved H₂ by Zhivonitko et al.¹³¹ during the investigation of pH₂ activation with *ansa*-amino-boranes and by Kiryutin and Sauer et al.¹³² during the investigation of the PHIP enhancement of small oligopeptides (Figure 4b). This phenomenon is now referred to as the partially negative line (PNL) shape of the dihydrogen NMR signal.

The observation of the PNL effect is unusual because dihydrogen is a symmetric molecule that should exhibit only a single resonance in 1H NMR (A_2 spin system). However, the antiphase character of the H₂ signal in PHIP experiments implies the presence of two resonances (Figure 4b).¹³² Kiryutin et al. showed that the PNL effect is independent of the presence of a substrate and occurs via the transient interaction of the pH₂ molecules with the catalyst under fast chemical exchange.¹³² Using the double quantum coherence filter variant of the only parahydrogen spectroscopy (OPSY) NMR pulse sequence,^{133–136} it was demonstrated that the PNL effect results from the longitudinal two-spin order, $\hat{I}_z\hat{S}_z$, originating from pH₂ (eq 1). Normally, this term is NMR-invisible for a pair of magnetically equivalent protons in H₂.

However, the rapid exchange between free H₂ and two chemically inequivalent hydride ligands in a transient catalyst–dihydrogen complex (Figure 4a) leads to a slight frequency difference for two 1H resonance components of free H₂. In addition, the interaction of pH₂ with the catalyst promotes so-called singlet–triplet mixing^{119,137–139} when pH₂-originating protons become transiently inequivalent in the complex. This mixing, accompanied by the hydride–H₂ exchange, leads to the averaging of the ZQC terms derived from the initial pH₂ singlet spin order (eq 1 and discussion thereof) to zero, while the longitudinal term, $\hat{I}_z\hat{S}_z$, is preserved in both free H₂ and bound hydride ligands. Altogether, PASADENA-type exchange broadened antiphase signals are observed for dihydride ligands, while the H₂ signal is revealed as a superposition of two enhanced PNL resonance lines with opposite phases and slightly shifted frequencies. In the original work, the former species were difficult to detect directly due to heavy line broadening and their low concentration, while PNL was clearly visible and therefore useful.¹³²

Based on this explanation of the PNL effect, the partially negative line experiment (PANEL)¹³² was developed (Figure 5a), which employed continuous-wave low-power radio frequency irradiation to detect transient species indirectly by observing the change in the PNL signal of free H₂. When the frequency of the continuous radiofrequency pulse is in resonance with one of the hydrogens bound to the short-lived catalyst or free hydrogen, this nucleus is saturated, and




Figure 5. PANEL – A combination of the CEST method and the PNL effect boosts the sensitivity of catalytic intermediate detection. (a) Scheme of the PANEL (partially negative line) experiment for indirectly detecting the hidden hydrogen catalyst complex (L_nMH_2). (b) Calculated NMR spectrum of the hidden complex, magnified by a factor of 10^7 . (c) Experimental and (d) simulated PANEL spectra showing the signal of the hidden complex (lines at –16.5 ppm and –13.5 ppm). Adapted with permission from Kiryutin et al.¹³² Copyright 2017 American Chemical Society.

the PNL is strongly affected. This experiment is similar to chemical exchange saturation transfer (CEST) NMR experiments, which have also been used to study short-lived exchanging intermediates.¹⁴⁰ Using PANEL, a sensitivity gain of at least 3 orders of magnitude, compared to routine NMR experiments, is achievable (Figure 5b). As with CEST, the sensitivity and spectral resolution of PANEL depend on the amplitude of the RF saturation field.¹⁴¹ Since hydride nuclei often resonate at chemical shifts below -10 ppm, while the PNL signal for H_2 appears at around 4.5 ppm, the associated resonance separations reach several kHz in high magnetic fields (e.g., 5.8 kHz at 9.4 T). Consequently, RF pulses of 1 kHz amplitude can be applied without intrinsically distorting the H_2 signal. Hence, when used to probe the hydride region, the PANEL experiment can detect otherwise invisible intermediates with lifetimes of less than 1 ms. The achievable spectral resolution of the PANEL depends on the RF power used. Resolutions of 1 ppm are achievable by employing RF amplitudes of ca. 100 Hz.

The presence of two signals of the same intensity at -16.5 ppm and -13.5 ppm in the PANEL spectrum (Figure 5c) points to the sizable chemical shift difference between two inequivalent hydrides and the corresponding complex asymmetry. In such a weak coupling case ($|\omega_I - \omega_S|/2\pi \gg |J_{IS}|$) the H_2 resonance splitting that leads to PNL is given by eq 2¹³²

$$\Delta\nu \approx \frac{K_{eq}k_d^2}{2} J_{IS} \left(\frac{k_d^2 - (\omega_I - \omega_H)^2}{(k_d^2 + (\omega_I - \omega_H)^2)^2} + \frac{k_d^2 - (\omega_S - \omega_H)^2}{(k_d^2 + (\omega_S - \omega_H)^2)^2} \right) \quad (2)$$

where ω_I , ω_S , ω_H are frequencies of the spins in the complex (I and S) and of free dihydrogen (H), J_{IS} is the constant of spin–spin interaction in the complex, $K_{eq} = k_f[ML_n]/k_d$ is the equilibrium constant of the binding of dihydrogen to the complex, k_d is the dissociation rate constant, and k_f is the formation rate constant.

The same theoretical model of PNL¹³² reproduced the experimental NMR line-shapes, the nutation angle dependence, and the dependence on the frequency of the resonance position of the PNL. It also permitted the determination of chemical shift values for exchanging protons in the transient complex and the sign of the scalar coupling constant between those protons. Typically, hydride resonances of transient complexes are hardly observable directly in 1H NMR, whereas PNL allows for indirect but sensitive detection of their presence.¹³² In parallel to that, Johnson et al.¹⁴² observed PNL and then managed to detect a hyperpolarized dihydrogen complex by cooling the catalyst sample with pH_2 down to 238 K, confirming the presence and rapid exchange of the dihydrogen. The observation of reaction intermediates is highly valuable as such species are difficult to discern in standard 1H NMR experiments due to their low signal intensity and broad line widths. These techniques were later applied to study several exchange pathways with pH_2 .¹⁴³

Bernatowicz et al.¹⁴⁴ proposed an alternative explanation for the creation of PNL, which suggests it is caused by residual dipolar couplings (RDCs) stemming from the partial ordering of the hydrogen molecules in the external magnetic field. However, further experimental studies into this mechanism are warranted.

Czarnota et al.¹⁴⁵ and Alam et al.¹⁴⁶ investigated the conditions for a PNL effect employing iridium complexes or metal–organic frameworks (MOFs). PNL effects were found in the catalytic hydrogenation of eptifibatide, a disintegrin derivative based on a protein from the rattlesnake venom,¹⁴⁷ or trivinyl orthoacetate,¹⁰⁰ as well as in SABRE studies employing Zintl cluster-supported rhodium centers¹⁴⁸ or nickel diazadi-phosphacyclooctane complexes.¹⁴³ Finally, PNL effects are also a possible loss channel for hyperpolarization in hPHIP or SABRE applications.^{119,138,139,149} PNL not only allows for ultrasensitive detection of intermediates but can also serve in the future as a tool to monitor spin dynamics and, thus, chemical kinetics in intermediate complexes.

2.3. OneH-PHIP. Pairwise addition of pH_2 to an unsaturated substrate is the typical requirement for generating hyperpolarization in hPHIP (Figure 1), meaning that both hydrogen atoms must end up in the same product molecule and their spins remain correlated. This implies that hydrogens must follow each other throughout all catalytic steps. However, this requirement can be lifted when the pairwise addition of pH_2 to an active catalytic center leads to the formation of a dihydride intermediate where the chemical shift difference between the hydride ligands is small, and a strongly coupled spin pair therefore results. Then, if the lifetime of this intermediate dihydride complex is sufficient, the initial singlet spin order of the pH_2 -derived hydrogen pair (eq 1) can evolve into individual single-spin net polarizations ($\mp \hat{I}_z$ and $\mp \hat{S}_z$, Zeeman orders) of the hydrogen atoms that will consequently be transferred into the final reaction products. In such cases, the hyperpolarization is associated with each of the two pH_2 -derived protons separately, without requiring their spin correlation. Therefore, the initial pH_2 pair can be separated in subsequent steps of the catalytic cycle, while the hyperpolarization observed in the final product can be derived from only one hydrogen of the pair. This effect is referred to as the oneH-PHIP effect.

OneH-PHIP was first observed by Permin and Eisenberg during their stoichiometric studies of hydroformylation by platinum–tin and iridium carbonyl species.¹⁰⁸ Here, *trans*-PtCl(COEt)(PPh₃)₂ proved to react with SnCl₂ and pH_2 to form propanal, where only the slowly relaxing aldehydic proton (Figure 6a) exhibited NMR signal enhancement. This is reflective of the creation of single-spin net polarization associated with an $\hat{I}_z$ type term (indicated by H in the following products), which differs significantly from the more usual pH_2 -derived longitudinal two-spin order $\hat{I}_z\hat{S}_z$ term (the last term in eq 1), which is destroyed when the coupling between the spins is lost. As this section will illustrate, such effects are relatively common, although the signal enhancements are relatively low. For example, a signal enhancement of 5-fold at 9.4 T using 50% pH_2 was achieved for the aldehydic proton of propanal.¹⁰⁸ Hence, oneH-PHIP observation has been limited to slowly relaxing species or high-turnover catalysis.

Permin and Eisenberg investigated the catalytic production of propanal (CH_3CH_2CHO) using $PtCl_2(CO)(PPh_3)_2$ –SnCl₂¹⁰⁸ and $[Ir(COEt)(CO)_2(dppe)]$ (dppe is 1,2-bis-(diphenylphosphino)ethane) wherein only the aldehydic proton was hyperpolarized (Figure 6a). In the case of Ir systems, hyperpolarized hydride ligand signals were also detected for the intermediate $[Ir(H)_2(COEt)(CO)(dppe)]$, where the hydride ligand *cis* to the phosphines (H' in the figure) was then found to become the hyperpolarized




Figure 6. The oneH-PHIP effect has allowed the detection of several single-spin hyperpolarized products. This effect results from the creation of a hyperpolarized AB-type spin system for the $p\text{-H}_2$ -derived protons (indicated with pink H atoms in (a) and (b)), with subsequent further reaction producing single spin hyperpolarized products. Examples shown here include (a) aldehydes,¹⁵⁸ (b) and (c) metal hydride complexes,^{156,157} and (d) vinyl-containing species.¹⁵⁸ The NMR traces illustrate the typical appearance of a single spin-polarized resonance under ^{31}P decoupling in the case of (c).

aldehydic proton in the final product. Hydride signals for this intermediate appear at very close resonances (−8.696 and −8.905 ppm in benzene), forming an AB-type spin system. In Pople notation, the AB-type spin system corresponds to two spins exhibiting a chemical shift difference comparable to their spin–spin coupling. The observed hyperpolarization of only a single proton in the final aldehyde product was called oneH-PHIP.

Subsequent solvent variation of a benzene–acetone mixture led to an inversion in the oneH-PHIP phase of the aldehydic proton as a consequence of the relative change in chemical shifts of the AB spin system of the hydride ligands inverting. Hence, the oneH-PHIP effect was linked to strong coupling. In this case, a rigorous theoretical description, enunciating the role of chemical shift difference and mutual spin–spin interaction in the AB spin system that created the $\hat{I}_z$ – $\hat{S}_z$ type magnetization was reported, which set Bargon's earlier description into a firm chemical context.²⁸ However, it does not exclude the possibility of a relaxation-driven polarization transfer mechanism. For instance, in section 2.5, we discuss such mechanisms in the hydrogenation of alkynes and imines with $p\text{-H}_2$ using *ansa*-aminoborane catalysts.^{150–152} To verify the exact mechanism, one can study the magnetic field dependence of the oneH-PHIP effect: cross-correlated relaxation is expected to be stronger at higher fields, while coherent mechanisms dominate at lower fields.^{28,153,154} Another rationale for the oneH-PHIP effect appears in ref 155, although this time in the context of the hyperpolarization of water and alcohols by either homogeneous or heterogeneous catalysis, as described in section 2.4.

An additional example of a hydride resonance exhibiting oneH-PHIP was observed during catalytic studies of alkyne hydrogenation by $[\text{Pd}(\text{bcope})(\text{OTf})_2]$ (bcope = $(\text{-C}_8\text{H}_{14}\text{-1,5})\text{PCH}_2\text{CH}_2\text{P}(\text{-C}_8\text{H}_{14}\text{-1,5})$; OTf = $\text{CF}_3\text{SO}_2\text{O}^-$) where

species like $[\text{Pd}(\text{bcope})(\text{pyridine})(\text{H})](\text{OTf})$ were detected.¹⁵⁷ This study extended into the detection of critical reaction intermediates like $[\text{Pd}(\text{bcope})(\text{CPh}=\text{C}(\text{H})\text{Ph})\text{-(pyridine)}](\text{OTf})$, where the single vinyl proton exhibited strong hyperpolarization, alongside *cis*- $\text{PhH}=\text{CHPh}$ (Figure 6b). During these studies, it was the strongly coupled spin system of the reversibly formed intermediate $[\text{Pd}(\text{bcope})(\text{CHPhC}(\text{H})_2\text{Ph})](\text{OTf})$ that led to this behavior.¹⁵⁷ The related complex, the alkene insertion product, $[\text{Pd}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PCy}_2)(\text{-C}(\text{Ph})\text{H}-\text{CHPh}-\text{CPh}=\text{CH}-\text{Ph})](\text{OTf})$ has also been observed thanks to the ^1H NMR signal enhancement of oneH-PHIP¹⁵⁹ and vinyl ethers have been produced during platinum-catalyzed reactions that exhibit this effect.¹⁶⁰

Furthermore, the addition of CO to drive palladium-catalyzed carbonylation has extended the hyperpolarized observations to include the ketone $\text{MeOCO}(\text{CPh})=\text{CHPh}$ proton resonance, alongside further signals in the acyl bearing reaction intermediate $[\text{Pd}(\text{bcope})(\text{CO}-\text{CPh}=\text{C}(\text{H})\text{Ph})(\text{CO})](\text{OTf})$, the novel hydride complex $[\text{Pd}(\text{bcope})(\text{CO})(\text{H})](\text{OTf})$, the alkene complex $[\text{Pd}(\text{bcope})(\text{CHPh}=\text{CPh}(\text{COOMe}))]$ and free HD (alongside H_2) (Figure 6d).¹⁵⁸ Here again, detecting a hyperpolarized response for the released H_2 reflects the oneH-PHIP that results from strong coupling effects in species that led to it.

Later, Guan et al. reported on studies of related $[\text{Ir}(\eta^3\text{-C}_3\text{H}_3)(\text{CO})(\text{PMe}_3)_2]$ type species and described how the hydride ligand signal for $[\text{HfIr}(\text{CO})(\text{PMe}_3)_2]$ exhibited oneH-PHIP (Figure 6c).¹⁵⁶

It should be apparent from these discussions that the sharing of hyperpolarization between species can occur via numerous processes with dramatically different efficiencies. The polarization of a single proton was also reported in other reactions with H_2 involving metal-free catalysts or heterogeneous catalysts and led, among others, to a polarization of water, as discussed in the following sections.

2.4. SWAMP and NEPTUN. Hyperpolarized water is an important target molecule as it can be used for angiography and perfusion biomedical imaging^{161–163} and as a polarization source for heteronuclear signal enhancement in biomolecular NMR spectroscopy.^{10,164–168} While high polarization levels (>60%) have been shown to result from dDNP,¹⁶⁹ $p\text{-H}_2$ -based methods offer an alternate route that is both rapid and less expensive, thus making the approach more widely accessible.

Water had been an elusive target for PHIP until 2017, when it was hyperpolarized in D_2O mixtures of L-histidine and a water-soluble iridium complex, $[\text{Ir}(\text{Cl})(\text{IDEG})(\text{COD})]$ (IDEG = 1,3-bis(3,4,5-tris(diethylene glycol)benzyl)-imidazole-2-ylidene).¹⁷⁰ In this system, the hyperpolarized HDO and HD proton signals appear in the emission and absorption phases, respectively (Figure 7a,b). Consistent with the oneH-PHIP theory (see section 2.3),^{110,155} the enhanced HD and HDO signals have opposite phases and very similar field dependencies of their signal amplitudes, which reach a maximum near 45 mT where the *J*-coupling and chemical shift difference between the dihydride protons (Figure 7c) are matched. The corresponding oneH-PHIP mechanism, mediated by (i) H/D exchange with coordinated D_2O , (ii) dissociation of HDO, and (iii) H–D recombination, was named nuclear exchange polarization by transposing unattached nuclei (NEPTUN).¹¹⁰ The role of L-histidine in this work remains unclear.




Figure 7. (a) H/D exchange, in the presence of a water-soluble iridium catalyst and histidine, leading to hyperpolarization of HD and HDO and (b) the resulting NMR spectrum, in comparison to the thermally polarized spectrum. (c) The proposed “NEPTUN” mechanism underpins the spectrum. (a, b) Adapted with permission from Lehmkuhl et al.¹⁷⁰ Copyright 2017 John Wiley and Sons. (c) Adapted with permission from Emondts et al.¹¹⁰ Copyright 2018 John Wiley and Sons.

A NEPTUN-type mechanism was also suspected to be active in relayed hyperpolarization experiments where the goal was to further extend the SABRE hyperpolarization to heteronuclei in noncoordinating substrates like alcohols (e.g., methanol, ethanol) via proton exchange with a carrier amine.¹⁷¹ The possible involvement of such a mechanism, in addition to OH/NH exchange, was inferred from magnetic field dependencies of ¹³C distortionless enhancement by polarization transfer (DEPT) signals, which, in addition to showing a peak at 6.5 mT, as expected for the conventional SABRE matching condition (relayed to the target via NH/OH exchange), there is an even more prominent peak after transfer at 19.2 mT, which is hypothesized to stem from the NEPTUN effect.¹⁷¹ However, attempts to observe the hydride resonances indicative of NEPTUN directly were unsuccessful.

While most PHIP studies utilize dissolved organometallic catalysts, heterogeneous catalysis offers facile separation of the hyperpolarized products from the catalyst and can even be used in a packed-bed flow-reactor configuration.^{172,173} Supported noble metals are among the most active hydrogenation catalysts. Unfortunately, only a tiny fraction of adducts (ca. 1–5%) are formed by pairwise addition, depending on nanoparticle size and reaction conditions. Rapid H adatom diffusion and facile exchange with gaseous H₂ conspire to destroy the singlet order in the nascent H adatom pair.^{15,51,174,175} Intermetallic phases incorporating an active and inactive metal, such as Pt and Sn, allow for the tuning of molecular adsorption and diffusion dynamics through a combination of geometric and electronic effects.^{176–178} Thus, as the fraction of Sn increases across the series Pt → Pt₃Sn → PtSn, the pairwise selectivity for hydrogenation of propene increases by more than 3 orders of magnitude.¹⁷⁶ After bubbling p-H₂ through a D₂O suspension of Pt₃Sn@mSiO₂ (Pt₃Sn nanoparticles encapsulated in mesoporous silica) for 30 s, Zhao et al. observed hyperpolarization of the residual protons of solvent molecules.¹⁵⁵ The effect was dubbed surface waters are magnetized from parahydrogen (SWAMP). Proton hyperpolarization in methanol-*d*₄ and ethanol-*d*₆ was also observed. The surface properties of Pt₃Sn now balance the necessary facile H₂ activation and suppression of diffusion.

The NEPTUN and SWAMP systems share a few similarities: (i) emission phase of the HDO peak. (ii) absence of signal enhancement at zero or high magnetic field, revealing a role of Zeeman interactions; (iii) monotonic growth of [HDO] with total p-H₂ bubbling time; (iv) emergence of a dissolved HD (triplet) NMR signal. The last two are accounted for by the net isotope exchange reaction described in eq 3.



For Pt surfaces, H/D exchange is mediated by reversible electron transfer from an H adatom to the metal and proton transfer to surface water to yield a hydronium-like species where H/D exchange occurs.¹⁷⁹ However, the oneH-PHIP/NEPTUN mechanism was only tentatively excluded by preliminary data revealing the different dependences of the SWAMP signals for exchangeable and nonexchangeable protons of HOCD₃ and DOCHD₂, respectively, on the total amount of H/D exchange. Plausible mechanisms are illustrated in Figure 8a.

While the monometallic Pt@mSiO₂ nanoparticles of ref 155 were found to be inactive as SWAMP catalysts, Norcott




Figure 8. (a) Possible mechanisms underpinning surface-mediated hyperpolarization of liquid water. (b) Mechanism of benzoquinone scavenging of depolarized H adatoms on Pt/C, and its effect on the SWAMP signals. (a) Adapted with permission from Zhao et al.¹⁵⁵ Copyright 2018 Elsevier. (b) Reproduced from Norcott.¹⁸⁰ Copyright 2023 American Chemical Society. This publication is licensed under CC-BY-NC-ND 4.0.




Figure 9. An overview of metal-free PHIP (MF-PHIP). The central top part highlights that MF-PHIP effects have been demonstrated in activation of pH_2 using frustrated Lewis pairs (FLPs) and using bi- and tetra-radicaloids (BRs and TRs). Correspondingly, the structures of *ansa*-aminoborane (AAB) and *ansa*-phosphinoborane (APB) FLPs are presented in (a). AABs are referred to as molecular tweezers for pH_2 . pH_2 activation under ambient conditions using *ortho*-phenylene AABs ($n = 0$) accompanied by the corresponding hyperpolarized NMR spectra is illustrated in (b). See (a) for the definition of n . In addition to 1H , ^{15}N and ^{11}B nuclei are also hyperpolarized spontaneously at high magnetic fields in this process. Alkyne and imine hydrogenation reactions catalyzed by AABs HCAT ($n = 0$; $NR_2 = NMe_2$; $Ar = C_6F_5$; $R' = H$) and QCAT ($n = 1$; $NR_2 = THQ$; $Ar = C_6F_5$), respectively, are shown in (c). Additionally, the catalytic cycle of alkyne hydrogenation using the HCAT AAB catalyst is presented in (d). Typical 1H NMR signals of the reaction intermediate and the reaction product are shown next to the corresponding structures in the cycle. Structures of BR and TR molecules that demonstrated hyperpolarization effects in pH_2 activations are depicted in (e). Corresponding examples of enhanced 1H and ^{31}P NMR signals observed in reactions with BR molecules are shown in (f). The structures of the corresponding BR- H_2 adducts are depicted next to the spectra. Abbreviations: TMP = N-2,2,6,6-tetramethylpiperidiny; THQ = N-tetrahydroquinolinyl; iPrPh = 2-isopropylphenyl; Mes = mesityl; Cy = cyclohexyl; Ter = 2,6-dimesitylphenyl; Dmp = 2,6-dimethylphenyl; Emind = 1,1,7,7-tetraethyl-3,3,5,5-tetramethyl-s-hydrindacenyl. (d) Adapted from Zakharov et al.¹⁵⁰ Copyright 2022 John Wiley and Sons. This publication is licensed under CC-BY 3.0.

discovered that one could hyperpolarize water and methanol using a commercially available carbon-supported Pt nanoparticle catalyst when benzoquinone was added to a D_2O suspension of the catalyst.¹⁸⁰ Maximum 1H NMR signal enhancements (relative to thermal equilibrium at 1.4 T) approaching 45-fold were observed for methanol with ten equivalents of benzoquinone (w/w with respect to Pt/C). Benzoquinone is converted to hydroquinone during this process, which assists in increasing the turnover of fresh pH_2 on the surface and thereby increasing the level of polarization of methanol and water (Figure 8b).

While such signal enhancements are likely to increase with further catalyst development and optimization of experimental conditions, it remains to be seen whether the PHIP approach can rival the very high polarization levels achievable by dDNP for water.^{161,162,164–167} As pH_2 -based hyperpolarization techniques are inherently rapid, continuous, and low-cost, their use to achieve sufficient levels of water hyperpolarization could provide advantages to dDNP methods and significantly

advance the application range of hyperpolarized water in biomedical research.

2.5. Metal-Free PHIP: Molecular Tweezers and pH_2 Activators. The chemical activation of pH_2 is crucial to derive enhanced NMR signals in PHIP. Commonly, transition metal catalysts are employed to mediate such activations and produce hyperpolarized substances. At the same time, the use of metal-free activators and catalysts for pH_2 -based hyperpolarization, collectively named metal-free PHIP (MF-PHIP), is also documented.^{131,150,151,181–188} This section focuses on several types of MF catalysts, their structures, hyperpolarization effects, and their unique mechanistic features (Figure 9).

MF activations of H_2 are less common than those that rely on transition metal centers. They have recently attracted a lot of attention due to the possibility of using sustainable main-group elements to design less toxic and more environmentally friendly catalysts. MF catalysts for PHIP is an emerging research field that is still in its infancy. In this regard, frustrated

Lewis pairs (FLPs)¹⁸⁹ are the most studied class of MF activators for pH_2 . Specifically, various *ansa*-aminoborane (AAB) FLPs show pronounced hyperpolarization effects (Figure 9a). These compounds are referred to in the literature as “molecular tweezers” that stretch but do not split H_2 molecules.¹⁹⁰ Recent studies revealed that the stretched H–H bond is relatively weak, making it possible to form various rotameric forms in solution, including those with large H...H separations.¹⁸⁵ Nevertheless, AAB– H_2 adducts have motionally averaged J -coupling constants (2–4 Hz) between the ^1H – ^1H pair, allowing for PHIP effects under high field (PASADENA) conditions.^{131,185}

Unlike homolytic oxidative addition to metal centers, AABs activate pH_2 heterolytically with a clear charge separation on the Lewis acidic boron and the Lewis basic nitrogen sites (Figure 9b), although the two protons remain spin correlated. Depending on the AAB structure, hyperpolarization of ^1H , ^{11}B , and ^{15}N can be observed in simple pH_2 bubbling experiments without harnessing dedicated pulse sequences.³⁷ Signal enhancements as large as 2000-fold at 9.4 T and room temperature have been observed in the resulting ^1H NMR spectra for the pH_2 -originating protons of AAB– H_2 . The size of this ^1H NMR signal gain depends strongly on the experimental conditions and is defined by relaxation and kinetic parameters.¹⁸⁵ Density functional theory (DFT) calculations have also revealed various conformational forms of AAB adducts and their transformations.

In addition to simple pH_2 activation, *ansa*-aminoboranes can be used in catalytic hydrogenations of alkynes¹⁵⁰ and imines¹⁵¹ with pH_2 (Figure 9c). Interestingly, the catalytic cycles that lead to hyperpolarized alkenes and amines are nonpairwise, implying that the pH_2 -derived protons end up in different product molecules (Figure 9d). The hyperpolarization effects in this case are not expected to be observable in PHIP. However, due to the strong chemical shift anisotropy (CSA) of NH protons in the catalytic intermediates, a net negative polarization is generated from the pH_2 spin order through CSA–dipole–dipole cross-correlated relaxation.¹⁵¹ For instance, in alkyne hydrogenations, this mechanism is revealed by the negative in-phase resonance of the NH group of HCAT–alkyne– H_2 intermediate. As this proton transfers to the final alkene product, a two-orders-of-magnitude enhanced negative signal of one of the added protons at the double bond of the resulting alkene appears in the ^1H NMR spectra at 9.4 T. This effect is related to oneH-PHIP in hydroformylation reactions catalyzed by metals (section 2.3), but the underlying mechanisms of hyperpolarization, as well as chemical processes, are different. It is worth noting that the ability of cross-correlated relaxation to transform pH_2 spin order to a net polarization was also observed on metal complexes, e.g., by Aime et al. in pH_2 activations using Os and Ru clusters.^{191,192}

AABs are reported to be generally water-intolerant, which is a significant obstacle to the wide application of MF-PHIP. Valuable steps have been made to resolve this issue recently.¹⁹³ In addition, *ansa*-phosphinoboranes (APBs) were demonstrated to show PHIP effects in the presence of several equivalents of H_2O (Figure 9a).¹⁸⁴ Other FLPs showing hyperpolarization effects include Sn/P systems,¹⁹⁴ though it is strictly not a MF compound and will not be discussed here. In addition, aromatic triphosphabenzene was also shown to reveal hyperpolarization in the reaction with pH_2 at elevated temperatures (375 K).¹⁸¹ However, the resulting H_2 adduct is prone to decomposition.

Singlet pnictogen radicaloids are another class of MF pH_2 activators that demonstrate prominent hyperpolarization effects. Electron spins in these open-shell molecules are coupled into a singlet state that does not possess free electron angular momentum, which excludes the deleterious influence of the radical centers on the nuclear spins. Typical examples include cyclic species with P and/or As radical centers isolated by surrounding bulky substituents for stabilization. This configuration maintains high reactivity toward small molecules, such as H_2 , while preserving an open-shell structure. In the context of PHIP, biradicaloids (BRs) with four- or five-membered cycles are studied more extensively (Figure 9e), and any observed hyperpolarization effects strongly depend on BR symmetry.¹⁸⁷ With symmetric four-membered biradicaloids, pH_2 forms symmetric adducts. For instance, four-membered biradicaloids form the AA'XX' spin system, which leads to enhanced ^1H and ^{31}P NMR signals in PASADENA experiments (Figure 9f).^{186,187} Nonsymmetric five-membered species form a system with weakly coupled protons, leading to only ^1H hyperpolarization. However, the transfer of ^1H hyperpolarization to ^{31}P can be achieved using ESOTHERIC NMR pulse sequences with ^{31}P NMR signal enhancements exceeding 3 orders of magnitude at 9.4 T.^{187,195} Tetradicaloids (TRs) are represented by a single example (Figure 9e),¹⁸⁸ which showed less pronounced but interesting hyperpolarization for the addition of the first and second equivalents of pH_2 . Radicaloid systems are generally more reactive than FLPs, which makes them especially interesting for future developments that may lead to active catalysts, e.g., for hydrogenation or hydroformylation reactions.

MF-PHIP represents an exciting frontier in hyperpolarization method development. Using MF compounds such as FLPs and radicaloids introduces new mechanistic pathways and structural features that differentiate them from traditional metal-catalyzed systems. The unique hyperpolarization effects and mechanisms of MF-PHIP, including two-centered activation and nonpairwise hydrogen transfer, offer potential for innovative applications. Future research in this area promises to further enrich our understanding and utilization of these novel catalysts for hyperpolarization.

2.6. Revealing PHIP in Subsequent Chemical Transformations. The involvement of hyperpolarized molecules in subsequent chemical transformations is an interesting application of PHIP that is nicely illustrated using metabolic reactions, such as pyruvate-to-lactate conversion¹⁹⁶ and fumarate-to-malate¹⁹⁷ conversions. Other examples include oxidation of hyperpolarized pyruvate with H_2O_2 ¹⁹⁸ or decarboxylation with yttrium polyaminoacrylate adducts,¹⁹⁹ methylation of N-heterocycles,²⁰⁰ and conversion of hyperpolarized $^{15}\text{NO}_2^-$ via several reactions to make a range of products with enhanced ^{15}N NMR signals,²⁰¹ and other transformations.²⁰²

Typically, PHIP-enhanced NMR signals are visible without any subsequent reaction and can serve as a tool for kinetic measurements and identifying intermediates or additional products of the following reactions. However, if we consider symmetric molecules produced in reactions with pH_2 , the product, although in a far from thermodynamic equilibrium nuclear spin state, may not exhibit observable hyperpolarization if the pH_2 addition site is at the center of symmetry. In this case, a subsequent symmetry-breaking chemical reaction(s) may be required to reveal otherwise unobservable hyperpolarization (Figure 10a). The quintessential example of




Figure 10. (a) A general scheme of formation of compounds with hyperpolarized spins in chemical reactions of symmetric molecules, such as pH₂, ethylene, DMM, and para-¹⁵N₂, accommodating otherwise unobservable nuclear spin orders. (b) The chemical synthesis of *Z*- and *E*-ethylene from acetylene and pH₂ over various heterogeneous catalysts, as well as subsequent reactions revealing hyperpolarization. (c) The chemical synthesis of dimethyl maleate (DMM) molecules in hydrogenation with pH₂ over a cationic Rh⁺ catalyst, followed by their subsequent reaction with thiol molecules to reveal hyperpolarization. Abbreviations: DMM - dimethyl maleate.

this kind is the pH₂ molecule, which accommodates singlet nuclear spin order and does not yield NMR signals. As first shown by Bowers and Weitekamp,^{17,18} one needs to break the symmetry of pH₂ in a chemical reaction to observe hyperpolarization. Notably, symmetric molecules can typically host long-lived spin orders.^{203,204} Such molecules can be synthesized using pH₂ in hydrogenative and non-hydrogenative reactions. For example, the syntheses of ethylene from acetylene and pH₂ (Figure 10b), dimethyl maleate (DMM) from dimethyl acetylene dicarboxylate (DMAD) (Figure 10c), and para-¹⁵N₂ using SABRE were described.^{205–212}

The case of ethylene is fundamentally interesting since it, like H₂, has nuclear spin isomers of molecules (NSIMs) that differ by rotational and spin degrees of freedom due to the coupling of nuclear spin and rotational states through the symmetry properties of their respective wave functions.²¹³ Briefly, there are four NSIMs for ethylene that can be classified according to the symmetries of the nuclear spin state for the D_{2h} molecular point group using Mulliken symbols: A_g (one quintet and two singlets), B_{1u} (triplet), B_{2u} (triplet), and B_{3g} (triplet). Depending on the stereoselectivity of the hydrogenation of acetylene, *syn* and *anti* pH₂ addition products, *Z*- and *E*-ethylene, respectively, can be produced (Figure 10c),²⁰⁵ which is primarily determined by the hydrogenation catalyst employed. For instance, supported Pd nanoparticles are less selective and produce both *Z*- and *E*-ethylene products,²⁰⁵ whereas immobilized complexes of Ir are more selective, leading primarily to *Z*-ethylene.²⁰⁸ Interestingly, the subsequent reactions of ethylene produced using different catalysts with sulfonyl chlorides²⁰⁵ or Br₂/D₂O^{208,214} reveal different lifetimes of the nonequilibrium spin states in ethylene. This was rationalized based on the interconversion between

different NSIMs of ethylene,²⁰⁷ providing insights that in the gas phase, *Z*-ethylene has only one long-lived component. In contrast, *E*-ethylene has two long-lived components due to the imbalance of NSIMs with different inversion symmetry in the latter case.^{205,208} Lifetime constants of more than 15 min were measured by unlocking the hyperpolarization in the subsequent reactions for gaseous *E*-ethylene.

Another example of revealing the latent polarization inherited from pH₂ is documented for the reaction of thiols with DMM produced from DMAD in a liquid-state hydrogenation over a Rh(I) cationic catalyst (Figure 10c).²⁰⁶ As in the case of ethylene, storage of the nonequilibrium nuclear spin order after the hydrogenation was demonstrated in this study. The thiol reaction allowed for the lifetime measurement of the populated long-lived singlet spin order at a high field of up to 4.7 min. Interestingly, the unique symmetry of the DMM molecule, which induces slight magnetic inequivalence in the added pH₂-derived proton pair, allows alternative methods that do not require chemical transformation to reveal hidden singlet spin state populations. Instead, such hidden spin states can be converted into observable magnetization using magnetic field cycling²¹⁵ or applying RF fields.^{216–218} These methods do not apply to ethylene, as all its protons are magnetically equivalent.

In addition to ethylene and DMM, para-¹⁵N₂ has also been reported to form from SABRE-hyperpolarized ¹⁵N-labeled tetrazine²¹¹ and diazirines²¹² in chemical reactions involving these agents. However, the successful formation of para-¹⁵N₂ in these cases was inferred only from the absence of a ¹⁵N₂ signal in ¹⁵N NMR spectra after its production step. Similarly to pH₂, para-¹⁵N₂ is NMR silent. So far, no subsequent symmetry-breaking reaction of para-¹⁵N₂ that reveals its singlet spin order has been reported to our knowledge.

Overall, the availability of methods that use pH₂ to produce symmetric molecules with long-lived nuclear spin orders can expand the range of chemical reactions studied by providing flexible time windows. For example, it can enable one to go beyond conventional hydrogenation studies using pH₂ in PHIP, extending to electrophilic additions to double bonds, as demonstrated in the cases of ethylene and DMM. Furthermore, the analysis of the generated spin order lifetimes in molecules such as ethylene and para-¹⁵N₂ can provide essential insights into the fundamentals of NSIMs and underlying molecular physics.

2.7. Spreading Hyperpolarization via Chemical Exchange: PHIP-X and SABRE-RELAY. In recent years, a new approach has emerged to boost the NMR signal for molecules that do not contain functionalities suitable for direct PHIP or SABRE (natively or on a side arm). Two versions of this approach are based on reversible proton exchange between a hyperpolarized carrier and a to-be-hyperpolarized molecule and are termed PHIP by chemical exchange (PHIP-X)²¹⁹ and SABRE-Relay.²²⁰ In both methods, traditional PHIP or SABRE is used to polarize a “transfer” or “carrier” molecule, whose polarization is then transferred to a secondary target molecule via the exchange of hyperpolarized OH/NH protons (Figures 11 and 12a). These approaches have allowed a significant expansion of the substrate scope of PHIP and SABRE in recent years.

In the case of PHIP-X²¹⁹ (also referred to as PHIP-Relay),²²¹ propargyl alcohol, propionic acid, or propargyl amine were hydrogenated with pH₂ using a homogeneous Rh catalyst in an aprotic solvent, such as acetone, to produce a hyperpolarized “transfer” agent (Figure 11).²¹⁹ Strong spin–




Figure 11. Schematic view of parahydrogen-induced polarization by chemical exchange (PHIP-X). PHIP-X consists of four essential steps: hydrogenation of the carrier agent (step 1), the polarization of the exchanging protons (step 2), transfer of the exchanging protons from the carrier to the target molecule (step 3), and polarization of the target nucleus (step 4) using RF-induced spin order transfer technique or free evolution at low and ultralow magnetic fields. Adapted from Them et al.²²² Copyright 2024 Springer Nature. This publication is licensed under CC BY 4.0.

spin interactions distribute the polarization among the protons of the transfer agent, including the labile OH (or NH) proton. The polarization of this labile proton is relayed to the spin system of a third molecule by chemical exchange, where, again, spin–spin couplings, low magnetic field, or RF spin order transfer sequences (RF-SOT) facilitate the transfer of the polarization to other nuclei such as ^1H , ^{13}C , or ^{15}N .^{221,222} PHIP-X was shown to spontaneously polarize labile protons to ca. 0.4% for ethanol and water, 0.07% for lactic acid, 0.005% for pyruvic acid, and at least 0.009% ^{13}C polarization for glucose. Using RF-SOTs, 1.2% ^{15}N polarization was achieved for urea, where the ^{15}N coupling to the labile proton is large, 0.024% for ^{13}C glucose,²²¹ 0.026% for ^{13}C lactate²²² and ca. 0.007% for ^{13}C methanol.²²² The balance here is reached when proton exchange is slow enough to allow the J -coupling to transfer polarization to the labile proton of the carrier first and then from the labile proton to other nuclei of the target. At the same time, the exchange must be fast enough such that spin relaxation would not destroy polarization before the target is polarized. Therefore, even higher polarization values are expected to be achievable after thoroughly tuning the exchange parameters. Also, the polarization transfer is faster and more efficient when transferred to nuclei directly bound to the labile proton, such as ^{15}N or ^{13}C , with the strongest J -coupling constants. An advantage of using PHIP to hyperpolarize the transfer agent compared to SABRE is that the molecule can be polarized up to unity by the direct addition of pH_2 ; a disadvantage is that it is irreversible as the addition step can be performed only once (one addition of pH_2 per transfer agent).

In SABRE-Relay, classical SABRE is used to hyperpolarize ligating carriers, typically NH_3 and amines, with secondary NH/OH exchange effectively relaying polarization to non-ligating target substrates.^{223,224} Consequently, SABRE-Relay has been used to hyperpolarize alcohols,^{171,225} sugars,²²⁶ silanols,⁷⁹ lactate esters,²²⁷ natural products,²²⁸ and many other functional groups^{220,229} that do not interact with the SABRE catalyst directly. As SABRE-Relay, like PHIP-X, depends on transferring proton magnetization, direct polarization of heteronuclei (analogous to SABRE-SHEATH) is impossible. However, the polarization of the exchanging OH group allows the polarization of heteronuclei either spontaneously (i.e., by free evolution), or with the help of RF-SOT. So far, SABRE-Relay has achieved 2.6% ^1H ,²²⁵ 2.3% ^{29}Si ,⁷⁹ 1.1% ^{13}C ,^{225,226} 0.2% ^{19}F ,²²⁵ and 0.04% ^{31}P ²²⁵ polarization levels. These NMR signal enhancements are generally lower than typical values achieved by conventional SABRE because the relayed polarization is derived from a finite carrier




Figure 12. Demonstration of the SABRE-Relay effect. (a) Depiction of the SABRE-Relay method. (b) SABRE-Relay can allow quantification of isomer ratios of fructose in a single scan ^{13}C measurement or (c) quantification of diastereomer ratios of the natural product (–)-carveol from a single scan ^1H measurement. Details for (b): $^{13}\text{C}\{^1\text{H}\}$ NMR spectra (right) acquired for 40 mM of D-fructose (natural ^{13}C abundance) with 23.8 mM benzyl- d_7 -amine and 4.8 mM of $[\text{Ir}(\text{Cl})(\text{COD})(\text{SIMes-}d_{22})]$ (where COD is *cis,cis*-cyclooctadiene and SIMes is 1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene) in a 0.65 mL $\text{DCM-}d_2$:DMF (1.6:1) mixture measured at 9.4 T. The bottom spectrum shows the result of a thermally polarized signal averaging over 1024 scans (approximately 17 h), and the middle spectrum represents the single scan SABRE-Relay hyperpolarization measurement recorded after shaking the sample with pH_2 at 6.5 mT. Details for (c): Exemplar single scan thermally polarized (above) and ^1H SABRE-Relay hyperpolarized (lower) ^1H NMR spectra for a sample of $[\text{Ir}(\text{Cl})(\text{COD})(\text{IMes})]$ (5 mM), NH_3 (30 mM), (–)-carveol (25 mM) and pH_2 (3 bar) in $\text{DCM-}d_2$ (0.6 mL). The resonance labels in red and blue correspond to the two diastereomers. The hyperpolarized NMR spectrum is recorded immediately after shaking the sample for 10 s with fresh pH_2 at 6.5 mT. (a, c) Adapted from Alshehri et al.²²⁸ Copyright 2023 Royal Society of Chemistry. This publication is licensed under CC BY 3.0. (b) Adapted from Richardson et al.²²⁶ Copyright 2019 Royal Society of Chemistry. This publication is licensed under CC BY 3.0.

polarization affected by spin relaxation during the chemical exchange. Current studies have optimized factors such as amine type and concentration ratios to increase the target polarization.^{225,227,228}

A major limitation of PHIP-X and SABRE-Relay is that they cannot be performed in alcohol or aqueous solvents as their exchangeable protons will compete with the ones of the target molecule. Accordingly, they are commonly performed in dry dichloromethane or chloroform, which may pose a significant challenge for insoluble substrates in these media. SABRE-Relay has already shown potential in molecular analysis. It can enhance the NMR signals of sugars²²⁶ and natural products²²⁸ at concentrations as low as tens of micromolar with a single NMR scan. Notably, it can give single-scan quantification of isomeric ratios for OH-containing molecules, such as α and β forms of glucose and fructose (Figure 12b)²²⁶ or diastereomers of natural oils like (–)-carveol (Figure 12c).²²⁸ Thanks to the continuous nature of polarization production in SABRE, relayed polarization transfer is expected to become better understood, improved to increase polarization levels, and applied to an ever-increasing scope of target molecules in the years ahead.

Compared to direct pH_2 addition (PHIP-X), using reversible exchange (SABRE) to polarize the transfer agent has the advantage that the transfer agent can be continuously repolarized; a disadvantage is that the polarization yield is usually lower. Despite recent progress,^{222,227,230} the detailed and quantitative description of chemical exchange and the spread of polarization within the target is still not fully understood.

2.8. Perspectives: PHIP in Enzymatically Catalyzed Reactions. The use of pH_2 and hydrogen–deuterium scrambling has been explored in a non-NMR context for several decades to study kinetics and intermediates of enzymes thanks to the slow conversion of pH_2 to oH_2 in pure water of about tens to hundreds of minutes.¹³⁸ Such scrambling can thereby help to understand the chemisorption and exchange process by forming HD based on a hydrogen and deuterium source.²³¹ Detection of oH_2 formation after supplying pH_2 yields information on the splitting and recombination of H_2 .²³² This information can then give a clearer kinetic picture of hydrogen-activating enzymes.²³³ The particular focus, therefore, is on hydrogenases,^{231–234} an important class of enzymes involved in the hydrogen activation process of, e.g., anaerobic organisms.^{235–238} Hydrogenases promise to be blueprints for eco-friendly catalysts for hydrogen activation on the way to produce green energy.²³⁹ Therefore, understanding the detailed hydrogen activation of hydrogenases could facilitate the design of potent eco-friendly catalysts. Three different types of hydrogenases are currently known: [FeFe]-hydrogenases, [NiFe]-hydrogenases, and the [Fe]-hydrogenase.²⁴⁰ While reaction intermediates of the first two hydrogenases could be well characterized by available methodologies such as EPR, X-ray diffraction, and IR, investigating the [Fe]-hydrogenase has proven to be more challenging. This is because the iron center of the [Fe]-hydrogenase is Fe^{II+} encapsulated in a guanylylpyridinol (FeGP) and remains in a diamagnetic state through the whole catalytic cycle (Figure 13a). Under catalytic conditions, methenyl-tetrahydromethanopterin (methenyl- H_4MPT^+) is bound by the protein, bringing together FeGP and methenyl- H_4MPT^+ . This is the active site that heterolytically cleaves molecular hydrogen into a proton and a hydride, whereby the hydride is stereospecific-




Figure 13. Active site of the mono iron hydrogenase and the hydrogenation reaction. (a) The active site of the [Fe]-hydrogenase, including the iron guanylyl pyridinol and methenyl- H_4MPT^+ . (b) Hydride transfer to the Hpro-R position of the substrate forming methylene- H_4MPT . Adapted from Kaltschnee et al.¹⁵³ Copyright 2024 Springer Nature. This publication is licensed under CC-BY 4.0.

cally transferred to the Hpro-R position of the methylene carbon of methylene- H_4MPT ²⁴¹ (Figure 13b). So far, computational models have predicted several iron–hydrogen species in the catalytic cycle, none of which could be experimentally verified. The use of pH_2 in hyperpolarization experiments has recently changed this.¹⁵³

When pH_2 was supplied to an aqueous buffer containing [Fe]-hydrogenase and methenyl- H_4MPT^+ , the appearance of a hyperpolarized PNL (section 2.2) as well as HD and HDO NEPTUN PHIP signals (section 2.4) was observed (Figure 14a).¹⁵³ The former state is created when pH_2 reversibly binds to the enzyme. In addition, hyperpolarized HD signals, first observed with iridium catalysts in the context of the NEPTUN effect,²⁴² were also observed in the presence of hydrogenase when the buffer was partially deuterated. This finding further indicates an isotope exchange with the solvent. In addition, rapid exchange between an enzyme-bound ensemble of hydrogen where both hydrogens are distinguishable, and a state where both hydrogens are indistinguishable on the NMR time scale, needs to occur. An estimate for the lifetime range of 1–100 μs was found, and chemical shifts and 1H – 1H J -coupling constants between these hydrogens were estimated. Optimized structural models based on the X-ray crystal structure of the hydrogenase allowed for the computation of 1H chemical shifts and 1H – 1H J -coupling constants for the hydrogen atoms within the active site and correlation with the experimental observations. Considering all this, an intermediate could be identified that had only been predicted previously (Figure 14b).²⁴³ The optimized structure reveals the presence of an iron hydride and the involvement of the oxopyridine site during the activation process. The determined intermediate supports the previously theoretically predicted process during




Figure 14. Using PHIP to reveal hitherto unobservable intermediates of [Fe] hydrogenase. (a) Hyperpolarized signals were observed when $p\text{H}_2$ was supplied to an enzymatic solution of the hydrogenase containing methenyl- H_4MPT^+ . A PNL effect and oneH-PHIP hyperpolarized HD signals demonstrate the activation of $p\text{H}_2$ by hydrogenase. (b) The reaction intermediate was determined using observed NMR parameters and analysis of chemical exchange, and it was derived from a crystal structure quantum mechanical optimization of the structure. Adapted from Kaltschnee et al.¹⁵³ Copyright 2024 Springer Nature. This publication is licensed under CC-BY 4.0.

which an oxo-pyridine moiety serves as a base for hydrogen activation during the catalytic process.

With this demonstration, the use of $p\text{H}_2$ -enhanced magnetic resonance has evolved into a tool to study the biochemistry and catalysis of hydrogenases and to investigate so far undetectable reaction intermediates of this important class of enzymes. It is envisioned that the concept can be used to study additional hydrogen-activating enzymes, such as the other two types of hydrogenases and potentially nitrogenases.

3. OUTLOOK

Almost 40 years after its first discovery, PHIP remains a source for ever-surprising novel applications and unique revelations, some of which are discussed in this review. In the wake of biomedical hyperpolarized MRI, recent discoveries of efficient ways to hyperpolarize ^{13}C -labeled pyruvate led to its *in vivo* imaging demonstrations.^{86,87,89,90} The high levels of polarization achieved using PHIP led to observations of effects such as radio amplification by stimulated emission of radiation (RASER) and spin diffusion in the solid state that attracted attention beyond the chemical community.^{59,244–250} Continued exploration and development of PHIP has and will likely continue to yield tools and techniques that complement more traditional methods, which have their shortcomings.

One promising direction lies in the continued development of catalysts and reaction conditions that enable PHIP with catalytic systems previously considered incompatible with PHIP. The ability to perform PHIP in these settings can lead to more sustainable and versatile applications. Hence, the field of applied catalysis will be enriched among others through synergetic development with PHIP as it was when the homogeneous catalysts were tuned to reach exclusive *trans*

hydrogenation for a direct hyperpolarization of fumarate,⁴⁵ or when PHIP was demonstrated using homogeneous,^{17–19} heterogeneous,^{15,43,44,251–253} and metal-free^{150,151,182} catalytic systems. Contrast agents produced by heterogeneous PHIP are currently progressing toward *in vivo* lung imaging.^{48,49}

It is worth highlighting the innovative combination of hyperpolarization for boosting NMR sensitivity while maintaining its quantitative nature. Tessari and others proposed two compelling approaches to utilize SABRE in analytical chemistry. In one method, analytes are introduced and hyperpolarized via SABRE, enabling concentration evaluations with near-nanomolar sensitivity.^{254–258} This demonstrates the potential of SABRE as a powerful quantitative tool. The second approach focuses on hyperpolarizing the hydride signals of the catalyst, exploiting the chemical shifts induced by the coordination of trace analytes to the Ir-complex.^{105,12,259–263} This method transforms the hyperpolarized hydrides into dynamic probes for chemical analysis, opening avenues for studying even minute concentrations of analytes with precision.

Furthermore, integrating PHIP with advanced NMR techniques, such as ultrafast sequences and ultralow and high-field instrumentation, may allow for detecting and studying transient intermediates and reaction dynamics with unprecedented sensitivity, and spectral and temporal resolution.^{264–268} This can significantly enhance our understanding of complex chemical processes, lead to the discovery of new reaction pathways, and uncover details of catalytic mechanisms.

All discussed here was made possible by utilizing the correlation of spins in $p\text{H}_2$ and its effects on the nuclear spin polarization of other interacting neighboring atomic nuclei. Observing nonequilibrium polarization with NMR provides chemical resolution, enabling chemical analysis on an atomic level. NMR, however, is relatively slow, although, in some cases, like in photo-PHIP, the time resolution was boosted while preserving the chemical resolution. For example, a similar boost in time resolution was achieved when photo-induced radical pairs could be generated with a short UV irradiation, resulting in time-resolved chemically induced dynamic nuclear polarization.²⁶⁹ Many methods discussed here are still under development and require significant expertise that currently limits their wider use. However, it is worth being aware of such effects, as, for example, they could provide invaluable information regarding the mobility of H_2 on catalytic centers that are inaccessible to other methods, as was exemplified here with hydrogenase studies.

AUTHOR INFORMATION

Corresponding Authors

Andrey N. Pravdivtsev – Department Section Biomedical Imaging, Molecular Imaging North Competence Center (MOIN CC), Department of Radiology and Neuroradiology, University Medical Center Kiel, Kiel University, 24118 Kiel, Germany; orcid.org/0000-0002-8763-617X; Email: andrey.pravdivtsev@rad.uni-kiel.de

Vladimir V. Zhivonitko – NMR Research Unit, University of Oulu, Oulu 90014, Finland; orcid.org/0000-0003-2919-8690; Email: vladimir.zhivonitko@oulu.fi

Authors

Ben J. Tickner – Centre for Hyperpolarization in Magnetic Resonance (CHyM), Department of Chemistry, University of York, Heslington YO10 5NY, United Kingdom

Stefan Glöggler – Max-Planck-Institute for Multidisciplinary Sciences, 37077 Göttingen, Germany; Center for Biostructural Imaging of Neurodegeneration (BIN), 37075 Göttingen, Germany; Advanced Imaging Research Center, The University of Texas Southwestern Medical Center, Dallas, Texas 75390, United States

Jan-Bernd Hövener – Department Section Biomedical Imaging, Molecular Imaging North Competence Center (MOIN CC), Department of Radiology and Neuroradiology, University Medical Center Kiel, Kiel University, 24118 Kiel, Germany; orcid.org/0000-0001-7255-7252

Gerd Buntkowsky – Eduard-Zintl-Institut für Anorganische und Physikalische Chemie, Technische Universität Darmstadt, D-64287 Darmstadt, Germany; orcid.org/0000-0003-1304-9762

Simon B. Duckett – Centre for Hyperpolarization in Magnetic Resonance (CHyM), Department of Chemistry, University of York, Heslington YO10 5NY, United Kingdom; orcid.org/0000-0002-9788-6615

Clifford R. Bowers – Department of Chemistry and National High Magnetic Field Laboratory, University of Florida, Gainesville, Florida 32611, United States

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acscatal.4c07870>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

ANP and J-BH acknowledge support from German Federal Ministry of Education and Research (BMBF) within the framework of the e:Med research and funding concept (01ZX1915C, 03WIR6208A hyperquant), DFG (555951950, 527469039, 469366436, HO-4602/2-2, HO-4602/3, HO-4602/4, EXC2167, FOR5042, TRR287). MOIN CC was founded by a grant from the European Regional Development Fund (ERDF) and the Zukunftsprogramm Wirtschaft of Schleswig-Holstein (Project no. 122-09-053). SBD and BJT acknowledge the UK Research and Innovation (UKRI) under the UK government's Horizon Europe funding guarantee (grant number EP/X023672/1). GB gratefully acknowledges financial support from the DFG (BU 911/22-2). CRB acknowledges NSF grants CHE-2108306, CBET-1933723, and the National High Magnetic Field Laboratory User Collaborative Grants Program, which is supported by NSF DMR-2128556 and the State of Florida. SG acknowledges funding from the DFG (SFB1633 project B03) and the Max Planck Society. VVZ is grateful for the support from the Research Council of Finland (grant number 362959) and the University of Oulu (Kvantum Institute).

REFERENCES

- (1) Luchinat, E.; Barbieri, L.; Cremonini, M.; Banci, L. Protein in-cell NMR spectroscopy at 1.2 GHz. *J. Biomol. NMR* **2021**, *75* (2), 97–107.
- (2) Kovacs, H.; Moskau, D.; Spraul, M. Cryogenically cooled probes—a leap in NMR technology. *Prog. Nucl. Magn. Reson. Spectrosc.* **2005**, *46* (2), 131–155.
- (3) Tickner, B. J.; Singh, K.; Zhivonitko, V. V.; Telkki, V.-V. Ultrafast Nuclear Magnetic Resonance as a Tool to Detect Rapid Chemical Change in Solution. *ACS Phys. Chem. Au* **2024**, *4* (5), 453–463.
- (4) Eills, J.; Budker, D.; Cavagnero, S.; Chekmenev, E. Y.; Elliott, S. J.; Jannin, S.; Lesage, A.; Matysik, J.; Meersmann, T.; Prisner, T.; Reimer, J. A.; Yang, H.; Koptug, I. V. Spin Hyperpolarization in Modern Magnetic Resonance. *Chem. Rev.* **2023**, *123* (4), 1417–1551.
- (5) Aime, S.; Dastrù, W.; Gobetto, R.; Santelia, D.; Viale, A. *Agents for Polarization Enhancement in MRI*; Semmler, W., Schwaiger, M., Eds.; Springer: Berlin, Heidelberg, 2008; pp 247–272. DOI: [10.1007/978-3-540-72718-7_12](https://doi.org/10.1007/978-3-540-72718-7_12).
- (6) Nelson, S. J.; Kurhanewicz, J.; Vigneron, D. B.; Larson, P. E. Z.; Harzstark, A. L.; Ferrone, M.; van Criekinge, M.; Chang, J. W.; Bok, R.; Park, I.; Reed, G.; Carvajal, L.; Small, E. J.; Munster, P.; Weinberg, V. K.; Ardenkjaer-Larsen, J. H.; Chen, A. P.; Hurd, R. E.; Odegardstuen, L.-I.; Robb, F. J.; Tropp, J.; Murray, J. A. Metabolic Imaging of Patients with Prostate Cancer Using Hyperpolarized [$1-^{13}\text{C}$]Pyruvate. *Sci. Transl. Med.* **2013**, *5* (198), 198ra108.
- (7) Gutte, H.; Hansen, A. E.; Larsen, M. M. E.; Rahbek, S.; Henriksen, S. T.; Johannesen, H. H.; Ardenkjaer-Larsen, J.; Kristensen, A. T.; Højgaard, L.; Kjær, A. Simultaneous Hyperpolarized ^{13}C -Pyruvate MRI and ^{18}F -FDG PET (HyperPET) in 10 Dogs with Cancer. *J. Nucl. Med.* **2015**, *56* (11), 1786–1792.
- (8) Chung, B. T.; Chen, H.-Y.; Gordon, J.; Mammoli, D.; Sriram, R.; Autry, A. W.; Le Page, L. M.; Chaumeil, M.; Shin, P.; Slater, J.; Tan, C. T.; Suszczynski, C.; Chang, S.; Li, Y.; Bok, R. A.; Ronen, S. M.; EZ Larson, P.; Kurhanewicz, J.; Vigneron, D. B. First Hyperpolarized [$2-^{13}\text{C}$]Pyruvate MR Studies of Human Brain Metabolism. *J. Magn. Reson.* **2019**, *309*, 106617.
- (9) Qi, C.; Mankinen, O.; Telkki, V.-V.; Hilty, C. Measuring Protein-Ligand Binding by Hyperpolarized Ultrafast NMR. *J. Am. Chem. Soc.* **2024**, *146* (8), 5063–5066.
- (10) Epasto, L. M.; Che, K.; Kozak, F.; Selimovic, A.; Kadeřávek, P.; Kurzbach, D. Toward protein NMR at physiological concentrations by hyperpolarized water—Finding and mapping uncharted conformational spaces. *Sci. Adv.* **2022**, *8* (31), No. eabq5179.
- (11) Pham, P.; Biswas, O.; Hilty, C. Parahydrogen Polarization in Reverse Micelles and Application to Sensing of Protein-Ligand Binding. *J. Am. Chem. Soc.* **2024**, *146*, 34274.
- (12) Hermkens, N. K. J.; Eshuis, N.; van Weerdenburg, B. J. A.; Feiters, M. C.; Rutjes, F. P. J. T.; Wijmenga, S. S.; Tessari, M. NMR-Based Chemosensing via p- H_2 Hyperpolarization: Application to Natural Extracts. *Anal. Chem.* **2016**, *88* (6), 3406–3412.
- (13) Dey, A.; Charrier, B.; Martineau, E.; Deborde, C.; Gandriaux, E.; Moing, A.; Jacob, D.; Eshchenko, D.; Schnell, M.; Melzi, R.; Kurzbach, D.; Ceillier, M.; Chappuis, Q.; Cousin, S. F.; Kempf, J. G.; Jannin, S.; Dumez, J.-N.; Giraudeau, P. Hyperpolarized NMR Metabolomics at Natural ^{13}C Abundance. *Anal. Chem.* **2020**, *92* (22), 14867–14871.
- (14) Duckett, S. B.; Sleight, C. J. Applications of the parahydrogen phenomenon: A chemical perspective. *Prog. Nucl. Magn. Reson. Spectrosc.* **1999**, *34* (1), 71–92.
- (15) Kovtunov, K. V.; Zhivonitko, V. V.; Skovpin, I. V.; Barskiy, D. A.; Koptug, I. V. Parahydrogen-Induced Polarization in Heterogeneous Catalytic Processes. *Topics in Current Chemistry*; Springer: Berlin, Heidelberg, 2012; Vol. 338, pp 123–180. DOI: [10.1007/128_2012_371](https://doi.org/10.1007/128_2012_371).
- (16) Tickner, B. J.; Zhivonitko, V. V. Advancing homogeneous catalysis for parahydrogen-derived hyperpolarisation and its NMR applications. *Chem. Sci.* **2022**, *13* (17), 4670–4696.
- (17) Bowers, C. R.; Weitekamp, D. P. Transformation of Symmetrization Order to Nuclear-Spin Magnetization by Chemical Reaction and Nuclear Magnetic Resonance. *Phys. Rev. Lett.* **1986**, *57* (21), 2645–2648.
- (18) Bowers, C. R.; Weitekamp, D. P. Parahydrogen and synthesis allow dramatically enhanced nuclear alignment. *J. Am. Chem. Soc.* **1987**, *109* (18), 5541–5542.

- (19) Eisenschmid, T. C.; Kirss, R. U.; Deutsch, P. P.; Hommeltoft, S. I.; Eisenberg, R.; Bargon, J.; Lawler, R. G.; Balch, A. L. Para hydrogen induced polarization in hydrogenation reactions. *J. Am. Chem. Soc.* **1987**, *109* (26), 8089–8091.
- (20) Green, R. A.; Adams, R. W.; Duckett, S. B.; Mewis, R. E.; Williamson, D. C.; Green, G. G. R. The theory and practice of hyperpolarization in magnetic resonance using parahydrogen. *Prog. Nucl. Magn. Reson. Spectrosc.* **2012**, *67*, 1–48.
- (21) Feng, B.; Coffey, A. M.; Colon, R. D.; Chekmenev, E. Y.; Waddell, K. W. A pulsed injection parahydrogen generator and techniques for quantifying enrichment. *J. Magn. Reson.* **2012**, *214*, 258–262.
- (22) Hövener, J.-B.; Bär, S.; Leupold, J.; Jenne, K.; Leibfritz, D.; Hennig, J.; Duckett, S. B.; von Elverfeldt, D. A continuous-flow, high-throughput, high-pressure parahydrogen converter for hyperpolarization in a clinical setting. *NMR Biomed.* **2013**, *26* (2), 124–131.
- (23) Du, Y.; Zhou, R.; Ferrer, M.-J.; Chen, M.; Graham, J.; Malphurs, B.; Labbe, G.; Huang, W.; Bowers, C. R. An Inexpensive Apparatus for up to 97% Continuous-Flow Parahydrogen Enrichment Using Liquid Helium. *J. Magn. Reson.* **2020**, *321*, 106869.
- (24) Nantogma, S.; Joalland, B.; Wilkens, K.; Chekmenev, E. Y. Clinical-Scale Production of Nearly Pure (>98.5%) Parahydrogen and Quantification by Benchtop NMR Spectroscopy. *Anal. Chem.* **2021**, *93*, 3594.
- (25) Jeong, K.; Min, S.; Chae, H.; Namgoong, S. K. Detecting low concentrations of unsaturated C—C bonds by parahydrogen-induced polarization using an efficient home-built parahydrogen generator. *Magn. Reson. Chem.* **2018**, *56* (11), 1089–1093.
- (26) Ellermann, F.; Pravdivtsev, A.; Hövener, J.-B. Open-source, partially 3D-printed, high-pressure (50-bar) liquid-nitrogen-cooled parahydrogen generator. *Magn. Reson.* **2021**, *2* (1), 49–62.
- (27) Chapman, B.; Joalland, B.; Meersman, C.; Ettedgui, J.; Swenson, R. E.; Krishna, M. C.; Nikolaou, P.; Kovtunov, K. V.; Salnikov, O. G.; Koptug, I. V.; Gemeinhardt, M. E.; Goodson, B. M.; Shchepin, R. V.; Chekmenev, E. Y. Low-Cost High-Pressure Clinical-Scale 50% Parahydrogen Generator Using Liquid Nitrogen at 77 K. *Anal. Chem.* **2021**, *93*, 8476.
- (28) Natterer, J.; Bargon, J. Parahydrogen induced polarization. *Prog. Nucl. Magn. Reson. Spectrosc.* **1997**, *31* (4), 293–315.
- (29) Buntkowsky, G.; Theiss, F.; Lins, J.; Miloslavina, Y. A.; Wienands, L.; Kiryutin, A.; Yurkovskaya, A. Recent advances in the application of parahydrogen in catalysis and biochemistry. *RSC Adv.* **2022**, *12* (20), 12477–12506.
- (30) Buljubasich, L.; Franzoni, M. B.; Münnemann, K. Parahydrogen Induced Polarization by Homogeneous Catalysis: Theory and Applications. *Top. Curr. Chem.* **2013**, *338*, 33–74.
- (31) Bowers, C. R. Sensitivity Enhancement Utilizing Parahydrogen. In *eMagRes*; Harris, R. K., Wasylishen, R. L., Eds.; Wiley, 2007. DOI: 10.1002/9780470034590.emrstm0489
- (32) Pravica, M. G.; Weitekamp, D. P. Net NMR alignment by adiabatic transport of parahydrogen addition products to high magnetic field. *Chem. Phys. Lett.* **1988**, *145* (4), 255–258.
- (33) Imamoto, T. New P-chirogenic diphosphines and their use in catalytic asymmetric reactions. *Pure Appl. Chem.* **2001**, *73* (2), 373–376.
- (34) Bhattacharya, P.; Chekmenev, E. Y.; Perman, W. H.; Harris, K. C.; Lin, A. P.; Norton, V. A.; Tan, C. T.; Ross, B. D.; Weitekamp, D. P. Towards hyperpolarized ^{13}C -succinate imaging of brain cancer. *J. Magn. Reson.* **2007**, *186* (1), 150–155.
- (35) Hövener, J.-B.; Chekmenev, E. Y.; Harris, K. C.; Perman, W. H.; Tran, T. T.; Ross, B. D.; Bhattacharya, P. Quality assurance of PASADENA hyperpolarization for ^{13}C biomolecules. *Magn. Reson. Mater. Phys.* **2009**, *22* (2), 123–134.
- (36) Huynh, M. T.; Kovacs, Z. Homogeneous Catalysts for Hydrogenative PHIP Used in Biomedical Applications. *Anal. Sens.* **2025**, *5*, No. e202400044.
- (37) Klages, S.; Permin, A. B.; Petrosyan, V. S.; Bargon, J. Parahydrogen induced polarization in the Pt(0)-mediated hydrogenation of alkynes and alkenes. *J. Org. Chem.* **1997**, *545*–546, 201–205.
- (38) Sulman, E.; Deibele, C.; Bargon, J. Study of homogenous hydrogenation of acetylene compounds with para-hydrogen and Pd(0) and Pt(0) complexes by in situ NMR spectroscopy. *React. Kinet. Catal. Lett.* **1999**, *67* (1), 117–122.
- (39) Tokmic, K.; Fout, A. R. Alkyne Semihydrogenation with a Well-Defined Nonclassical Co-H₂ Catalyst: A H₂ Spin on Isomerization and E-Selectivity. *J. Am. Chem. Soc.* **2016**, *138* (41), 13700–13705.
- (40) Zhivonitko, V. V.; Skovpin, I. V.; Szeto, K. C.; Taoufik, M.; Koptug, I. V. Parahydrogen-Induced Polarization Study of the Silica-Supported Vanadium Oxo Organometallic Catalyst. *J. Phys. Chem. C* **2018**, *122* (9), 4891–4900.
- (41) Muhammad, S. R.; Greer, R. B.; Ramirez, S. B.; Goodson, B. M.; Fout, A. R. Cobalt-Catalyzed Hyperpolarization of Structurally Intact Olefins. *ACS Catal.* **2021**, *11*, 2011–2020.
- (42) Najera, D. C.; Fout, A. R. Iron-Catalyzed Parahydrogen Induced Polarization. *J. Am. Chem. Soc.* **2023**, *145* (38), 21086–21095.
- (43) Koptug, I. V.; Kovtunov, K. V.; Burt, S. R.; Anwar, M. S.; Hilty, C.; Han, S.-I.; Pines, A.; Sagdeev, R. Z. para-Hydrogen-Induced Polarization in Heterogeneous Hydrogenation Reactions. *J. Am. Chem. Soc.* **2007**, *129* (17), 5580–5586.
- (44) Kovtunov, K. V.; Salnikov, O. G.; Zhivonitko, V. V.; Skovpin, I. V.; Bukhtiyarov, V. I.; Koptug, I. V. Catalysis and Nuclear Magnetic Resonance Signal Enhancement with Parahydrogen. *Top. Catal.* **2016**, *59* (19–20), 1686–1699.
- (45) Ripka, B.; Eills, J.; Kouřilová, H.; Leutzsch, M.; Levitt, M. H.; Münnemann, K. Hyperpolarized fumarate via parahydrogen. *Chem. Commun.* **2018**, *54* (86), 12246–12249.
- (46) Stewart, N. J.; Nakano, H.; Sugai, S.; Tomohiro, M.; Kase, Y.; Uchio, Y.; Yamaguchi, T.; Matsuo, Y.; Naganuma, T.; Takeda, N.; Nishimura, I.; Hirata, H.; Hashimoto, T.; Matsumoto, S. Hyperpolarized ^{13}C Magnetic Resonance Imaging of Fumarate Metabolism by Parahydrogen-induced Polarization: A Proof-of-Concept in vivo Study. *ChemPhysChem* **2021**, *22* (10), 915–923.
- (47) Gallagher, F. A.; Kettunen, M. I.; Hu, D.-E.; Jensen, P. R.; Zandt, R.; Karlsson, M.; Gisselsson, A.; Nelson, S. K.; Witney, T. H.; Bohndiek, S. E.; Hansson, G.; Peetersen, T.; Lerche, M. H.; Brindle, K. M. Production of hyperpolarized [1,4- $^{13}\text{C}_2$]malate from [1,4- $^{13}\text{C}_2$]-fumarate is a marker of cell necrosis and treatment response in tumors. *Proc. Nat. Acad. Sci.* **2009**, *106* (47), 19801–19806.
- (48) Ariyasingha, N. M.; Samoilenko, A.; Birchall, J. R.; Chowdhury, M. R. H.; Salnikov, O. G.; Kovtunova, L. M.; Bukhtiyarov, V. I.; Zhu, D. C.; Qian, C.; Bradley, M.; Gelovani, J. G.; Koptug, I. V.; Goodson, B. M.; Chekmenev, E. Y. Ultra-Low-Cost Disposable Hand-Held Clinical-Scale Propane Gas Hyperpolarizer for Pulmonary Magnetic Resonance Imaging Sensing. *ACS Sens.* **2023**, *8* (10), 3845–3854.
- (49) Chowdhury, M. R. H.; Oladun, C.; Ariyasingha, N. M.; Samoilenko, A.; Bawardi, T.; Burueva, D. B.; Salnikov, O. G.; Kovtunova, L. M.; Bukhtiyarov, V. I.; Shi, Z.; Luo, K.; Tan, S.; Gelovani, J. G.; Koptug, I. V.; Goodson, B. M.; Chekmenev, E. Y. Rapid lung ventilation MRI using parahydrogen-induced polarization of propane gas. *Analyst* **2024**, *149* (24), 5832–5842.
- (50) Zhou, R.; Zhao, E. W.; Cheng, W.; Neal, L. M.; Zheng, H.; Quinones, R. E.; Hagelin-Weaver, H. E.; Bowers, C. R. Parahydrogen-Induced Polarization by Pairwise Replacement Catalysis on Pt and Ir Nanoparticles. *J. Am. Chem. Soc.* **2015**, *137* (5), 1938–1946.
- (51) Zhou, R.; Cheng, W.; Neal, L. M.; Zhao, E. W.; Ludden, K.; Hagelin-Weaver, H. E.; Bowers, C. R. Parahydrogen enhanced NMR reveals correlations in selective hydrogenation of triple bonds over supported Pt catalyst. *Phys. Chem. Chem. Phys.* **2015**, *17* (39), 26121–26129.
- (52) Salnikov, O. G.; Pokochueva, E. V.; Burueva, D. B.; Kovtunova, L. M.; Kovtunov, K. V.; Koptug, I. V. Parahydrogen-Induced Polarization in Gas-Phase Heterogeneous Hydrogenation of Epoxides. *J. Phys. Chem. C* **2023**, *127* (49), 23634–23644.
- (53) Gyton, M. R.; Royle, C. G.; Beaumont, S. K.; Duckett, S. B.; Weller, A. S. Mechanistic Insights into Molecular Crystalline

Organometallic Heterogeneous Catalysis through Parahydrogen-Based Nuclear Magnetic Resonance Studies. *J. Am. Chem. Soc.* **2023**, *145* (4), 2619–2629.

(54) Jang, M.; Duckett, S. B.; Eisenberg, R. Definitive Evidence for a Pairwise Addition of Hydrogen to a Platinum Bis(phosphine) Complex Using Parahydrogen-Induced Polarization. *Organometallics* **1996**, *15* (13), 2863–2865.

(55) Adams, R. W.; Aguilar, J. A.; Atkinson, K. D.; Cowley, M. J.; Elliott, P. I. P.; Duckett, S. B.; Green, G. G. R.; Khazal, I. G.; López-Serrano, J.; Williamson, D. C. Reversible Interactions with parahydrogen Enhance NMR Sensitivity by Polarization Transfer. *Science* **2009**, *323* (5922), 1708–1711.

(56) Adams, R. W.; Duckett, S. B.; Green, R. A.; Williamson, D. C.; Green, G. G. R. A theoretical basis for spontaneous polarization transfer in non-hydrogenative parahydrogen-induced polarization. *J. Chem. Phys.* **2009**, *131* (19), 194505.

(57) Hövener, J.-B.; Schwaderlapp, N.; Lickert, T.; Duckett, S. B.; Mewis, R. E.; Highton, L. A. R.; Kenny, S. M.; Green, G. G. R.; Leibfritz, D.; Korvink, J. G.; Hennig, J.; von Elverfeldt, D. A hyperpolarized equilibrium for magnetic resonance. *Nat. Commun.* **2013**, *4*, 2946.

(58) Pravdivtsev, A. N.; Yurkovskaya, A. V.; Vieth, H.-M.; Ivanov, K. L. RF-SABRE: A Way to Continuous Spin Hyperpolarization at High Magnetic Fields. *J. Phys. Chem. B* **2015**, *119* (43), 13619–13629.

(59) Suefke, M.; Lehmkuhl, S.; Liebisch, A.; Blümich, B.; Appelt, S. Para-hydrogen raser delivers sub-millihertz resolution in nuclear magnetic resonance. *Nat. Phys.* **2017**, *13* (6), 568–572.

(60) Pravdivtsev, A. N.; Yurkovskaya, A. V.; Vieth, H.-M.; Ivanov, K. L.; Kaptein, R. Level Anti-Crossings are a Key Factor for Understanding para-Hydrogen-Induced Hyperpolarization in SABRE Experiments. *ChemPhysChem* **2013**, *14* (14), 3327–3331.

(61) Theis, T.; Truong, M.; Coffey, A. M.; Chekmenev, E. Y.; Warren, W. S. LIGHT-SABRE enables efficient in-magnet catalytic hyperpolarization. *J. Magn. Reson.* **2014**, *248*, 23–26.

(62) Theis, T.; Truong, M. L.; Coffey, A. M.; Shchepin, R. V.; Waddell, K. W.; Shi, F.; Goodson, B. M.; Warren, W. S.; Chekmenev, E. Y. Microtesla SABRE Enables 10% Nitrogen-15 Nuclear Spin Polarization. *J. Am. Chem. Soc.* **2015**, *137* (4), 1404–1407.

(63) Pravdivtsev, A. N.; Yurkovskaya, A. V.; Zimmermann, H.; Vieth, H.-M.; Ivanov, K. L. Transfer of SABRE-derived hyperpolarization to spin-1/2 heteronuclei. *RSC Adv.* **2015**, *5* (78), 63615–63623.

(64) Ivanov, K. L.; Pravdivtsev, A. N.; Yurkovskaya, A. V.; Vieth, H.-M.; Kaptein, R. The role of level anti-crossings in nuclear spin hyperpolarization. *Prog. Nucl. Magn. Reson. Spectrosc.* **2014**, *81*, 1–36.

(65) Soon, P. C.; Xu, X.; Zhang, B.; Gruppi, F.; Canary, J. W.; Jerschow, A. Hyperpolarization of amino acid precursors to neurotransmitters with parahydrogen induced polarization. *Chem. Commun.* **2013**, *49* (46), 5304–5306.

(66) Reineri, F.; Boi, T.; Aime, S. ParaHydrogen Induced Polarization of ^{13}C carboxylate resonance in acetate and pyruvate. *Nat. Commun.* **2015**, *6*, 5858.

(67) Cavallari, E.; Carrera, C.; Di Matteo, G.; Bondar, O.; Aime, S.; Reineri, F. In-vitro NMR Studies of Prostate Tumor Cell Metabolism by Means of Hyperpolarized $[1-^{13}\text{C}]\text{Pyruvate}$ Obtained Using the PHIP-SAH Method. *Front. Oncol.* **2020**, *10*, 497.

(68) Carrera, C.; Cavallari, E.; Digilio, G.; Bondar, O.; Aime, S.; Reineri, F. ParaHydrogen polarized ethyl- $[1-^{13}\text{C}]\text{pyruvate}$ in water, a key substrate for fostering the PHIP-SAH approach to metabolic imaging. *ChemPhysChem* **2021**, *22*, 1042.

(69) Brahm, A.; Pravdivtsev, A. N.; Stamp, T.; Ellermann, F.; Sönnichsen, F.; Hövener, J.-B.; Herges, R. Synthesis of ^{13}C and ^2H Labeled Vinyl Pyruvate and Hyperpolarization of Pyruvate. *Chem.—Eur. J.* **2022**, *28*, No. e202201210.

(70) Ding, Y.; Korchak, S.; Mamone, S.; Jagtap, A. P.; Stevanato, G.; Sternkopf, S.; Moll, D.; Schroeder, H.; Becker, S.; Fischer, A.; Gerhardt, E.; Outeiro, T. F.; Opazo, F.; Griesinger, C.; Glöggler, S. Rapidly Signal-enhanced Metabolites for Atomic Scale Monitoring of

Living Cells with Magnetic Resonance. *Chem. Methods* **2022**, *2*, No. e202200023.

(71) Brahm, A.; Pravdivtsev, A. N.; Thorns, L.; Sönnichsen, F. D.; Hövener, J.-B.; Herges, R. Exceptionally Mild and High-Yielding Synthesis of Vinyl Esters of Alpha-Ketocarboxylic Acids, Including Vinyl Pyruvate, for Parahydrogen-Enhanced Metabolic Spectroscopy and Imaging. *J. Org. Chem.* **2023**, *88* (21), 15018–15028.

(72) Mei, R.; Fries, L. M.; Hune, T. L. K.; Santi, M. D.; Rodriguez, G. G.; Sternkopf, S.; Glöggler, S. Hyperpolarization of ^{15}N -Pyridinium by Using Parahydrogen Enables Access to Reactive Oxygen Sensors and Pilot In Vivo Studies. *Angew. Chem., Int. Ed.* **2024**, *63*, No. e202403144.

(73) Fernandez Diaz-Rullo, F.; Zamberlan, F.; Mewis, R. E.; Fekete, M.; Broche, L.; Cheyne, L. A.; Dall'Angelo, S.; Duckett, S. B.; Dawson, D.; Zanda, M. Synthesis and hyperpolarisation of eNOS substrates for quantification of NO production by ^1H NMR spectroscopy. *Bioorg. Med. Chem.* **2017**, *25* (10), 2730–2742.

(74) Tickner, B. J.; Borozdina, Y.; Duckett, S. B.; Angelovski, G. Exploring the hyperpolarisation of EGTA-based ligands using SABRE. *Dalton Trans.* **2021**, *50* (7), 2448–2461.

(75) Barskiy, D. A.; Knecht, S.; Yurkovskaya, A. V.; Ivanov, K. L. SABRE: Chemical kinetics and spin dynamics of the formation of hyperpolarization. *Prog. Nucl. Magn. Reson. Spectrosc.* **2019**, *114–115*, 33–70.

(76) Iali, W.; Roy, S. S.; Tickner, B. J.; Ahwal, F.; Kennerley, A. J.; Duckett, S. B. Hyperpolarising Pyruvate through Signal Amplification by Reversible Exchange (SABRE). *Angew. Chem., Int. Ed.* **2019**, *58* (30), 10271–10275.

(77) Zhivonitko, V. V.; Skovpin, I. V.; Koptuyg, I. V. Strong ^{31}P nuclear spin hyperpolarization produced via reversible chemical interaction with parahydrogen. *Chem. Commun.* **2015**, *51* (13), 2506–2509.

(78) Shchepin, R. V.; Barskiy, D. A.; Coffey, A. M.; Goodson, B. M.; Chekmenev, E. Y. NMR Signal Amplification by Reversible Exchange of Sulfur-Heterocyclic Compounds Found In Petroleum. *ChemistrySelect* **2016**, *1* (10), 2552–2555.

(79) Rayner, P. J.; Richardson, P. M.; Duckett, S. B. The Detection and Reactivity of Silanols and Silanes Using Hyperpolarized ^{29}Si Nuclear Magnetic Resonance. *Angew. Chem., Int. Ed.* **2020**, *59* (7), 2710–2714.

(80) Bhattacharya, P.; Harris, K.; Lin, A. P.; Mansson, M.; Norton, V. A.; Perman, W. H.; Weitekamp, D. P.; Ross, B. D. Ultra-fast three dimensional imaging of hyperpolarized ^{13}C in vivo. *Magn. Reson. Mater. Phys.* **2005**, *18* (5), 245–256.

(81) Olsson, L. E.; Chai, C.-M.; Axelsson, O.; Karlsson, M.; Golman, K.; Petersson, J. S. MR coronary angiography in pigs with intraarterial injections of a hyperpolarized ^{13}C substance. *Magn. Reson. Med.* **2006**, *55* (4), 731–737.

(82) Golman, K.; Axelsson, O.; Jóhannesson, H.; Månsson, S.; Olofsson, C.; Petersson, J. S. Parahydrogen-induced polarization in imaging: Subsecond ^{13}C angiography. *Magn. Reson. Med.* **2001**, *46* (1), 1–5.

(83) Goldman, M.; Jóhannesson, H.; Axelsson, O.; Karlsson, M. Design and implementation of ^{13}C hyper polarization from parahydrogen, for new MRI contrast agents. *C. R. Chim.* **2006**, *9* (3), 357–363.

(84) Hövener, J.-B.; Chekmenev, E. Y.; Harris, K. C.; Perman, W. H.; Robertson, L. W.; Ross, B. D.; Bhattacharya, P. PASADENA hyperpolarization of ^{13}C biomolecules: equipment design and installation. *Magn. Reson. Mater. Phys.* **2009**, *22* (2), 111–121.

(85) Gierse, M.; Nagel, L.; Keim, M.; Lucas, S.; Speidel, T.; Lobmeyer, T.; Winter, G.; Josten, F.; Karaali, S.; Fellermann, M.; Scheuer, J.; Müller, C.; van Heijster, F.; Skinner, J.; Löffler, J.; Parker, A.; Handwerker, J.; Marshall, A.; Salhov, A.; El-Kassem, B.; Vassiliou, C.; Blanchard, J. W.; Picazo-Frutos, R.; Eills, J.; Barth, H.; Jelezko, F.; Rasche, V.; Schilling, F.; Schwartz, I.; Knecht, S. Parahydrogen-Polarized Fumarate for Preclinical in Vivo Metabolic Magnetic Resonance Imaging. *J. Am. Chem. Soc.* **2023**, *145* (10), 5960–5969.

- (86) Nagel, L.; Gierse, M.; Gottwald, W.; Ahmadova, Z.; Grashe, M.; Wolff, P.; Josten, F.; Karaali, S.; Müller, C. A.; Lucas, S.; Scheuer, J.; Müller, C.; Blanchard, J.; Topping, G. J.; Wendlinger, A.; Setzer, N.; Sühnel, S.; Handwerker, J.; Vassiliou, C.; van Heijster, F. H. A.; Knecht, S.; Keim, M.; Schilling, F.; Schwartz, I. Parahydrogen-Polarized $[1-^{13}\text{C}]$ Pyruvate for Reliable and Fast Preclinical Metabolic Magnetic Resonance Imaging. *Adv. Sci.* **2023**, *10* (30), 2303441.
- (87) Hune, T. L. K.; Mamone, S.; Schmidt, A. B.; Mahú, I.; D'Apolito, N.; Wiedermann, D.; Brüning, J.; Glöggler, S. Hyperpolarized Multi-organ Spectroscopy of Liver and Brain Using $1-^{13}\text{C}$ -Pyruvate Enhanced via Parahydrogen. *Appl. Magn. Reson.* **2023**, *54* (11), 1283–1295.
- (88) Cavallari, E.; Carrera, C.; Sorge, M.; Bonne, G.; Muchir, A.; Aime, S.; Reineri, F. The ^{13}C hyperpolarized pyruvate generated by ParaHydrogen detects the response of the heart to altered metabolism in real time. *Sci. Rep.* **2018**, *8* (1), 8366.
- (89) de Maissin, H.; Groß, P. R.; Mohiuddin, O.; Weigt, M.; Nagel, L.; Herzog, M.; Wang, Z.; Willing, R.; Reichardt, W.; Pichotka, M.; Heß, L.; Reinheckel, T.; Jessen, H. J.; Zeiser, R.; Bock, M.; von Elverfeldt, D.; Zaitsev, M.; Korchak, S.; Glöggler, S.; Hövener, J.-B.; Chekmenev, E. Y.; Schilling, F.; Knecht, S.; Schmidt, A. B. In Vivo Metabolic Imaging of $[1-^{13}\text{C}]$ Pyruvate- d_3 Hyperpolarized By Reversible Exchange With Parahydrogen. *Angew. Chem., Int. Ed.* **2023**, *62*, No. e202306654.
- (90) MacCulloch, K.; Browning, A.; Guarín Bedoya, D. O.; McBride, S. J.; Abdulmojeed, M. B.; Dedesma, C.; Goodson, B. M.; Rosen, M. S.; Chekmenev, E. Y.; Yen, Y.-F.; TomHon, P.; Theis, T. Facile hyperpolarization chemistry for molecular imaging and metabolic tracking of $[1-^{13}\text{C}]$ pyruvate in vivo. *J. Magn. Reson. Open* **2023**, *16*–17, 100129.
- (91) Wolber, J.; Cherubini, A.; Dzik-Jurasz, A. S. K.; Leach, M. O.; Bifone, A. Spin-lattice relaxation of laser-polarized xenon in human blood. *Proc. Natl. Acad. Sci. U.S.A.* **1999**, *96* (7), 3664–3669.
- (92) Schröder, L.; Lowery, T. J.; Hilty, C.; Wemmer, D. E.; Pines, A. Molecular Imaging Using a Targeted Magnetic Resonance Hyperpolarized Biosensor. *Science* **2006**, *314* (5798), 446–449.
- (93) Khan, A. S.; Harvey, R. L.; Birchall, J. R.; Irwin, R. K.; Nikolaou, P.; Schrank, G.; Emami, K.; Dummer, A.; Barlow, M. J.; Goodson, B. M.; Chekmenev, E. Y. Enabling Clinical Technologies for Hyperpolarized ^{129}Xe Magnetic Resonance Imaging and Spectroscopy. *Angew. Chem., Int. Ed.* **2021**, *133* (41), 22298–22319.
- (94) Hövener, J.-B.; Pravdivtsev, A. N.; Kidd, B.; Bowers, C. R.; Glöggler, S.; Kovtunov, K. V.; Plaumann, M.; Katz-Brull, R.; Buckenmaier, K.; Jerschow, A.; Reineri, F.; Theis, T.; Shchepin, R. V.; Wagner, S.; Bhattacharya, P.; Zacharias, N. M.; Chekmenev, E. Y. Parahydrogen-Based Hyperpolarization for Biomedicine. *Angew. Chem., Int. Ed.* **2018**, *57* (35), 11140–11162.
- (95) Kovtunov, K. V.; Pokochueva, E.; Salnikov, O.; Cousin, S.; Kurzbach, D.; Vuichoud, B.; Jannin, S.; Chekmenev, E.; Goodson, B.; Barskiy, D.; Koptuyg, I. Hyperpolarized NMR Spectroscopy: d-DNP, PHIP, and SABRE Techniques. *Chem.—Asian J.* **2018**, *13* (15), 1857–1871.
- (96) Pravdivtsev, A. N.; Buntkowsky, G.; Duckett, S. B.; Koptuyg, I. V.; Hövener, J.-B. Parahydrogen-Induced Polarization of Amino Acids. *Angew. Chem., Int. Ed.* **2021**, *60*, 23496.
- (97) Reineri, F.; Cavallari, E.; Carrera, C.; Aime, S. Hydrogenative-PHIP polarized metabolites for biological studies. *Magn. Reson. Mater. Phys.* **2021**, *34* (1), 25–47.
- (98) Schmidt, A. B.; Bowers, C. R.; Buckenmaier, K.; Chekmenev, E. Y.; de Maissin, H.; Eills, J.; Ellermann, F.; Glöggler, S.; Gordon, J. W.; Knecht, S.; Koptuyg, I. V.; Kuhn, J.; Pravdivtsev, A. N.; Reineri, F.; Theis, T.; Them, K.; Hövener, J.-B. Instrumentation for Hydrogenative Parahydrogen-Based Hyperpolarization Techniques. *Anal. Chem.* **2022**, *94* (1), 479–502.
- (99) Dagys, L.; Ripka, B.; Leutzsch, M.; Moustafa, G. A. I.; Eills, J.; Colell, J. F. P.; Levitt, M. H. Geminal parahydrogen-induced polarization: accumulating long-lived singlet order on methylene proton pairs. *Magn. Reson.* **2020**, *1* (2), 175–186.
- (100) Pravdivtsev, A. N.; Brahms, A.; Kienitz, S.; Sönnichsen, F. D.; Hövener, J.-B.; Herges, R. Catalytic Hydrogenation of Trivinyl Orthoacetate: Mechanisms Elucidated by Parahydrogen Induced Polarization. *ChemPhysChem* **2021**, *22* (4), 370–377.
- (101) Duckett, S. B.; Newell, C. L.; Eisenberg, R. More than INEPT: parahydrogen and INEPT + give unprecedented resonance enhancement to carbon-13 by direct proton polarization transfer. *J. Am. Chem. Soc.* **1993**, *115* (3), 1156–1157.
- (102) Pravdivtsev, A. N. SABRE Hyperpolarization of Bipyridine Stabilized Ir-Complex at High, Low and Ultralow Magnetic Fields. *Z. Phys. Chem.* **2017**, *231* (3), 497–511.
- (103) Robinson, A. D.; Hill-Casey, F.; Duckett, S. B.; Halse, M. E. Quantitative reaction monitoring using parahydrogen-enhanced benchtop NMR spectroscopy. *Phys. Chem. Chem. Phys.* **2024**, *26* (19), 14317–14328.
- (104) Majchrzak, R. M.; Norcott, P. L. Iridium Pyridylpyrrolides Hyperpolarised by Para-Hydrogen. *Chem.—Eur. J.* **2024**, *30*, No. e202400472.
- (105) Sellies, L.; Reile, I.; Aspers, R. L. E. G.; Feiters, M. C.; Rutjes, F. P. J. T.; Tessari, M. Parahydrogen induced hyperpolarization provides a tool for NMR metabolomics at nanomolar concentrations. *Chem. Commun.* **2019**, *55* (50), 7235–7238.
- (106) Skovpin, I. V.; Zhivonitko, V. V.; Koptuyg, I. V. Parahydrogen-Induced Polarization in Heterogeneous Hydrogenations over Silica-Immobilized Rh Complexes. *Appl. Magn. Reson.* **2011**, *41* (2), 393–410.
- (107) Harthun, A.; Giernoth, R.; Elsevier, C. J.; Bargon, J. Rhodium- and palladium-catalysed proton exchange in styrene detected in situ by para-hydrogen induced polarization. *Chem. Commun.* **1996**, No. 21, 2483–2484.
- (108) Permin, A. B.; Eisenberg, R. One-Hydrogen Polarization in Hydroformylation Promoted by Platinum-Tin and Iridium Carbonyl Complexes: A New Type of Parahydrogen-Induced Effect. *J. Am. Chem. Soc.* **2002**, *124* (42), 12406–12407.
- (109) Zhivonitko, V. V.; Skovpin, I. V.; Crespo-Quesada, M.; Kiwi-Minsker, L.; Koptuyg, I. V. Acetylene Oligomerization over Pd Nanoparticles with Controlled Shape: A Parahydrogen-Induced Polarization Study. *J. Phys. Chem. C* **2016**, *120* (9), 4945–4953.
- (110) Emonds, M.; Schikowski, D.; Klankermayer, J.; Schleker, P. P. M. Non-Pairwise Interactions in Parahydrogen Experiments: Nuclear Exchange of Single Protons Enables Bulk Water Hyperpolarization. *ChemPhysChem* **2018**, *19* (20), 2614–2620.
- (111) Procacci, B.; Duckett, S. B.; George, M. W.; Hanson-Heine, M. W. D.; Horvath, R.; Perutz, R. N.; Sun, X.-Z.; Vuong, K. Q.; Welch, J. A. Competing Pathways in the Photochemistry of $\text{Ru}(\text{H})_2(\text{CO})(\text{PPh}_3)_3$. *Organometallics* **2018**, *37* (6), 855–868.
- (112) Halse, M. E.; Procacci, B.; Perutz, R. N.; Duckett, S. Towards measuring reactivity on micro-to-millisecond timescales with laser pump, NMR probe spectroscopy. *Faraday Discuss.* **2019**, *220*, 28–44.
- (113) Halse, M. E.; Procacci, B.; Henshaw, S.-L.; Perutz, R. N.; Duckett, S. B. Coherent evolution of parahydrogen induced polarisation using laser pump, NMR probe spectroscopy: Theoretical framework and experimental observation. *J. Magn. Reson.* **2017**, *278*, 25–38.
- (114) Torres, O.; Procacci, B.; Halse, M. E.; Adams, R. W.; Blazina, D.; Duckett, S. B.; Eguillor, B.; Green, R. A.; Perutz, R. N.; Williamson, D. C. Photochemical Pump and NMR Probe: Chemically Created NMR Coherence on a Microsecond Time Scale. *J. Am. Chem. Soc.* **2014**, *136* (28), 10124–10131.
- (115) Procacci, B.; Aguiar, P. M.; Halse, M. E.; Perutz, R. N.; Duckett, S. B. Photochemical pump and NMR probe to monitor the formation and kinetics of hyperpolarized metal dihydrides. *Chem. Sci.* **2016**, *7* (12), 7087–7093.
- (116) Adams, R. W.; John, R. O.; Blazina, D.; Eguillor, B.; Cockett, M. C. R.; Dunne, J. P.; López-Serrano, J.; Duckett, S. B. Contrasting Photochemical and Thermal Catalysis by Ruthenium Arsine Complexes Revealed by Parahydrogen Enhanced NMR Spectroscopy. *Eur. J. Inorg. Chem.* **2022**, *2022* (5), No. e202100991.

- (117) Blazina, D.; Duckett, S. B.; Halstead, T. K.; Kozak, C. M.; Taylor, R. J. K.; Anwar, M. S.; Jones, J. A.; Carteret, H. A. Generation and interrogation of a pure nuclear spin state by parahydrogen-enhanced NMR spectroscopy: a defined initial state for quantum computation. *Magn. Reson. Chem.* **2005**, *43* (3), 200–208.
- (118) Pravdivtsev, A. N.; Ivanov, K. L.; Yurkovskaya, A. V.; Vieth, H.-M.; Sagdeev, R. Z. Spontaneous transfer of para-hydrogen induced polarization to ^{13}C spins in symmetric molecules. *Dokl. Phys. Chem.* **2015**, *464* (2), 247–250.
- (119) Berner, S.; Schmidt, A. B.; Zimmermann, M.; Pravdivtsev, A. N.; Glögger, S.; Hennig, J.; von Elverfeldt, D.; Hövener, J.-B. SAMBADENA Hyperpolarization of ^{13}C -Succinate in an MRI: Singlet-Triplet Mixing Causes Polarization Loss. *ChemistryOpen* **2019**, *8* (6), 728–736.
- (120) Hasnip, S.; Duckett, S. B.; Taylor, D. R.; Taylor, M. J. Parahydrogen enhanced NMR studies on thermally and photochemically generated products from $[\text{IrH}_3(\text{CO})(\text{PPh}_3)_2]$. *Chem. Commun.* **1998**, No. 8, 923–924.
- (121) Hasnip, S. K.; Colebrooke, S. A.; Sleight, C. J.; Duckett, S. B.; Taylor, D. R.; Barlow, G. K.; Taylor, M. J. Activation of H_2 by halocarbonyl bis-phosphine and bis-arsine iridium(I) complexes. The use of parahydrogen induced polarisation to detect species present at low concentration and investigate their reactivity. *J. Chem. Soc., Dalton Trans.* **2002**, No. 5, 743–751.
- (122) Natterer, J.; Schedletsky, O.; Barkemeyer, J.; Bargon, J.; Glaser, S. J. Investigating Catalytic Processes with Parahydrogen: Evolution of Zero-Quantum Coherence in AA'X Spin Systems. *J. Magn. Reson.* **1998**, *133* (1), 92–97.
- (123) Nasibulov, E. A.; Pravdivtsev, A. N.; Yurkovskaya, A. V.; Lukzen, N. N.; Vieth, H.-M.; Ivanov, K. L. Analysis of Nutation Patterns in Fourier-Transform NMR of Non-Thermally Polarized Multispin Systems. *Z. Phys. Chem.* **2013**, *227* (6–7), 929–953.
- (124) Goldman, M.; Jóhannesson, H.; Axelsson, O.; Karlsson, M. Hyperpolarization of ^{13}C through order transfer from parahydrogen: A new contrast agent for MRI. *Magn. Reson. Imag.* **2005**, *23* (2), 153–157.
- (125) Pravdivtsev, A. N.; Yurkovskaya, A. V.; Petrov, P. A.; Vieth, H.-M. Coherent evolution of singlet spin states in PHOTO-PHIP and M2S experiments. *Phys. Chem. Chem. Phys.* **2017**, *19* (38), 25961–25969.
- (126) Anwar, M. S.; Blazina, D.; Carteret, H. A.; Duckett, S. B.; Halstead, T. K.; Jones, J. A.; Kozak, C. M.; Taylor, R. J. K. Preparing High Purity Initial States for Nuclear Magnetic Resonance Quantum Computing. *Phys. Rev. Lett.* **2004**, *93* (4), 040501.
- (127) Kiryutin, A. S.; Yurkovskaya, A. V.; Lukzen, N. N.; Vieth, H.-M.; Ivanov, K. L. Exploiting adiabatically switched RF-field for manipulating spin hyperpolarization induced by parahydrogen. *J. Chem. Phys.* **2015**, *143* (23), 234203.
- (128) Brown, E. E.; Mandzhieva, I.; TomHon, P. M.; Theis, T.; Castellano, F. N. Triplet Photosensitized para-Hydrogen Induced Polarization. *ACS Cent. Sci.* **2022**, *8* (11), 1548–1556.
- (129) Kiryutin, A. S.; Kozinenko, V. P.; Yurkovskaya, A. V. Photo-SABRE: Nuclear Spin Hyperpolarization of cis-trans Photoswitchable Molecules by Parahydrogen*. *ChemPhotoChem.* **2024**, *8* (1), No. e202300151.
- (130) Brown, E. E.; Harrison, R. J.; Havrylyuk, D.; McBride, S. J.; Glaser, E. C.; Castellano, F. N.; Theis, T. Photo-ejected ligands hyperpolarized by parahydrogen in reversible exchange. *Chem. Commun.* **2025**, published online. DOI: 10.1039/D4CC06807F
- (131) Zhivonitko, V. V.; Sorochkina, K.; Chernichenko, K.; Kótai, B.; Földes, T.; Pápai, L.; Telkki, V.-V.; Repo, T.; Koptyug, I. Nuclear spin hyperpolarization with ansa-aminoboranes: a metal-free perspective for parahydrogen-induced polarization. *Phys. Chem. Chem. Phys.* **2016**, *18* (40), 27784–27795.
- (132) Kiryutin, A. S.; Sauer, G.; Yurkovskaya, A. V.; Limbach, H.-H.; Ivanov, K. L.; Buntkowsky, G. Parahydrogen Allows Ultrasensitive Indirect NMR Detection of Catalytic Hydrogen Complexes. *J. Phys. Chem. C* **2017**, *121* (18), 9879–9888.
- (133) Aguilar, J. A.; Elliott, P. I. P.; López-Serrano, J.; Adams, R. W.; Duckett, S. B. Only para-hydrogen spectroscopy (OPSY), a technique for the selective observation of para-hydrogen enhanced NMR signals. *Chem. Commun.* **2007**, 0 (11), 1183–1185.
- (134) Aguilar, J. A.; Adams, R. W.; Duckett, S. B.; Green, G. G. R.; Kandiah, R. Selective detection of hyperpolarized NMR signals derived from para-hydrogen using the Only Para-hydrogen Spectroscopy (OPSY) approach. *J. Magn. Reson.* **2011**, *208* (1), 49–57.
- (135) Pravdivtsev, A. N.; Kozinenko, V. P.; Hövener, J.-B. Only Para-Hydrogen Spectroscopy (OPSY) Revisited: In-Phase Spectra for Chemical Analysis and Imaging. *J. Phys. Chem. A* **2018**, *122* (45), 8948–8956.
- (136) Pravdivtsev, A. N.; Sönnichsen, F.; Hövener, J.-B. OnlyParahydrogen Spectroscopy (OPSY) pulse sequences - One does not fit all. *J. Magn. Reson.* **2018**, *297*, 86–95.
- (137) Buntkowsky, G.; Bargon, J.; Limbach, H.-H. A Dynamic Model of Reaction Pathway Effects on Parahydrogen-Induced Nuclear Spin Polarization. *J. Am. Chem. Soc.* **1996**, *118* (36), 8677–8683.
- (138) Schmidt, A. B.; Worner, J.; Pravdivtsev, A.; Knecht, S.; Scherer, H.; Weber, S.; Hennig, J.; von Elverfeldt, D.; Hövener, J.-B. Lifetime of Parahydrogen in Aqueous Solutions and Human Blood. *ChemPhysChem* **2019**, *20* (19), 2408–2412.
- (139) Markelov, D. A.; Kozinenko, V. P.; Knecht, S.; Kiryutin, A. S.; Yurkovskaya, A. V.; Ivanov, K. L. Singlet to triplet conversion in molecular hydrogen and its role in parahydrogen induced polarization. *Phys. Chem. Chem. Phys.* **2021**, *23* (37), 20936–20944.
- (140) Knecht, S.; Hadjiali, S.; Barskiy, D. A.; Pines, A.; Sauer, G.; Kiryutin, A. S.; Ivanov, K. L.; Yurkovskaya, A. V.; Buntkowsky, G. Indirect Detection of Short-Lived Hydride Intermediates of Iridium N-Heterocyclic Carbene Complexes via Chemical Exchange Saturation Transfer Spectroscopy. *J. Phys. Chem. C* **2019**, *123* (26), 16288–16293.
- (141) Ward, K. M.; Aletras, A. H.; Balaban, R. S. A New Class of Contrast Agents for MRI Based on Proton Chemical Exchange Dependent Saturation Transfer (CEST). *J. Magn. Reson.* **2000**, *143* (1), 79–87.
- (142) Johnson, A.; Royle, C. G.; Brodie, C. N.; Martínez-Martínez, A. J.; Duckett, S. B.; Weller, A. S. η^2 -Alkene Complexes of $[\text{Rh}(\text{PONOP-}^i\text{Pr})(\text{L})]^+$ Cations (L = COD, NBD, Ethene). Intramolecular Alkene-Assisted Hydrogenation and Dihydrogen Complex $[\text{Rh}(\text{PONOP-}^i\text{Pr})(\eta\text{-H}_2)]^+$. *Inorg. Chem.* **2021**, *60* (18), 13903–13912.
- (143) Kireev, N. V.; Kiryutin, A. S.; Pavlov, A. A.; Yurkovskaya, A. V.; Musina, E. I.; Karasik, A. A.; Shubina, E. S.; Ivanov, K. L.; Belkova, N. V. Nickel(II) Dihydrogen and Hydride Complexes as the Intermediates of H_2 Heterolytic Splitting by Nickel Diazadiphosphacyclooctane Complexes. *Eur. J. Inorg. Chem.* **2021**, 2021 (41), 4265–4272.
- (144) Bernatowicz, P.; Ratajczyk, T.; Szymański, S. Non-uniform orientation of H_2 molecules in a magnetic field as a critical condition for the appearance of partially negative NMR lines of o- H_2 in hyperpolarization experiments using p- H_2 . *J. Chem. Phys.* **2023**, *159* (12), 124304.
- (145) Czarnota, M.; Mames, A.; Pietrzak, M.; Jopa, S.; Theiß, F.; Buntkowsky, G.; Ratajczyk, T. A Straightforward Method for the Generation of Hyperpolarized Orthohydrogen with a Partially Negative Line. *Angew. Chem., Int. Ed.* **2024**, *63* (12), No. e202309188.
- (146) Alam, M. S.; Li, X.; Brittin, D. O.; Islam, S.; Deria, P.; Chekmenev, E. Y.; Goodson, B. M. Anomalous Large Antiphase Signals from Hyperpolarized Orthohydrogen Using a MOF-Based SABRE Catalyst. *Angew. Chem., Int. Ed.* **2023**, *62* (8), No. e202213581.
- (147) Fleckenstein, M.; Herr, K.; Theiß, F.; Knecht, S.; Wienands, L.; Brodrecht, M.; Reggelen, M.; Buntkowsky, G. A disintegrin derivative as a case study for PHIP labeling of disulfide bridged biomolecules. *Sci. Rep.* **2022**, *12* (1), 2337.
- (148) Townrow, O. P. E.; Duckett, S. B.; Weller, A. S.; Goicoechea, J. M. Zintl cluster supported low coordinate Rh(I) centers for

catalytic H/D exchange between H₂ and D₂. *Chem. Sci.* **2022**, *13* (25), 7626–7633.

(149) Vaneeckhaute, E.; De Ridder, S.; Tyburn, J.-M.; Kempf, J. G.; Taulelle, F.; Martens, J. A.; Breynaert, E. Long-Term Generation of Longitudinal Spin Order Controlled by Ammonia Ligation Enables Rapid SABRE Hyperpolarized 2D NMR. *ChemPhysChem* **2021**, *22* (12), 1170–1177.

(150) Zakharov, D. O.; Chernichenko, K.; Sorochkina, K.; Yang, S.; Telkki, V.-V.; Repo, T.; Zhivonitko, V. V. Parahydrogen-Induced Polarization in Hydrogenation Reactions Mediated by a Metal-Free Catalyst. *Chem.—Eur. J.* **2022**, *28* (8), No. e202103501.

(151) Zakharov, D. O.; Chernichenko, K.; Sorochkina, K.; Repo, T.; Zhivonitko, V. V. Parahydrogen-induced polarization study of imine hydrogenations mediated by a metal-free catalyst. *Dalton Trans.* **2022**, *51* (36), 13606–13611.

(152) Zakharov, D. O. Parahydrogen-induced polarization in H₂ activations and hydrogenations mediated by metal-free catalysts. PhD thesis, University of Oulu Graduate School, Faculty of Science, NMR Research Unit, 2023.

(153) Kaltschnee, L.; Pravdivtsev, A. N.; Gehl, M.; Huang, G.; Stoychev, G. L.; Riplinger, C.; Keitel, M.; Neese, F.; Hövener, J.-B.; Auer, A. A.; Griesinger, C.; Shima, S.; Glöggler, S. Sensitivity-enhanced magnetic resonance reveals hydrogen intermediates during active [Fe]-hydrogenase catalysis. *Nat. Catal.* **2024**, *7*, 1417–1429.

(154) Kowalewski, J.; Mäler, L. *Nuclear spin relaxation in liquids: theory, experiments, and applications*; Taylor & Francis: 2006.

(155) Zhao, E. W.; Maligal-Ganesh, R.; Du, Y.; Zhao, T. Y.; Collins, J.; Ma, T.; Zhou, L.; Goh, T.-W.; Huang, W.; Bowers, C. R. Surface-Mediated Hyperpolarization of Liquid Water from Parahydrogen. *Chem.* **2018**, *4* (6), 1387–1403.

(156) Guan, D.; Godard, C.; Polas, S. M.; Tooze, R. P.; Whitwood, A. C.; Duckett, S. B. Using para hydrogen induced polarization to study steps in the hydroformylation reaction. *Dalton Trans.* **2019**, *48* (8), 2664–2675.

(157) López-Serrano, J.; Duckett, S. B.; Aiken, S.; Almeida Leñero, K. Q.; Drent, E.; Dunne, J. P.; Konya, D.; Whitwood, A. C. A para-Hydrogen Investigation of Palladium-Catalyzed Alkyne Hydrogenation. *J. Am. Chem. Soc.* **2007**, *129* (20), 6513–6527.

(158) Guan, D.; Holmes, A. J.; López-Serrano, J.; Duckett, S. B. Following palladium catalyzed methoxycarbonylation by hyperpolarized NMR spectroscopy: a parahydrogen based investigation. *Catal. Sci. Technol.* **2017**, *7* (10), 2101–2109.

(159) López-Serrano, J.; Duckett, S. B.; Dunne, J. P.; Godard, C.; Whitwood, A. C. Palladium catalysed alkyne hydrogenation and oligomerisation: a parahydrogen based NMR investigation. *Dalton Trans.* **2008**, No. 32, 4270–4281.

(160) Boutain, M.; Duckett, S. B.; Dunne, J. P.; Godard, C.; Hernández, J. M.; Holmes, A. J.; Khazal, I. G.; López-Serrano, J. A parahydrogen based NMR study of Pt catalysed alkyne hydrogenation. *Dalton Trans.* **2010**, *39* (14), 3495–3500.

(161) Wigh Lipsø, K.; Hansen, E. S. S.; Tougaard, R. S.; Laustsen, C.; Ardenkjær-Larsen, J. H. Renal MR angiography and perfusion in the pig using hyperpolarized water. *Magn. Reson. Med.* **2017**, *78* (3), 1131–1135.

(162) Lipsø, K. W.; Hansen, E. S. S.; Tougaard, R. S.; Laustsen, C.; Ardenkjær-Larsen, J. H. Dynamic coronary MR angiography in a pig model with hyperpolarized water. *Magn. Reson. Med.* **2018**, *80* (3), 1165–1169.

(163) Kharbanda, Y.; Urbańczyk, M.; Zhivonitko, V. V.; Mailhot, S.; Kettunen, M. I.; Telkki, V.-V. Sensitive, Efficient and Portable Analysis of Molecular Exchange Processes by Hyperpolarized Ultrafast NMR. *Angew. Chem., Int. Ed.* **2022**, *61* (28), No. e202203957.

(164) Harris, T.; Szekely, O.; Frydman, L. On the Potential of Hyperpolarized Water in Biomolecular NMR Studies. *J. Phys. Chem. B* **2014**, *118* (12), 3281–3290.

(165) Stern, Q.; Milani, J.; Vuichoud, B.; Bornet, A.; Gossert, A. D.; Bodenhausen, G.; Jannin, S. Hyperpolarized Water to Study Protein-Ligand Interactions. *J. Phys. Chem. Lett.* **2015**, *6* (9), 1674–1678.

(166) Kurzbach, D.; Canet, E.; Flamm, A. G.; Jhajharia, A.; Weber, E. M. M.; Konrat, R.; Bodenhausen, G. Investigation of Intrinsically Disordered Proteins through Exchange with Hyperpolarized Water. *Angew. Chem., Int. Ed.* **2017**, *56* (1), 389–392.

(167) Hilty, C.; Kurzbach, D.; Frydman, L. Hyperpolarized water as universal sensitivity booster in biomolecular NMR. *Nat. Protoc.* **2022**, *17* (7), 1621–1657.

(168) Pradhan, N.; Hilty, C. Cross-Polarization of Insensitive Nuclei from Water Protons for Detection of Protein-Ligand Binding. *J. Am. Chem. Soc.* **2024**, *146*, 24754.

(169) Pinon, A. C.; Capozzi, A.; Ardenkjær-Larsen, J. H. Hyperpolarized water through dissolution dynamic nuclear polarization with UV-generated radicals. *Commun. Chem.* **2020**, *3* (1), 1–9.

(170) Lehmkuhl, S.; Emondts, M.; Schubert, L.; Spanning, P.; Klankermayer, J.; Blümich, B.; Schleker, P. P. M. Hyperpolarizing Water with Parahydrogen. *ChemPhysChem* **2017**, *18* (18), 2426–2429.

(171) Van Dyke, E. T.; Eills, J.; Picazo-Frutos, R.; Sheberstov, K. F.; Hu, Y.; Budker, D.; Barskiy, D. A. Relayed hyperpolarization for zero-field nuclear magnetic resonance. *Sci. Adv.* **2022**, *8* (29), No. eabp9242.

(172) Hale, W. G.; Zhao, T. Y.; Choi, D.; Ferrer, M.-J.; Song, B.; Zhao, H.; Hagelin-Weaver, H. E.; Bowers, C. R. Toward Continuous-Flow Hyperpolarisation of Metabolites via Heterogeneous Catalysis, Side-Arm-Hydrogenation, and Membrane Dissolution of Parahydrogen. *ChemPhysChem* **2021**, *22*, 822.

(173) Zhao, T. Y.; Lapak, M. P.; Behera, R.; Zhao, H.; Ferrer, M.-J.; Weaver, H. E. H.; Huang, W.; Bowers, C. R. Perpetual hyperpolarization of allyl acetate from parahydrogen and continuous flow heterogeneous hydrogenation with recycling of unreacted propargyl acetate. *J. Magn. Reson. Open* **2022**, *12*–13, 100076.

(174) Zhivonitko, V. V.; Kovtunov, K. V.; Beck, I. E.; Ayupov, A. B.; Bukhtiyarov, V. I.; Koptiyug, I. V. Role of Different Active Sites in Heterogeneous Alkene Hydrogenation on Platinum Catalysts Revealed by Means of Parahydrogen-Induced Polarization. *J. Phys. Chem. C* **2011**, *115* (27), 13386–13391.

(175) Pokochueva, E. V.; Burueva, D. B.; Salnikov, O. G.; Koptiyug, I. V. Heterogeneous Catalysis and Parahydrogen-Induced Polarization. *ChemPhysChem* **2021**, *22* (14), 1421–1440.

(176) Zhao, E. W.; Maligal-Ganesh, R.; Xiao, C.; Goh, T.-W.; Qi, Z.; Pei, Y.; Hagelin-Weaver, H. E.; Huang, W.; Bowers, C. R. Silica-Encapsulated Pt-Sn Intermetallic Nanoparticles: A Robust Catalytic Platform for Parahydrogen-Induced Polarization of Gases and Liquids. *Angew. Chem., Int. Ed.* **2017**, *56* (14), 3925–3929.

(177) Du, Y.; Behera, R.; Maligal-Ganesh, R. V.; Chen, M.; Chekmenev, E. Y.; Huang, W.; Bowers, C. R. Cyclopropane Hydrogenation vs. Isomerization over Pt and Pt-Sn Intermetallic Nanoparticle Catalysts: A Parahydrogen Spin Labelling Study. *J. Phys. Chem. C* **2020**, *124* (15), 8304–8309.

(178) Chen, M.; Bowers, C. R.; Huang, W. Silica-Encapsulated Intermetallic Nanoparticles for Highly Active and Selective Heterogeneous Catalysis. *Acc. Mater. Res.* **2021**, *2* (12), 1190–1202.

(179) Pan, M.; Pozun, Z. D.; Yu, W.-Y.; Henkelman, G.; Mullins, C. B. Structure Revealing H/D Exchange with Co-Adsorbed Hydrogen and Water on Gold. *J. Phys. Chem. Lett.* **2012**, *3* (14), 1894–1899.

(180) Norcott, P. L. Benzoquinone Enhances Hyperpolarization of Surface Alcohols with Para-Hydrogen. *J. Am. Chem. Soc.* **2023**, *145* (18), 9970–9975.

(181) Longobardi, L. E.; Russell, C. A.; Green, M.; Townsend, N. S.; Wang, K.; Holmes, A. J.; Duckett, S. B.; McGrady, J. E.; Stephan, D. W. Hydrogen Activation by an Aromatic Triphosphabenzene. *J. Am. Chem. Soc.* **2014**, *136* (38), 13453–13457.

(182) Zhivonitko, V. V.; Telkki, V.-V.; Chernichenko, K.; Repo, T.; Leskelä, M.; Sumerin, V.; Koptiyug, I. V. Tweezers for Parahydrogen: A Metal-Free Probe of Nonequilibrium Nuclear Spin States of H₂Molecules. *J. Am. Chem. Soc.* **2014**, *136* (2), 598–601.

(183) Sorochkina, K.; Zhivonitko, V. V.; Chernichenko, K.; Telkki, V.-V.; Repo, T.; Koptiyug, I. V. Spontaneous ¹⁵N Nuclear Spin

Hyperpolarization in Metal-Free Activation of Parahydrogen by Molecular Tweezers. *J. Phys. Chem. Lett.* **2018**, *9*, 903–907.

(184) Sorochkina, K.; Chernichenko, K.; Zhivonitko, V. V.; Nieger, M.; Repo, T. J. Water Reduction and Dihydrogen Addition in Aqueous Conditions With ansa-Phosphinoborane. *Chem.—Eur. J.* **2022**, *28*, No. e202201927.

(185) Konsewicz, K.; Laczkó, G.; Pápai, I.; Zhivonitko, V. V. Activation of H₂ using ansa-aminoboranes: solvent effects, dynamics, and spin hyperpolarization. *Phys. Chem. Chem. Phys.* **2024**, *26* (4), 3197–3207.

(186) Zhivonitko, V. V.; Bresien, J.; Schulz, A.; Koptug, I. V. Parahydrogen-induced polarization with a metal-free P-P biradicaloid. *Phys. Chem. Chem. Phys.* **2019**, *21* (11), 5890–5893.

(187) Zhivonitko, V. V.; Beer, H.; Zakharov, D. O.; Bresien, J.; Schulz, A. Hyperpolarization Effects in Parahydrogen Activation with Phosphonium Biradicaloids: Metal-free PHIP and SABRE. *ChemPhysChem* **2021**, *22* (9), 813–817.

(188) Zander, E.; Bresien, J.; Zhivonitko, V. V.; Fessler, J.; Villinger, A.; Michalik, D.; Schulz, A. Rational Design of Persistent Phosphorus-Centered Singlet Tetradicals and Their Use in Small-Molecule Activation. *J. Am. Chem. Soc.* **2023**, *145* (26), 14484–14497.

(189) Stephan, D. W. Frustrated Lewis Pairs. *J. Am. Chem. Soc.* **2015**, *137* (32), 10018–10032.

(190) Sumerin, V.; Schulz, F.; Atsumi, M.; Wang, C.; Nieger, M.; Leskelä, M.; Repo, T.; Pyykkö, P.; Rieger, B. Molecular Tweezers for Hydrogen: Synthesis, Characterization, and Reactivity. *J. Am. Chem. Soc.* **2008**, *130* (43), 14117–14119.

(191) Aime, S.; Gobetto, R.; Canet, D. Longitudinal Nuclear Relaxation in an A2 Spin System Initially Polarized through Parahydrogen. *J. Am. Chem. Soc.* **1998**, *120* (27), 6770–6773.

(192) Aime, S.; Dastrù, W.; Gobetto, R.; Russo, A.; Viale, A.; Canet, D. A Novel Application of para H₂: the Reversible Addition/Elimination of H₂ at a Ru₃ Cluster Revealed by the Enhanced NMR Emission Resonance from Molecular Hydrogen. *J. Phys. Chem. A* **1999**, *103* (48), 9702–9705.

(193) Konsewicz, K.; Repo, T.; Zhivonitko, V. V. Towards water-tolerant ansa-aminoboranes for parahydrogen-induced polarization and beyond. *Chem. Commun.* **2025**, published online. DOI: 10.1039/D5CC00528K

(194) Holtkamp, P.; Schwabedissen, J.; Neumann, B.; Stämmler, G.; Koptug, I.; Zhivonitko, V.; Mitzel, N. W. A Zwitterionic Phosphonium Stannate(II) via Hydrogen Splitting by a Sn/P Frustrated Lewis-Pair and Reductive Elimination. *Chem.—Eur. J.* **2020**, *26*, 17381.

(195) Korchak, S.; Mamone, S.; Glöggler, S. Over 50% ¹H and ¹³C Polarization for Generating Hyperpolarized Metabolites—A parahydrogen Approach. *ChemistryOpen* **2018**, *7* (9), 672–676.

(196) Reineri, F.; Daniele, V.; Cavallari, E.; Aime, S. Assessing the transport rate of hyperpolarized pyruvate and lactate from the intra- to the extracellular space. *NMR Biomed.* **2016**, *29* (8), 1022–1027.

(197) Eills, J.; Cavallari, E.; Kircher, R.; Matteo, G. D.; Carrera, C.; Dagys, L.; Levitt, M. H.; Ivanov, K. L.; Aime, S.; Reineri, F.; Münnemann, K.; Budker, D.; Buntkowsky, G.; Knecht, S. Singlet-Contrast Magnetic Resonance Imaging: Unlocking Hyperpolarization with Metabolism*. *Angew. Chem., Int. Ed.* **2021**, *60* (12), 6791–6798.

(198) Tickner, B. J.; Rayner, P. J.; Duckett, S. B. Using SABRE Hyperpolarized ¹³C NMR Spectroscopy to Interrogate Organic Transformations of Pyruvate. *Anal. Chem.* **2020**, *92* (13), 9095–9103.

(199) Tickner, B. J.; Platas-Iglesias, C.; Duckett, S. B.; Angelovski, G. In Situ Ternary Adduct Formation of Yttrium Polyaminocarboxylates Leads to Small Molecule Capture and Activation. *Chem.—Eur. J.* **2022**, *28* (57), No. e202201780.

(200) Tickner, B. J.; Dennington, M.; Collins, B. G.; Gater, C. A.; Tanner, T. F. N.; Whitwood, A. C.; Rayner, P. J.; Watts, D. P.; Duckett, S. B. Metal-Mediated Catalytic Polarization Transfer from para Hydrogen to 3,5-Dihalogenated Pyridines. *ACS Catal.* **2024**, *14*, 994–1004.

(201) Rayner, P. J.; Fekete, M.; Gater, C. A.; Ahwal, F.; Turner, N.; Kennerley, A. J.; Duckett, S. B. Real-Time High-Sensitivity Reaction Monitoring of Important Nitrogen-Cycle Synthons by ¹⁵N Hyperpolarized Nuclear Magnetic Resonance. *J. Am. Chem. Soc.* **2022**, *144* (19), 8756–8769.

(202) Angelovski, G.; Tickner, B. J.; Wang, G. Opportunities and challenges with hyperpolarized bioresponsive probes for functional imaging using magnetic resonance. *Nat. Chem.* **2023**, *15* (6), 755–763.

(203) Grant, A. K.; Vinogradov, E. Long-lived states in solution NMR: Theoretical examples in three- and four-spin systems. *J. Magn. Reson.* **2008**, *193* (2), 177–190.

(204) Levitt, M. H. Singlet Nuclear Magnetic Resonance. *Annu. Rev. Phys. Chem.* **2012**, *63* (1), 89–105.

(205) Zhivonitko, V. V.; Kovtunov, K. V.; Chapovsky, P. L.; Koptug, I. V. Nuclear Spin Isomers of Ethylene: Enrichment by Chemical Synthesis and Application for NMR Signal Enhancement. *Angew. Chem., Int. Ed.* **2013**, *52* (50), 13251–13255.

(206) Zhang, Y.; Soon, P. C.; Jerschow, A.; Canary, J. W. Long-Lived ¹H Nuclear Spin Singlet in Dimethyl Maleate Revealed by Addition of Thiols. *Angew. Chem., Int. Ed.* **2014**, *53* (13), 3396–3399.

(207) Chapovsky, P. L.; Zhivonitko, V. V.; Koptug, I. V. Conversion of Nuclear Spin Isomers of Ethylene. *J. Phys. Chem. A* **2013**, *117* (39), 9673–9683.

(208) Sviyazov, S. V.; Babenko, S. V.; Skovpin, I. V.; Kovtunova, L. M.; Chukanov, N. V.; Stakheev, A. Y.; Burueva, D. B.; Koptug, I. V. Manipulating stereoselectivity of parahydrogen addition to acetylene to unravel interconversion of ethylene nuclear spin isomers. *Phys. Chem. Chem. Phys.* **2024**, *26* (9), 7821–7829.

(209) Skovpin, I. V.; Sviyazov, S. V.; Burueva, D. B.; Kovtunova, L. M.; Nartova, A. V.; Kvon, R. I.; Bukhtiyarov, V. I.; Koptug, I. V. Nonequilibrium Nuclear Spin States of Ethylene during Acetylene Hydrogenation with Parahydrogen over Immobilized Iridium Complexes. *Dokl. Phys. Chem.* **2023**, *512* (2), 149–157.

(210) Pokochueva, E. V.; Svyatova, A. I.; Burueva, D. B.; Koptug, I. V. Chemistry of nuclear spin isomers of the molecules: from the past of the Universe to emerging technologies. *Russ. Chem. Bull.* **2023**, *72* (1), 1–19.

(211) Bae, J.; Zhou, Z.; Theis, T.; Warren, W. S.; Wang, Q. 15N4–1,2,4,5-tetrazines as potential molecular tags: Integrating bioorthogonal chemistry with hyperpolarization and unearthing para-N₂. *Sci. Adv.* **2018**, *4* (3), No. eaar2978.

(212) Procacci, B.; Roy, S. S.; Norcott, P.; Turner, N.; Duckett, S. B. Unlocking a diazine long-lived nuclear singlet state via photochemistry: NMR detection and lifetime of an unstabilized diazo-compound. *J. Am. Chem. Soc.* **2018**, *140* (48), 16855–16864.

(213) Landau, L. D.; Lifshitz, E. M. *Quantum Mechanics: Non-Relativistic Theory*, 3rd ed.; Pergamon Press: Oxford; New York, 1977.

(214) Babenko, S. V.; Sviyazov, S. V.; Burueva, D. B.; Koptug, I. V. Hyperpolarized long-lived spin state of methylene protons of 2-bromoethanol obtained from ethylene with non-equilibrium nuclear spin order. *J. Magn. Reson.* **2024**, *360*, 107648.

(215) Franzoni, M. B.; Buljubasich, L.; Spiess, H. W.; Münnemann, K. Long-Lived ¹H Singlet Spin States Originating from Parahydrogen in Cs-Symmetric Molecules Stored for Minutes in High Magnetic Fields. *J. Am. Chem. Soc.* **2012**, *134* (25), 10393–10396.

(216) Franzoni, M. B.; Graafen, D.; Buljubasich, L.; Schreiber, L. M.; Spiess, H. W.; Münnemann, K. Hyperpolarized ¹H long lived states originating from parahydrogen accessed by rf irradiation. *Phys. Chem. Chem. Phys.* **2013**, *15* (40), 17233–17239.

(217) Pravdivtsev, A. N.; Yurkovskaya, A. V.; Lukzen, N. N.; Vieth, H.-M.; Ivanov, K. L. Exploiting level anti-crossings (LACs) in the rotating frame for transferring spin hyperpolarization. *Phys. Chem. Chem. Phys.* **2014**, *16* (35), 18707–18719.

(218) Graafen, D.; Franzoni, M. B.; Schreiber, L. M.; Spiess, H. W.; Münnemann, K. Magnetic resonance imaging of ¹H long lived states derived from parahydrogen induced polarization in a clinical system. *J. Magn. Reson.* **2016**, *262*, 68–72.

- (219) Them, K.; Ellermann, F.; Pravdivtsev, A. N.; Salnikov, O. G.; Skovpin, I. V.; Koptuyug, I. V.; Herges, R.; Hövener, J.-B. Parahydrogen-Induced Polarization Relayed via Proton Exchange. *J. Am. Chem. Soc.* **2021**, *143* (34), 13694–13700.
- (220) Iali, W.; Rayner, P. J.; Duckett, S. B. Using parahydrogen to hyperpolarize amines, amides, carboxylic acids, alcohols, phosphates, and carbonates. *Sci. Adv.* **2018**, *4* (1), No. eaao6250.
- (221) Alciček, S.; Van Dyke, E.; Xu, J.; Pustelny, S.; Barskiy, D. A. ^{13}C and ^{15}N Benchtop NMR Detection of Metabolites via Relayed Hyperpolarization. *Chem. Methods* **2023**, *3* (7), No. e202200075.
- (222) Them, K.; Kuhn, J.; Pravdivtsev, A. N.; Hövener, J.-B. Nuclear spin polarization of lactic acid via exchange of parahydrogen-polarized protons. *Comm. Chem.* **2024**, *7*, 172.
- (223) Pravdivtsev, A. N.; Yurkovskaya, A. V.; Zimmermann, H.; Vieth, H.-M.; Ivanov, K. L. Enhancing NMR of insensitive nuclei by transfer of SABRE spin hyperpolarization. *Chem. Phys. Lett.* **2016**, *661*, 77–82.
- (224) Iali, W.; Rayner, P. J.; Alshehri, A.; Holmes, A. J.; Ruddlesden, A. J.; Duckett, S. B. Direct and indirect hyperpolarisation of amines using parahydrogen. *Chem. Sci.* **2018**, *9* (15), 3677–3684.
- (225) Rayner, P. J.; Tickner, B. J.; Iali, W.; Fekete, M.; Robinson, A. D.; Duckett, S. B. Relayed hyperpolarization from para-hydrogen improves the NMR detectability of alcohols. *Chem. Sci.* **2019**, *10* (33), 7709–7717.
- (226) Richardson, P. M.; Iali, W.; Roy, S. S.; Rayner, P. J.; Halse, M. E.; Duckett, S. B. Rapid ^{13}C NMR hyperpolarization delivered from para-hydrogen enables the low concentration detection and quantification of sugars. *Chem. Sci.* **2019**, *10* (45), 10607–10619.
- (227) Tickner, B. J.; Svensson, S. K.-M.; Vaara, J.; Duckett, S. B. Toward Optimizing and Understanding Reversible Hyperpolarization of Lactate Esters Relayed from para-Hydrogen. *J. Phys. Chem. Lett.* **2022**, *13* (29), 6859–6866.
- (228) Alshehri, A.; Tickner, B. J.; Iali, W.; Duckett, S. B. Enhancing the NMR signals of plant oil components using hyperpolarisation relayed via proton exchange. *Chem. Sci.* **2023**, *14* (36), 9843–9853.
- (229) Vaneckhaute, E.; Tyburn, J.-M.; Kilgour, D.; Kempf, J. G.; Taulelle, F.; Martens, J. A.; Breyneert, E. Hyperpolarised Magnetic Resonance of Exchangeable Protons Using Parahydrogen and Aminosilane. *J. Phys. Chem. C* **2020**, *124* (27), 14541–14549.
- (230) Knecht, S.; Barskiy, D. A.; Buntkowsky, G.; Ivanov, K. L. Theoretical Description of Hyperpolarization Formation in the SABRE-relay Method. *J. Chem. Phys.* **2020**, *153*, 164106.
- (231) Inokuchi, H. Catalytic activity of organic semiconductors and enzymes. *Discuss. Faraday Soc.* **1971**, *51*, 183.
- (232) Dementin, S.; Burlat, B.; Lacey, A. L. D.; Pardo, A.; Adryanczyk-Perrier, G.; Guigliarelli, B.; Fernandez, V. M.; Rousset, M. A Glutamate Is the Essential Proton Transfer Gate during the Catalytic Cycle of the [NiFe] Hydrogenase. *J. Biol. Chem.* **2004**, *279* (11), 10508–10513.
- (233) Yagi, T.; Tsuda, M.; Inokuchi, H. Kinetic studies on hydrogenase. Parahydrogen-Orthohydrogen Conversion and Hydrogen-Deuterium Exchange Reactions. *J. Biochem.* **1973**, *73* (5), 1069–1081.
- (234) Couper, A.; Eley, D. D.; Hayward, A. Hydrogen-activating enzymes of bacteria. *Discuss. Faraday Soc.* **1955**, *20*, 174.
- (235) Shima, S.; Thauer, R. K. A third type of hydrogenase catalyzing H_2 activation. *Chem. Record.* **2007**, *7* (1), 37–46.
- (236) Shima, S.; Pilak, O.; Vogt, S.; Schick, M.; Stagni, M. S.; Meyer-Klaucke, W.; Warkentin, E.; Thauer, R. K.; Ermler, U. The Crystal Structure of [Fe]-Hydrogenase Reveals the Geometry of the Active Site. *Science* **2008**, *321* (5888), 572–575.
- (237) Lubitz, W.; Ogata, H.; Rüdiger, O.; Reijerse, E. Hydrogenases. *Chem. Rev.* **2014**, *114* (8), 4081–4148.
- (238) Wang, C.; Lai, Z.; Huang, G.; Pan, H.-J. Current State of [Fe]-Hydrogenase and Its Biomimetic Models. *Chem.—Eur. J.* **2022**, *28* (57), No. e202201499.
- (239) Ji, H.; Wan, L.; Gao, Y.; Du, P.; Li, W.; Luo, H.; Ning, J.; Zhao, Y.; Wang, H.; Zhang, L.; Zhang, L. Hydrogenase as the basis for green hydrogen production and utilization. *J. Energy Chem.* **2023**, *85*, 348–362.
- (240) Vignais, P. M.; Billoud, B. Occurrence, Classification, and Biological Function of Hydrogenases: An Overview. *Chem. Rev.* **2007**, *107* (10), 4206–4272.
- (241) Schleucher, J.; Griesinger, C.; Schwoerer, B.; Thauer, R. K. H_2 -Forming $\text{N}^5, \text{N}^{10}$ -Methylenetetrahydromethanopterin Dehydrogenase from *Methanobacterium thermoautotrophicum* Catalyzes a Stereoselective Hydride Transfer As Determined by Two-Dimensional NMR Spectroscopy. *Biochemistry* **1994**, *33* (13), 3986–3993.
- (242) Glöggl, S.; Müller, R.; Colell, J.; Emondts, M.; Dabrowski, M.; Blümich, B.; Appelt, S. Para-hydrogen induced polarization of amino acids, peptides and deuterium-hydrogen gas. *Phys. Chem. Chem. Phys.* **2011**, *13* (30), 13759–13764.
- (243) Huang, G.; Wagner, T.; Wodrich, M. D.; Ataka, K.; Bill, E.; Ermler, U.; Hu, X.; Shima, S. The atomic-resolution crystal structure of activated [Fe]-hydrogenase. *Nat. Catal.* **2019**, *2* (6), 537–543.
- (244) Pravdivtsev, A. N.; Sönnichsen, F. D.; Hövener, J.-B. Continuous Radio Amplification by Stimulated Emission of Radiation using Parahydrogen Induced Polarization (PHIP-RASER) at 14 T. *ChemPhysChem* **2020**, *21* (7), 667–672.
- (245) Korchak, S.; Kaltschnee, L.; Dervisoglu, R.; Andreas, L.; Griesinger, C.; Glöggl, S. Spontaneous Enhancement of Magnetic Resonance Signals Using a RASER. *Angew. Chem., Int. Ed.* **2021**, *60* (38), 20984–20990.
- (246) Salnikov, O. G.; Trofimov, I. A.; Pravdivtsev, A. N.; Them, K.; Hövener, J.-B.; Chekmenev, E. Y.; Koptuyug, I. V. Through-Space Multinuclear Magnetic Resonance Signal Enhancement Induced by Parahydrogen and Radiofrequency Amplification by Stimulated Emission of Radiation. *Anal. Chem.* **2022**, *94* (43), 15010–15017.
- (247) Lehmkuhl, S.; Fleischer, S.; Lohmann, L.; Rosen, M. S.; Chekmenev, E. Y.; Adams, A.; Theis, T.; Appelt, S. RASER MRI: Magnetic resonance images formed spontaneously exploiting cooperative nonlinear interaction. *Sci. Adv.* **2022**, *8* (28), No. eabp8483.
- (248) Nelson, C.; Schmidt, A. B.; Adelabu, I.; Nantogma, S.; Kiselev, V. G.; Abdurraheem, A.; de Maissin, H.; Lehmkuhl, S.; Appelt, S.; Theis, T.; Chekmenev, E. Y. Parahydrogen-Induced Carbon-13 Radiofrequency Amplification by Stimulated Emission of Radiation. *Angew. Chem., Int. Ed.* **2023**, *62* (5), No. e202215678.
- (249) Gierse, M.; Dagys, L.; Keim, M.; Lucas, S.; Josten, F.; Plenio, M. B.; Schwartz, I.; Knecht, S.; Eills, J. Hyperpolarizing Small Molecules using Parahydrogen and Solid-State Spin Diffusion. *Angew. Chem., Int. Ed.* **2024**, *63*, No. e202319341.
- (250) Dagys, L.; Korzeczek, M. C.; Parker, A. J.; Eills, J.; Blanchard, J. W.; Bengs, C.; Levitt, M. H.; Knecht, S.; Schwartz, I.; Plenio, M. B. Robust parahydrogen-induced polarization at high concentrations. *Sci. Adv.* **2024**, *10* (30), No. eado0373.
- (251) Bouchard, L.-S.; Kovtunov, K. V.; Burt, S. R.; Anwar, M. S.; Koptuyug, I. V.; Sagdeev, R. Z.; Pines, A. Para-Hydrogen-Enhanced Hyperpolarized Gas-Phase Magnetic Resonance Imaging. *Angew. Chem., Int. Ed.* **2007**, *46* (22), 4064–4068.
- (252) Kovtunov, K. V.; Beck, I. E.; Bukhtiyarov, V. I.; Koptuyug, I. V. Observation of Parahydrogen-Induced Polarization in Heterogeneous Hydrogenation on Supported Metal Catalysts. *Angew. Chem., Int. Ed.* **2008**, *47* (8), 1492–1495.
- (253) Chen, S.; Tennakoon, A.; You, K.-E.; Paterson, A. L.; Yappert, R.; Alayoglu, S.; Fang, L.; Wu, X.; Zhao, T. Y.; Lapak, M. P.; et al. Ultrasmall amorphous zirconia nanoparticles catalyze polyolefin hydrogenolysis. *Nat. Catal.* **2023**, *6* (2), 161–173.
- (254) Eshuis, N.; Hermkens, N.; van Weerdenburg, B. J. A.; Feiters, M. C.; Rutjes, F. P. J. T.; Wijmenga, S. S.; Tessari, M. Toward Nanomolar Detection by NMR Through SABRE Hyperpolarization. *J. Am. Chem. Soc.* **2014**, *136* (7), 2695–2698.
- (255) Eshuis, N.; Aspers, R. L. E. G.; van Weerdenburg, B. J. A.; Feiters, M. C.; Rutjes, F. P. J. T.; Wijmenga, S. S.; Tessari, M. 2D NMR Trace Analysis by Continuous Hyperpolarization at High Magnetic Field. *Angew. Chem., Int. Ed.* **2015**, *54* (48), 14527–14530.

- (256) Eshuis, N.; van Weerdenburg, B. J. A.; Feiters, M. C.; Rutjes, F. P. J. T.; Wijmenga, S. S.; Tessari, M. Quantitative Trace Analysis of Complex Mixtures Using SABRE Hyperpolarization. *Angew. Chem., Int. Ed.* **2015**, *54* (5), 1481–1484.
- (257) Reile, I.; Eshuis, N.; Hermkens, N. K. J.; van Weerdenburg, B. J. A.; Feiters, M. C.; Rutjes, F. P. J. T.; Tessari, M. NMR detection in biofluid extracts at sub- μ M concentrations via para- H_2 induced hyperpolarization. *Analyst* **2016**, *141* (13), 4001–4005.
- (258) Tickner, B. J.; Komulainen, S.; Palosaari, S.; Heikkinen, J.; Lehenkari, P.; Zhivonitko, V. V.; Telkki, V.-V. Hyperpolarised NMR to aid molecular profiling of electronic cigarette aerosols. *RSC Adv.* **2022**, *12* (3), 1479–1485.
- (259) Hermkens, N. K. J.; Aspers, R. L. E. G.; Feiters, M. C.; Rutjes, F. P. J. T.; Tessari, M. Trace analysis in water-alcohol mixtures by continuous p- H_2 hyperpolarization at high magnetic field. *Magn. Reson. Chem.* **2018**, *56* (7), 633–640.
- (260) Sellies, L.; Aspers, R.; Feiters, M. C.; Rutjes, F.; Tessari, M. Para-hydrogen hyperpolarization allows direct NMR detection of α -amino acids in complex (bio)mixtures. *Angew. Chem., Int. Ed.* **2021**, *60* (52), 26954–26959.
- (261) Reimets, N.; Ausmees, K.; Vija, S.; Reile, I. Developing Analytical Applications for Parahydrogen Hyperpolarization: Urinary Elimination Pharmacokinetics of Nicotine. *Anal. Chem.* **2021**, *93* (27), 9480–9485.
- (262) Tickner, B.; John, R. O.; Roy, S. S.; Hart, S.; Whitwood, A. C.; Duckett, S. B. Using coligands to gain mechanistic insight into iridium complexes hyperpolarized with para-hydrogen. *Chem. Sci.* **2019**, *10* (20), 5235–5245.
- (263) Wood, N. J.; Brannigan, J. A.; Duckett, S. B.; Heath, S. L.; Wagstaff, J. Detection of Picomole Amounts of Biological Substrates by para-Hydrogen-Enhanced NMR Methods in Conjunction with a Suitable Receptor Complex. *J. Am. Chem. Soc.* **2007**, *129* (36), 11012–11013.
- (264) Barskiy, D. A.; Tayler, M. C. D.; Marco-Rius, I.; Kurhanewicz, J.; Vigneron, D. B.; Cikrikci, S.; Aydogdu, A.; Reh, M.; Pravdivtsev, A. N.; Hövener, J.-B.; Blanchard, J. W.; Wu, T.; Budker, D.; Pines, A. Zero-field nuclear magnetic resonance of chemically exchanging systems. *Nat. Commun.* **2019**, *10* (1), 3002.
- (265) Barskiy, D. A.; Blanchard, J. W.; Budker, D.; Eills, J.; Pustelny, S.; Sheberstov, K. F.; Tayler, M. C. D.; Trabesinger, A. H. Zero- to Ultralow-field Nuclear Magnetic Resonance. *Prog. Nucl. Magn. Reson. Spectrosc.* **2025**, *148–149*, 101558.
- (266) Myers, J. Z.; Bullinger, F.; Kempf, N.; Plaumann, M.; Ortmeier, A.; Theis, T.; Povolni, P.; Romanowski, J.; Engelmann, J.; Scheffler, K.; Hövener, J.-B.; Buckenmaier, K.; Körber, R.; Pravdivtsev, A. N. Zero to ultralow magnetic field NMR of $[1-^{13}C]$ pyruvate and $[2-^{13}C]$ pyruvate enabled by SQUID sensors and hyperpolarization. *Phys. Rev. B* **2024**, *109* (18), 184443.
- (267) Tickner, B. J.; Zhivonitko, V. V.; Telkki, V.-V. Ultrafast Laplace NMR to study metal-ligand interactions in reversible polarisation transfer from parahydrogen. *Phys. Chem. Chem. Phys.* **2021**, *23* (31), 16542–16550.
- (268) Kiryutin, A. S.; Sauer, G.; Tietze, D.; Brodrecht, M.; Knecht, S.; Yurkovskaya, A. V.; Ivanov, K. L.; Avrutina, O.; Kolmar, H.; Buntkowsky, G. Ultrafast Single-Scan 2D-NMR Spectroscopic Detection of a PHIP-Hyperpolarized Protease Inhibitor. *Chem.—Eur. J.* **2019**, *25* (16), 4025–4030.
- (269) Morozova, O. B.; Fishman, N. N.; Yurkovskaya, A. V. Indirect NMR detection of transient guanosyl radical protonation in neutral aqueous solution. *Phys. Chem. Chem. Phys.* **2017**, *19* (32), 21262–21266.